
TOPOLOGY PROCEEDINGS



Volume 2, 1977

Pages 233–241

<http://topology.auburn.edu/tp/>

TOPOLOGICAL PROPERTIES OF PRODUCT SPACES AND THE NOTION OF n -CARDINALITY

by

TEODOR C. PRZYMUSIŃSKI

Topology Proceedings

Web: <http://topology.auburn.edu/tp/>

Mail: Topology Proceedings
Department of Mathematics & Statistics
Auburn University, Alabama 36849, USA

E-mail: topolog@auburn.edu

ISSN: 0146-4124

COPYRIGHT © by Topology Proceedings. All rights reserved.

TOPOLOGICAL PROPERTIES OF PRODUCT SPACES AND THE NOTION OF n -CARDINALITY

Teodor C. Przymusiński*

Recently the author obtained several examples of product spaces with interesting topological properties. In this paper we discuss these examples and the notion of n -cardinality, which plays an important role in their construction.

1. Normality and Paracompactness of Products

Example 1. [P] For every k and m such that $k \leq m \leq \omega$ there exists a separable and first countable space X such that:

(a) X^n is paracompact (Lindelöf, subparacompact) iff $n < k$;

(b) X^n is (collectionwise) normal iff $n < m$.

Corollary 1. [P] There exists a paracompact space X such that X^2 is normal but X^2 is not paracompact.

Corollary 1 answers a question raised by E. Michael [M] and attributed by him to M. Maurice. Consistent examples of spaces with properties described in Corollary 1 have been obtained by the author [P₁] in 1973 under the assumption of $MA + \neg CH$ and later by K. Alster and P. Zenor [AZ] under the assumption of CH . On the other hand let us recall that the

* This paper was written while the author was a Visiting Assistant Professor at the University of Pittsburgh. It is a slightly expanded version of an invited talk given by the author at the March 1977 Topology Conference in Baton Rouge, Louisiana.

perfect but X^3 is not?

Our next examples give, assuming CH, a positive answer to Heath's question. Let us recall that a Lindelöf space is perfect if and only if it is hereditarily Lindelöf.

Example 3. (CH) $[P_3]$ For every $n < \omega$ there exists a separable and first countable space X such that:

- (a) X^n is hereditarily Lindelöf;
- (b) X^{n+1} is Lindelöf, but not hereditarily.

Example 4. (CH) $[P_3]$ For every $n < \omega$ there exists a separable and first countable, locally compact and locally countable space X such that:

- (a) X^n is perfectly normal;
- (b) X^{n+1} is normal, but not hereditarily.

Spaces described in the above examples cannot be both paracompact and locally compact. This follows from the fact that a locally compact and paracompact space whose square is perfect is metrizable (cf. [E]).

4. Subparacompactness of Product Spaces

In 1972, K. Alster and R. Engelking obtained the first example of a paracompact space X whose square is not subparacompact [AE]. Their example, however, was neither Lindelöf nor first countable. F. Tall asked [T] whether there exist such Lindelöf spaces and it was also unknown whether products of subparacompact spaces with metric spaces are again subparacompact. These questions are answered by our next example.

Example 5. [P] There exists a first countable and

separable Lindelöf space X and a metric separable space M such that the product space $X \times M$ is not countably θ -refinable (hence, it is not subparacompact).

Corollary 4. [P] *There exists a first countable and separable Lindelöf space Z such that Z^2 is not countably θ -refinable.*

Proof. Put $Z = X \oplus M$.

The following question remains open.

Question 2. *Does there exist a Lindelöf space X and a complete separable metric space M such that the product space $X \times M$ is not Lindelöf or-equivalently-normal?*

Such spaces exist if CH is assumed [M].

5. The Notion of n -Cardinality

In our examples an important role is played by the notion of n -cardinality, which is defined and investigated in [P₄]. The notion of n -cardinality turns out to be useful in constructions involving product spaces and seems to be a natural generalization of the notion of cardinality.

Definition 1. [P₄] Let X be an arbitrary set and let A be a subset of X^n , where n is a natural number. The n -cardinality $|A|_n$ of A (with respect to X^n) is defined by $|A|_n = \max \{|B| : B \subset A \text{ and } p_i \neq q_i, \text{ for every two distinct points } p = (p_1, \dots, p_n) \text{ and } q = (q_1, \dots, q_n) \text{ from } B \text{ and } i = 1, 2, \dots, n\}$.

The set A is n -countable (n -uncountable) if $|A|_n \leq \omega$ ($|A|_n > \omega$).

It is shown in [P₄] that $|A|_n$ is well-defined and that:

- (1) $|A|_n \leq |A|$;
- (2) $|A|_1 = |A|$;
- (3) $|A|_2 = \min \{|Y| : Y \subset X \text{ and } A \subset Y \times X \cup X \times Y\}$;
- (4) $|A|_n = \min \{|Y| : Y \subset X \text{ and } A \subset \bigcup_{i=1}^n (X^{i-1} \times Y \times X^{n-i})\}^{(2)}$.

The following Theorem generalizes a result of van Douwen.

Theorem 1. ([P₄]; cf. [vD]) *Let M be a separable complete metric space and B a Borel subset of M^m . The following conditions are equivalent:*

- (i) B is n -uncountable;
- (ii) $|B|_n = 2^\omega$;
- (iii) B contains a homeomorphic image $h(C)$ of the Cantor set C by the diagonal

$$h = \Delta_{i=1}^n h_i : C \rightarrow M^n$$

of homeomorphic embeddings $h_i : C \rightarrow M$.

The following result, which can be derived from Theorem 1, is probably most useful in applications.

Theorem 2. [P₄] *Let M be a separable, complete, and uncountable metric space. There exists a family $\{A_i\}_{i < \omega}$ of disjoint subsets A_i of M such that for every $n < \omega$, for every n -uncountable Borel subset B of M^n and for every $i < \omega$ we have $B \cap A_i^n \neq \emptyset$.*

As an application of Theorem 2, we shall give an independent proof of Corollary 2, i.e.; we shall construct a space X such that X^n is Lindelöf for all $n < \omega$, but X^ω is not normal.

(2) To be precise, equalities (3) and (4) are valid if the n -cardinality of A is infinite.

Construction: Let M be the real line R , and let A_i 's be as in Theorem 2. We can clearly assume that $R = \bigcup_{i < \omega} A_i$. Denote by τ_i the topology on R obtained by making all points from A_i isolated and put $X_i = (R, \tau_i)$ and $X = \bigoplus_{i < \omega} X_i$.

We shall show that the space X has the required properties.

A. X^n is Lindelöf, for every $n < \omega$.

The proof is by induction on $n = 0, 1, \dots$. Clearly X^0 is Lindelöf. Assume that $n > 0$, and that X^{n-1} is Lindelöf. In order to show that X^n is Lindelöf it suffices to prove that for every sequence i_1, \dots, i_n of indexes the space

$$Y = X_{i_1} \times \dots \times X_{i_n} = (R^n, \prod_{j=1}^n \tau_{i_j})$$

is Lindelöf. Choose $i < \omega$ different from all i_j 's, for $j = 1, 2, \dots, n$. Therefore for every j the topology τ_{i_j} coincides with the usual topology at points of A_i . Let \mathcal{U} be an open covering of Y . There exists a countable refinement \mathcal{V} of \mathcal{U} covering A_i^n and consisting of euclidean-open sets. The set $B = R^n \setminus \mathcal{U}$; is euclidean-closed and disjoint from A_i^n . By Theorem 2 the set B is n -countable and therefore by (4) there exists a countable subset Y of R such that

$$(5) \quad B \subset \bigcup_{j=1}^n (R^{j-1} \times Y \times R^{n-j}).$$

It follows easily from (5) and the inductive assumption that B is a Lindelöf subspace of Y and hence can be covered by a countable subcollection of \mathcal{U} .

B. X^ω is not normal.

The space X^ω contains the space $Z = \prod_{i < \omega} X_i = (R^\omega, \prod_{i < \omega} \tau_i)$ as a closed subset but the diagonal

$$\Delta = \{(x, x, \dots) \in R^\omega : x \in R\}$$

is a closed subset of Z and since $R = \bigcup_{i < \omega} A_i$ the subspace Δ is discrete. This implies that X^ω is not Lindelöf. The non-normality of X^ω follows from a theorem of K. Nagami [N] and P. Zenor [Z] stating that if X^n is Lindelöf, for every $n < \omega$ and X^ω is normal, then X^ω is Lindelöf. This completes the proof.

References

- [AE] K. Alster and R. Engelking, *Subparacompactness and product spaces*, Bull. Polon. Acad. Sci. 20 (1972), 763-767.
- [AP] K. Alster and T. Przymusiński, *Normality and Martin's Axiom*, Fundamenta Mathematicae 91 (1976), 123-131.
- [AZ] K. Alster and P. Zenor, *An example concerning the preservation of the Lindelöf property in product spaces*, Set-theoretic topology, edited by G. M. Reed, Academic Press, New York, 1976.
- [vD] E. van Douwen, *A technique for constructing honest, locally compact, submetrizable examples*, to appear.
- [E] R. Engelking, *General topology*, Polish Scientific Publishers, Warsaw, 1977.
- [H] R. W. Heath, *On p-spaces, q-spaces, n-spaces, and "s" spaces*, Proc. Auburn Topological Conference, March 1969, 123-134.
- [M] E. Michael, *Paracompactness and the Lindelöf property in finite and countable cartesian products*, Compositio Math. 23 (1971), 199-214.
- [N] K. Nagami, *A note on the normality of inverse limits and products*, Proc. International Symp. on Topology and its Appl. Herceg-Novri 1968, Beograd 1969, 261-264.
- [P] T. Przymusiński, *Normality and paracompactness in finite and countable cartesian products*, Fund. Math., to appear.
- [P₁] _____, *A Lindelöf space X such that X^2 is normal but not paracompact*, Fund. Math. 78 (1973), 291-296.

- [P₂] T. Przymusiński, *On the dimension of product spaces and an example of M. Wage*, to appear.
- [P₃] _____, *Products of perfectly normal spaces*, Fund. Math., to appear.
- [P₄] _____, *On the notion of n-cardinality*, Proc. Amer. Math. Soc., to appear.
- [RS] M. E. Rudin and M. Starbird, *Products with a metric factor*, Gen. Topology Appl. 5 (1975), 235-248.
- [T] F. D. Tall, *Problem session*, Colloq. Math. Soc. Janos Bolyai, 8, Topics in Topology, Amsterdam, North Holland, 1974.
- [W] M. Wage, *The dimension of product spaces*, to appear.
- [Z] P. Zenor, *Countable paracompactness in product spaces*, Proc. Amer. Math. Soc. 30 (1971), 199-201.

University of Pittsburgh

Pittsburgh, PA 15260

and

Institute of Mathematics of the Polish Academy of Sciences

00-950 Warsaw