

---

# TOPOLOGY PROCEEDINGS



Volume 2, 1977

Pages 479–482

---

<http://topology.auburn.edu/tp/>

## ON THE CANCELLATION OF SHAPES FROM PRODUCTS

by

J. G. HOLLINGSWORTH AND R. B. SHER

---

### Topology Proceedings

**Web:** <http://topology.auburn.edu/tp/>

**Mail:** Topology Proceedings  
Department of Mathematics & Statistics  
Auburn University, Alabama 36849, USA

**E-mail:** [topolog@auburn.edu](mailto:topolog@auburn.edu)

**ISSN:** 0146-4124

COPYRIGHT © by Topology Proceedings. All rights reserved.

## ON THE CANCELLATION OF SHAPES FROM PRODUCTS

J. G. Hollingsworth and R. B. Sher

K. Borsuk [1] has raised the following question concerning the "cancellation law" for shapes of compacta: Is it true that  $\text{Sh}(X) \times \text{Sh}(S^1) = \text{Sh}(Y) \times \text{Sh}(S^1)$  implies that  $\text{Sh}(X) = \text{Sh}(Y)$ ? The answer to this question is generally "no." Specifically, Charlap [2] has constructed compact manifolds  $M$  and  $N$  such that  $M \not\cong N$  and  $M \times S^1 \cong N \times S^1$ . (In fact,  $M$  and  $N$  are Riemannian manifolds and  $M \times S^1$  is diffeomorphic to  $N \times S^1$ .) As in the case of the analogous question in the homotopy theory of compact ANR's, the problem lies in the fundamental group. Thus, in this brief note we are able to establish the following.

*Theorem.* Suppose  $X$  and  $Y$  are approximately 1-connected compacta and  $\text{Sh}(X) \times \text{Sh}(S^1) = \text{Sh}(Y) \times \text{Sh}(S^1)$ . Then  $\text{Sh}(X) = \text{Sh}(Y)$ .

For our purposes, we shall find most suitable the ANR-systems approach to shape due to Mardešić and Segal [3]. By [4], this approach may be regarded as equivalent to Borsuk's.

We shall adopt, along with the terminology of [3], the following notations.

- (i)  $\mathbb{R}$  shall denote the real numbers;
- (ii)  $S^1$  is the set of complex numbers  $z$  such that  $|z| = 1$ ;
- (iii)  $\exp: \mathbb{R} \rightarrow S^1$  is the exponential map;
- (iv) If  $A$  is a space, then  $\alpha: A \rightarrow A \times S^1$  is defined by  $\alpha(a) = (a, 1)$  for all  $a \in A$ ;

(v) If  $A$  and  $B$  are spaces,  $\pi: A \times B \rightarrow A$  is defined by

$$\pi(a, b) = a \text{ for all } (a, b) \in A \times B.$$

Of course  $\alpha$  and  $\pi$  depend on the space  $A$  and, in the case of  $\pi$ ,  $B$ . However, context shall make the meaning clear in each case.

*Proof of the theorem.* We may regard  $X$  and  $Y$  as being subsets of  $Q$ , the Hilbert cube. Let  $\underline{X} : U_1 \supset U_2 \supset U_3 \supset \dots$  be an inclusion ANR-sequence for  $X$  in  $Q$  and  $\underline{Y} : V_1 \supset V_2 \supset V_3 \supset \dots$  be an inclusion ANR-sequence for  $Y$  in  $Q$ . Then  $\underline{X}' : U_1 \times S^1 \supset U_2 \times S^1 \supset U_3 \times S^1 \supset \dots$  and  $\underline{Y}' : V_1 \times S^1 \supset V_2 \times S^1 \supset V_3 \times S^1 \supset \dots$  are inclusion ANR-sequences for  $X \times S^1$  and  $Y \times S^1$ , respectively, in  $Q \times S^1$ . Hence, there exist system maps  $\underline{f}' = (f', f'_i) : \underline{X}' \rightarrow \underline{Y}'$  and  $\underline{g}' = (g', g'_i) : \underline{Y}' \rightarrow \underline{X}'$  such that  $\underline{g}' \underline{f}' \approx \underline{1}_{\underline{X}'}$ , and  $\underline{f}' \underline{g}' \approx \underline{1}_{\underline{Y}'}$ . Let  $f = f'$  and, for each positive integer  $i$ ,  $f_i = \pi f'_i \alpha : U_{f(i)} \rightarrow V_i$ . Similarly, let  $g = g'$  and  $g_i = \pi g'_i \alpha : V_{g(i)} \rightarrow U_i$ . The proof will be complete if we are able to establish that:

(1)  $\underline{f} = (f, f_i) : \underline{X} \rightarrow \underline{Y}$  and  $\underline{g} = (g, g_i) : \underline{Y} \rightarrow \underline{X}$  are system maps, and


(2)  $\underline{g} \underline{f} \approx \underline{1}_{\underline{X}}$  and  $\underline{f} \underline{g} \approx \underline{1}_{\underline{Y}}$ .

Assertion (1) is easily verified, and we shall not present the details. The reader familiar with shape theory will note that there are obvious system maps  $\underline{\alpha} : \underline{X} \rightarrow \underline{X}'$  and  $\underline{\pi} : \underline{Y}' \rightarrow \underline{Y}$ , associated with the maps  $\alpha : X \rightarrow X \times S^1$  and  $\pi : Y \times S^1 \rightarrow Y$  respectively, so that  $\underline{f} = \underline{\pi} \underline{f}' \underline{\alpha}$ . A similar comment applies to  $\underline{g}$ .

To confirm the first part of (2), we are required to show that if  $i$  is a positive integer, then there exists an

integer  $k \geq f(g(i))$  such that  $l_{U_k} \approx g_i f'_{g(i)} |_{U_k}$  in  $U_i$ . Since  $g' f' \approx l_X$ , we know that there exist a positive integer  $j \geq f(g(i))$  and a homotopy  $H : U_j \times S^1 \times I \rightarrow U_i \times S^1$  such that for all  $(x, y) \in U_j \times S^1$ ,  $H(x, y, 0) = (x, y)$  and  $H(x, y, 1) = g'_i f'_{g(i)}(x, y)$ . Since  $X$  is approximatively 1-connected, there exists an integer  $k \geq j$  such that each loop in  $U_k$  is contractible in  $U_j$ . Define  $G : U_k \times I \rightarrow U_i$  by  $G(x, t) = \pi H(x, 1, t)$  for all  $x \in U_k$  and  $t \in I$ . Then, if  $x \in U_k$ ,  $G(x, 0) = \pi H(x, 1, 0) = \pi(x, 1) = x$  and  $G(x, 1) = \pi H(x, 1, 1) = \pi g'_i f'_{g(i)}(x, 1) = \pi g'_i f'_{g(i)} \alpha(x)$ . We conclude that  $l_{U_k} \approx \pi g'_i f'_{g(i)} \alpha$  in  $U_i$ .

Since each loop in  $U_k$  is contractible in  $U_j$ , the map  $(f'_{g(i)} |_{U_k \times S^1}) \alpha : U_k \rightarrow V_{g(i)} \times S^1$  can be lifted through the covering  $l_{V_{g(i)}} \times \exp : V_{g(i)} \times R \rightarrow V_{g(i)} \times S^1$  to a map  $\tilde{f} : U_k \rightarrow V_{g(i)} \times R$ . Consider the following diagram.



The two "outermost" triangles commute, while the "innermost" triangle commutes up to homotopy via the map  $K : V_{g(i)} \times R \times I \rightarrow V_{g(i)} \times S^1$  defined by  $K(x, r, t) = (x, \exp(rt))$  for all  $(x, r, t) \in V_{g(i)} \times R \times I$ . It thus follows that, as maps into  $U_i$ ,  $l_{U_k} \approx \pi g'_i f'_{g(i)} \alpha = \pi g'_i (l \times \exp) \tilde{f} \approx \pi g'_i \alpha \pi \tilde{f} = \pi g'_i \alpha \pi (l \times \exp) \tilde{f} = (\pi g'_i \alpha) (\pi (f'_{g(i)} |_{U_k \times S^1}) \alpha) = g_i f_{g(i)} |_{U_k}$ . We have thus succeeded

in showing that  $\underline{g} \underline{f} \approx \underline{1}_X$ . Of course a symmetric argument shows that  $\underline{f} \underline{g} \approx \underline{1}_Y$ , and the proof is complete.

*Remark.* It is obvious that  $S^1$  may be replaced in the above proof by any complex  $K$  of type  $(II,1)$ . We need only replace  $\exp:R \rightarrow S^1$  by the universal cover  $P:\tilde{K} \rightarrow K$ , noting that  $\tilde{K}$  is contractible.

### **Bibliography**

1. Karol Borsuk, *On several problems of the theory of shape*, Studies in Topology, Academic Press, N.Y., 1975, 67-69.
2. Leonard S. Charlap, *Compact flat Riemannian manifolds: I*, Ann. of Math. (2) 81 (1965), 15-30.
3. Sibe Mardešić and Jack Segal, *Shapes of compacta and ANR-systems*, Fund. Math. 72 (1971), 41-59.
4. \_\_\_\_\_, *Equivalence of the Borsuk and ANR-system approach to shapes*, Fund. Math. 72 (1971), 61-68.

The University of Georgia

Athens, GA 30602

The University of North Carolina at Greensboro

Greensboro, NC 27412