
TOPOLOGY PROCEEDINGS



Volume 2, 1977

Pages 621–630

<http://topology.auburn.edu/tp/>

PL HOMOLOGY 3-SPHERES AND TRIANGULATIONS OF MANIFOLDS

by

RONALD J. STERN

Topology Proceedings

Web: <http://topology.auburn.edu/tp/>

Mail: Topology Proceedings
Department of Mathematics & Statistics
Auburn University, Alabama 36849, USA

E-mail: topolog@auburn.edu

ISSN: 0146-4124

COPYRIGHT © by Topology Proceedings. All rights reserved.

PL HOMOLOGY 3-SPHERES AND TRIANGULATIONS OF MANIFOLDS

Ronald J. Stern¹

One of the least understood but important groups arising in geometric topology is θ_3^H , the abelian group obtained from the set of oriented 3-dimensional PL homology spheres using the operation of connected sum, modulo those which bound acyclic PL 4-manifolds.

In this paper we will show how the group θ_3^H and the following theorem of Rohlin play an important role in triangulating topological manifolds.

V. A. Rohlin Signature Theorem ([4], [8], [12], [15]).
Every closed oriented smooth 4-manifold M^4 whose second Stiefel-Whitney class vanishes has signature $\sigma(M) \in \mathbb{Z}$ divisible by 16.

By classical smoothing theory the same theorem holds for PL 4-manifolds. However, the theorem is an important undecided result for topological 4-manifolds.

Given a PL homology 3-sphere H^3 , then H^3 bounds a parallelizable PL 4-manifold W^4 . Let $\alpha(H^3) \in \mathbb{Z}_2$ be the Kervaire-Milnor-Rohlin invariant given by

$$\alpha(H^3) = \alpha(\bar{W}^4)/8 \pmod{2}$$

where \bar{W}^4 is the closed (polyhedral) homology 4-manifold $\bar{W}^4 = W^4 \cup_{H^3} cH^3$. That $\sigma(\bar{W}^4)$ is divisible by 8 follows from the fact that the cup-product pairing of \bar{W}^4 is an even

¹Supported in part by NSF grant MCS 76-06393.

quadratic form (see [14]). Also, an application of Rohlin's theorem and the additivity properties of the signature show that $\alpha(H^3)$ is independent of the bounding parallelizable manifold W^4 , and in fact α determines a well-defined homomorphism $\alpha: \theta_3^H \rightarrow \mathbb{Z}_2$.

The classical example of a PL homology 3-sphere is the Poincaré homology 3-sphere $H^3 = SO(3)/A_5$, where A_5 is the even permutation group on five objects = the 60 orientation preserving symmetries of the dodecahedron. In fact $\pi_1(X) = \langle a, b \mid a^3 = b^5 = (ab)^2 \rangle \neq 0$. Now H^3 is also the boundary of the Milnor plumbing W^4 of 8 copies of the unit tangent disk bundle of S^2 according to the Dynkin diagram

$$\begin{array}{c} \text{o}-\text{o}-\text{o}-\text{o}-\text{o}-\text{o}-\text{o} \\ | \\ \text{o} \end{array}$$

(For this and other descriptions of H^3 see [9]). It is then an excellent exercise for an algebraic topology class to show that $\sigma(\overline{W}^4) = 8$, thus demonstrating that the homomorphism $\alpha: \theta_3^H \rightarrow \mathbb{Z}_2$ is surjective.

We now have a short exact sequence

$$0 \rightarrow \ker(\alpha) \rightarrow \theta_3^H \xrightarrow{\alpha} \mathbb{Z}_2 \rightarrow 0$$

Not one concrete fact is known about θ_3^H other than the existence of the surjection $\alpha: \theta_3^H \rightarrow \mathbb{Z}_2$. For instance, it is not even known if θ_3^H is finitely generated.

The importance of the group $\ker(\alpha)$ is explained by

Theorem 1. (Galewski-Stern [6], [7], T. Matumoto [13]).
Let M^m be a topological manifold with ∂M triangulated as a simplicial complex. If $m \geq 6$, then there is an element $\tau(M) \in H^5(M, \partial M; \ker(\alpha))$ such that $\tau(M) = 0$ if and only if

there is a simplicial triangulation K of M with $K|_{\partial M}$ compatible with the given triangulation on ∂M . Moreover, there are $|H^4(M, \partial M; \ker(\alpha))|$ such triangulations on M up to concordance rel ∂M .

Two triangulations K_0 and K_1 of M are concordant rel ∂M if there is a triangulation K of $M \times I$ such that $K|_{(\partial M \times I)}$ is compatible with the given triangulation on $\partial M \times I$ and such that $K|M \times \{i\}$ is compatible with K_i for $i = 0, 1$.

It should be noted that *a priori* there are two obstructions in Theorem 1 to triangulating M , but one of these obstructions vanish since

Theorem 2. (R. D. Edwards [3]). Let H^3 be any PL homology 3-sphere. Then $S^2 * H^3 = \Sigma^3 H^3$ is homeomorphic to S^6 .

The importance of even a little understanding of θ_3^H is explained by

Corollary 3. (Galewski-Stern [6], [7], T. Matumoto [13]). If there is an element $x \in \theta_3^H$ with $\alpha(x) = 1$ and $2x = 0$, then all topological m -manifolds, $m \geq 6$ (≥ 7 if $\partial M \neq \emptyset$) can be triangulated as simplicial complexes.

This leads us to the following well-known conjectures.

Conjecture 1. The group θ_3^H contains an element of order two.

Conjecture 2. The group $\ker(\alpha)$ is the zero group.

In the remainder of this paper we will first consider the relationship between the simplicial triangulation obstruction of Theorem 1 and the combinatorial triangulation

obstruction of Kirby-Siebenmann [10]. These results are implicit in [7]. We will then show how elements of θ_3^H yield exotic PL manifold structures on $I^k \times T^n$, $k + n \geq 5$, which are the standard PL structure on $\partial I^k \times T^n$, and how elements of $\ker(\alpha)$ yield *possibly* exotic simplicial manifold structures on $I^k \times T^n$, $k + n \geq 5$, which are the standard PL manifold structure on $\partial I^k \times T^n$. Furthermore, we will demonstrate that the exotic PL manifold structures on $I^k \times T^n$ are non-exotic when considered as simplicial manifold structures. Here, T^n is the cartesian product of S^1 n -times, and I^k is the cartesian product of the unit interval k -times.

Recall the short exact sequence

$$0 \rightarrow \ker(\alpha) \rightarrow \theta_3^H \xrightarrow{\alpha} \mathbb{Z}_2 \rightarrow 0$$

and the resulting Bockstein exact coefficient sequence

$$\cdots \rightarrow H^4(M; \mathbb{H}_3) \xrightarrow{\alpha_*} H^4(M; \mathbb{Z}_2) \xrightarrow{\beta} H^5(M; \ker(\alpha)) \rightarrow \cdots$$

If M is a polyhedral homology m -manifold, there is an element $r(M) \in H^4(M; \theta_3^H)$ such that $r(M) = 0$ if and only if there is a PL m -manifold N and a PL acyclic map $f: N \rightarrow M$ (see [1], [2], [11]). If M is also a topological manifold, let $\Delta(M) \in H^4(M; \mathbb{Z}_2)$ denote the Kirby-Siebenmann obstruction to putting a PL manifold structure on M [10]. Theorem 11.1 of [7] shows that

$$(4) \quad \alpha_* r(M) = \Delta(M)$$

Note that this identifies $\Delta(M)$ as a simplicial cohomology class in the case that M has a simplicial triangulation K . For then $\Delta(M)$ is represented by the cocycle $\alpha': C_4(M) \rightarrow \mathbb{Z}_2$, where $C_4(M)$ is the free abelian group generated by the dual 4-cells e_σ^4 of the $(m-4)$ -simplices σ of K and $\alpha'(e_\sigma^4) = \alpha(\text{link}(\sigma, K))$. In particular, a topological m -manifold M ,

$m \geq 5$ (≥ 6 if $\partial M \neq \emptyset$) has a PL manifold structure if and only if there is a simplicial triangulation K of M such that the Kervaire-Milnor-Rohlin invariant of every 3-dimensional link of K is zero.

Since the obstruction to putting a PL manifold structure on a topological manifold M and the obstruction to putting a simplicial manifold structure on M are lifting obstructions, standard obstruction theory shows that

$$(5) \quad \beta \Delta(M) = \tau(M)$$

Furthermore, we have by Corollary 12.5 of [7] that if $\beta_*: H^4(M; \mathbb{Z}_2) \rightarrow H^5(M; \mathbb{Z})$ is the integral Bockstein homomorphism, then $\beta_* \Delta(M) = 0$ implies that $\tau(M) = 0$.

Now to the construction of exotic triangulations on $I^k \times T^n$, $k + n \geq 5$. Let $\mathcal{S}_{PL}(I^k \times T^n, \partial)$ denote the set of concordance (hence isotopy) classes of PL manifold structures on $I^k \times T^n$ extending the standard PL manifold structure on $\partial I^k \times T^n$. Similarly, let $\mathcal{S}_{TRI}(I^k \times T^n, \partial)$ denote the set of concordance classes of simplicial manifold structures on $I^k \times T^n$ extending the standard PL manifold structure on $\partial I^k \times T^n$. There is a natural map $\mathcal{S}_{PL}(I^k \times T^n, \partial) \rightarrow \mathcal{S}_{TRI}(I^k \times T^n, \partial)$. By the work of Kirby-Siebenmann [10], $|\mathcal{S}_{PL}(I^k \times T^n, \partial)| = |H^{3-k}(T^n; \mathbb{Z}_2)|$ for $k + n \geq 5$; and by Theorem 1, $|\mathcal{S}_{TRI}(I^k \times T^n, \partial)| = |H^{4-k}(T^n; \ker(\alpha))|$ for $k + n \geq 6$. Our goal is to construct non-trivial elements of $\mathcal{S}_{PL}(I^k \times T^n, \partial)$ and $\mathcal{S}_{TRI}(I^k \times T^n, \partial)$ and show that the natural map $\mathcal{S}_{PL}(I^3 \times T^n, \partial) \rightarrow \mathcal{S}_{TRI}(I^3 \times T^n, \partial)$ is the zero map for $n \geq 2$.

Let H^3 be a PL homology 3-sphere and let $F^3 = H^3 - \text{int } I^3$. Then $F^3 \times I^n$ has as boundary a PL homology

$(n+2)$ -sphere H^{n+2} . Hence, by doing surgery on the interior of the parallelizable manifold $F^3 \times I^n$ we have that H^{n+2} bounds a contractible $PL(n+3)$ -manifold P^{n+3} if $n \geq 2$. By identifying the I^n factor in $\partial P^{n+3} = H^{n+2} = \partial(F^3 \times I^n)$ so as to get T^n , we have a $PL(n+3)$ -manifold M^{n+3} with boundary $S^2 \times T^n$. Note that M^{n+3} is homotopy equivalent rel ∂M^{n+3} to $I^3 \times T^n$. Since any manifold which is homotopy equivalent to $I^k \times T^n$ rel ∂ , $k + n \geq 5$, is homeomorphic to $I^k \times T^n$ rel ∂ [16], we have that M^{n+3} is homeomorphic to $I^3 \times T^n$ rel ∂ . The image of the PL manifold structure on M^{n+3} under this homeomorphism yields a PL manifold structure Γ_{H^3} on $I^3 \times T^n$ extending the standard PL manifold structure on $\partial I^3 \times T^n$, hence determines an element $[\Gamma_{H^3}] \in \mathcal{S}_{PL}(I^3 \times T^n, \partial)$.

Theorem 6. $[\Gamma_{H^3}] = 0$ if and only if $\alpha(H^3) = 0$. Furthermore, the natural map $\mathcal{Z}_2 \cong \mathcal{S}_{PL}(I^3 \times T^n, \partial) \rightarrow \mathcal{S}_{TRI}(I^3 \times T^n, \partial) \cong n \ker(\alpha)$ is the zero map.

Proof. Let M^{n+3} be as above and let $h: M^{n+3} \rightarrow I^3 \times T^n$ be a homeomorphism which is the identity over $\partial I^3 \times T^n$. Attach a copy of $I^3 \times T^n$ to M^{n+3} along ∂M^{n+3} to obtain a PL manifold M' homeomorphic to $S^3 \times T^n$. Note that M' bounds $Q^{n+4} = (CH^3 \times T^n) \cup \{\text{handles}\}$. By Theorem 2 Q^{n+4} is simplicially triangulated topological manifold which is an s -cobordism between M^{n+3} and $I^3 \times T^n$ rel ∂ . By the observations following (4), $CH^3 \times T^n$ possesses a PL manifold structure extending the natural one on $H^3 \times T^n$ if and only if $\alpha(H^3) = 0$. This also follows from Theorem C of [17].

Let Q_*^{n+4} be the topological manifold obtained by adjoining the mapping cylinder of h to Q^{n+4} along M^{n+3} . Then

Q_*^{n+4} is homotopy equivalent to $I^4 \times T^n \text{ rel } \partial Q_*^{n+4} = S^3 \times T^n$, hence Q_*^{n+4} is homeomorphic to $I^4 \times T^n \text{ rel } \partial I^4 \times T^n$. This homeomorphism induces a simplicial manifold structure Σ_{H^3} on $I^4 \times T^n$ and a simplicial manifold concordance between Γ_{H^3} and the standard structure on $I^3 \times T^n$. It now follows that $[\Gamma_{H^3}] = 0$ if and only if $\alpha(H^3) = 0$ and that $\mathcal{S}_{PL}(I^3 \times T^n, \partial) \rightarrow \mathcal{S}_{TRI}(I^3 \times T^n, \partial)$ is the zero map for $n \geq 2$.

Perhaps Theorem 6 morally justifies Conjecture 2 above.

Note that in the proof of Theorem 6, the simplicial manifold structure Σ_{H^3} on $I^4 \times T^n$ can be assumed to be the standard PL manifold structure on $\partial I^4 \times T^n$ if $\alpha(H^3) = 0$, for we can assume that $h: M^{n+3} \rightarrow I^3 \times T^n$ is a PL homeomorphism. There results for each PL homology 3-sphere H^3 with $\alpha(H^3) = 0$ an element $[\Sigma_{H^3}] \in \mathcal{S}_{TRI}(I^4 \times T^n, \partial) \cong \ker(\alpha)$.

Theorem 7. Let H^3 and \bar{H}^3 be PL homology 3-spheres with $\alpha(H^3) = \alpha(\bar{H}^3) = 0$. Then $[\Sigma_{H^3}] = [\Sigma_{\bar{H}^3}]$ if and only if there is a PL homology cobordism between H^3 and \bar{H}^3 .

Proof. Recall that Σ_{H^3} is the simplicial manifold structure on $I^4 \times T^n$ induced by a simplicial manifold structure on the topological manifold $Q^{n+4} = (cH^3 \times T^n) \cup \{\text{handles}\}$. Similarly, $\Sigma_{\bar{H}^3}$ is induced by a simplicial manifold structure on the topological manifold $\bar{Q}^{n+4} = (c\bar{H}^3 \times T^n) \cup \{\text{handles}\}$. Let X^4 be a PL homology cobordism between H^3 and \bar{H}^3 . Now $H^4 = cH^3 \cup X^4 \cup c\bar{H}^4$ is a polyhedral homology manifold having the homology of S^4 and with $H^4 \times \mathbb{R}^2$ a topological manifold. By Theorem 1.4 of [7], $cH^4 \times \mathbb{R}^2$ is a topological manifold so that $P^{n+5} = (Q^{n+4} \times I) \cup (cH^4 \times T^n) \cup (\bar{Q}^{n+4} \times I)$ is a

simplicially triangulated topological manifold. By attaching PL handles to $X^4 \times T^n \subset P^{n+5}$ we obtain a simplicially triangulated topological manifold $Y^{n+5} = (Q^{n+4} \times I) \cup (cH^4 \times T^n) \cup \{\text{handles}\} \cup (\bar{Q}^{n+4} \times I)$ homotopy equivalent to $I^5 \times T^n$. An application of the PL s-cobordism theorem to the boundary PL s-cobordism in Y^{n+5} between ∂Q^{n+4} and $\partial \bar{Q}^{n+4}$, each of which we identify with $\partial I^4 \times T^n$ via a PL homeomorphism, yields a simplicially triangulated topological manifold Y_*^{n+5} homotopy equivalent to $I^5 \times T^n$ and with $\partial Y_*^{n+5} = Q^{n+4} \cup S^3 \times T^n \times I \cup \bar{Q}^{n+4}$. There results a concordance between Σ_{H^3} and $\Sigma_{\bar{H}^3}$.

Conversely, suppose $[\Sigma_{H^3}] = [\Sigma_{\bar{H}^3}]$. Then there is a simplicially triangulated topological manifold W^{n+5} with $\partial W^{n+5} = Q^{n+4} \cup S^3 \times T^n \times I \cup \bar{Q}^{n+4}$ and with a homeomorphism $f: W^{n+5} \rightarrow I^4 \times T^n \times I$ which is PL over $\partial I^4 \times T^n \times I$. Let $\pi: cH^3 \times T^n \rightarrow I^4 \times T^n$ and $\bar{\pi}: c\bar{H}^3 \times T^n \rightarrow I^4 \times T^n$ be homology equivalences which fiber over T^n . There is no obstruction to extending π and $\bar{\pi}$ to a homotopy equivalence $p: W^{n+5} \rightarrow I^4 \times T^n \times I$, with $p|f^{-1}(\partial I^4 \times T^n \times I): f^{-1}(\partial I^4 \times T^n \times I) \rightarrow \partial I^4 \times T^n \times I$ a homotopy equivalence. Homotope p so that $p|f^{-1}(\partial I^4 \times T^n \times I)$ is PL transverse to $S^3 \times (\text{pt.}) \times I$. There results a PL cobordism X^4 between H^3 and \bar{H}^3 . There is a transversality theory for maps of homology manifolds provided that the target is a PL manifold (Theorem 3.7 of [5]). So homotope p rel $f^{-1}(\partial I^4 \times T^n \times I)$ to be homology transverse to $I^4 \times (\text{pt.}) \times I$. There results a homology manifold cobordism X^5 between cH^3 and $c\bar{H}^3$ extending X^4 . Now ∂X^5 bounds the homology manifold X^5 , so that $r(\partial X^5) = 0$. But the only non PL 3-sphere links of ∂X^5 are H^3 and \bar{H}^3 . Thus $H^3 \# -\bar{H}^3$

bounds a PL acyclic 4-manifold, hence H^3 and \bar{H}^3 are PL homology cobordant.

To construct non-trivial elements of $\mathcal{S}_{PL}(I^k \times T^n, \partial) \cong H^{3-k}(T^n; \mathbb{Z}_2)$, $k + n \geq 5$, take the non-trivial element of $\mathcal{S}_{PL}(I^3 \times T^{n-k}, \partial)$ constructed above and identify the opposite ends of k -interval factors of I^3 to derive a non-trivial element of $\mathcal{S}_{PL}(I^k \times T^n, \partial)$. Note that these elements are trivial when considered as elements of $\mathcal{S}_{TRI}(I^k \times T^n, \partial)$. Similarly one constructs (possibly) non-trivial elements of $\mathcal{S}_{TRI}(I^k \times T^n, \partial) \cong H^{4-k}(T^n; \ker(\alpha))$, $k + n \geq 6$, from the elements of $\mathcal{S}_{TRI}(I^4 \times T^n, \partial)$ constructed above.

Bibliography

1. M. Cohen, *Homeomorphism between homotopy manifolds and their resolutions*, Inv. Math. 10 (1970), 239-250.
2. A. Edmonds and R. Stern, *Resolutions of homology manifolds: A classification theorem*, J. London Math. Soc. (2), 11 (1975), 474-480.
3. R. D. Edwards,
4. M. Freedman and R. Kirby, *A geometric proof of Rohlin's theorem*, preprint, 1975.
5. D. Galewski and R. Stern, *The relationship between homology and topological manifolds via homology transversality*, to appear in Inv. Math.
6. _____, *Classification of simplicial triangulations of topological manifolds*, Bull. Amer. Math. Soc. 82 (1976), 916-918.
7. _____, *Classification of simplicial triangulations of topological manifolds*, preprint, 1976.
8. M. Kervaire and J. Milnor, *Bernoulli numbers, homotopy groups and a theorem of Rohlin*, Proc. Int. Congr. Math., Edinburg, 1958, 454-458.

9. R. Kirby and M. Scharlemann, *Eight faces of the dodecahedral 3-manifold*, preprint, 1976.
10. R. Kirby and L. Siebenmann, *Foundational essays on topological manifolds, smoothings, and triangulations*, Annals of Mathematics Studies No. 88, Princeton University Press, Princeton, NJ.
11. N. Martin, *On the difference between homology and piecewise linear block bundles*, J. London Math. Soc. 6 (1973), 197-204.
12. Y. Matsumoto, *An elementary proof of Rohlin's theorem*, preprint, 1976.
13. _____, *Variétés simpliciales d'homologie et variétés topologiques métrisables*, Thèse, Univ. de Paris-Sud, 91405 Orsay, 1976.
14. J. Milnor and J. Husemoller, *Symmetric bilinear forms*, Springer, 1973.
15. V. A. Rohlin, *A new result in the theory of 4-dimensional manifolds*, Soviet Math. Doklady 8 (1952), 221-224 (in Russian).
16. L. Siebenmann, *Disruption of low-dimensional handlebody theory by Rohlin's theorem*, in *Topology of Manifolds*, edited by Cantrell and Edwards, Univ. of Georgia, 1970.
17. L. Siebenmann, *Are non-triangulable manifolds triangulable?* in *Topology of Manifolds*, edited by Cantrell and Edwards, University of Georgia, 1970.

University of Utah

Salt Lake City, UT 84112