
TOPOLOGY PROCEEDINGS



Volume 3, 1978

Pages 319–333

<http://topology.auburn.edu/tp/>

ON LC^n -DIVISORS

by

JERZY DYDAK

Topology Proceedings

Web: <http://topology.auburn.edu/tp/>

Mail: Topology Proceedings
Department of Mathematics & Statistics
Auburn University, Alabama 36849, USA

E-mail: topolog@auburn.edu

ISSN: 0146-4124

COPYRIGHT © by Topology Proceedings. All rights reserved.

ON LC^n -DIVISORS

Jerzy Dydak

1. Introduction

In [19] D. M. Hyman introduced the class of ANR-divisors i.e. continua X such that Y/X is an ANR for each ANR-space Y containing X . By using Chapman's Complement Theorem [10] it is easy to show that being an ANR-divisor is a shape invariant (see [9]). More generally we have: being an ANR-divisor is a hereditary strong shape invariant in the sense of Edwards-Hastings [14] (see [9]). By using the work of Hyman [19] and characterizations of pointed FANR's it is clear that pointed FANR's are ANR-divisors. However an example in [12] shows that the class of ANR-divisors is wider than the class of pointed FANR's.

In this paper we introduce the class of LC^n -divisors in analogy to ANR-divisors. We give a characterization of LC^n -divisors which implies that being an LC^n -divisor is a hereditary shape invariant. As a consequence we infer that being an ANR-divisor is a hereditary shape invariant in the class of continua of finite fundamental dimension.

We assume that the reader is familiar with some elementary facts from shape theory (see [6], [22], [23] and [27]) and from the theory of pro-categories (see [2], [11] and [14]).

2. Some Algebraic Preliminaries

For a definition and basic properties of pro-categories (see [2], [11] and [14]).

Recall that an inverse sequence $\underline{A} = (A_n, p_n^{n+1})$ of groups is said to satisfy the Mittag-Leffler condition provided for each n there exists $k > n$ such that

$$\text{im } p_n^k = \text{im } p_n^m \quad \text{for } m > k$$

(see [3] and [26]).

\underline{A} is said to be stable iff \underline{A} is isomorphic to a group in the category of pro-groups pro-Gr (see [11] and [14]).

For a definition of $\varprojlim^1 \underline{A}$ and its properties see [8], pp. 250-252.

Lemma 2.1. Let $\underline{A} = (A_n, p_n^{n+1})$ and $\underline{B} = (B_n, q_n^{n+1})$ be inverse sequences of groups and let $f_n: A_n \rightarrow B_n$ be homomorphisms such that $q_n^{n+1} f_{n+1} = f_n p_n^{n+1}$ for $n \geq 1$.

If $\varprojlim^1 \underline{A} = *$, then $\varprojlim^1 \underline{C} = *$, where $\underline{C} = (\text{im } f_n, r_n^{n+1})$ and r_n^{n+1} is induced by q_n^{n+1} .

If \underline{A} and \underline{B} are stable, then $\underline{D} = (\ker f_n, s_n^{n+1})$ is stable, where s_n^{n+1} is induced by p_n^{n+1} .

Proof. Suppose $\varprojlim^1 \underline{A} = *$. Since

$$0 \rightarrow \ker f_n \rightarrow A_n \rightarrow \text{im } f_n \rightarrow 0$$

is exact for each n , then the following sequence is exact:

$$0 \rightarrow \varprojlim^1 \underline{D} \rightarrow \varprojlim^1 \underline{A} \rightarrow \varprojlim^1 \underline{C} \rightarrow \varprojlim^1 \underline{D} \rightarrow \varprojlim^1 \underline{A} \rightarrow \varprojlim^1 \underline{C} \rightarrow 0$$

(see [8], p. 252).

Hence, $\varprojlim^1 \underline{C} = *$.

Suppose \underline{A} and \underline{B} are stable. Then we may assume that

$$p_n^{n+1} / \text{im } p_{n+1}^{n+2}: \text{im } p_{n+1}^{n+2} \rightarrow \text{im } p_n^{n+1} \quad \text{and}$$

$$q_n^{n+1} / \text{im } q_{n+1}^{n+2}: \text{im } q_{n+1}^{n+2} \rightarrow \text{im } q_n^{n+1}$$

are isomorphisms for each n .

Let $x \in \ker f_{n+1}$. Then there is $y \in A_{n+3}$ with $p_n^{n+3}(y) = p_n^{n+1}(x)$. Hence, $q_n^{n+2}q_{n+2}^{n+3}f_{n+3}(y) = q_n^{n+3}f_{n+3}(y) = f_n p_n^{n+3}(y) = f_n p_n^{n+1}(y) = q_n^{n+1}f_{n+1}(x) = 0$. Therefore, $f_{n+2}p_{n+2}^{n+3}(y) = q_{n+2}^{n+3}f_{n+3}(y) = 0$ i.e. $p_{n+2}^{n+3}(y) \in \ker f_{n+2}$. Since $p_n^{n+2}p_{n+2}^{n+3}(y) = p_n^{n+3}(y) = p_n^{n+1}(x)$ we get $p_n^{n+2}(\ker f_{n+2}) = p_n^{n+1}(\ker f_{n+1})$. Hence

$$s_n^{n+1}/\text{im } s_{n+1}^{n+2}: \text{im } s_{n+1}^{n+2} \rightarrow \text{im } s_n^{n+1}$$

is an isomorphism for each n and \underline{D} is stable.

Let $p_n^{k-1,k}: G_n^k \rightarrow G_n^{k-1}$ and $q_{n,n+1}^k: G_{n+1}^k \rightarrow G_n^k$ be homomorphisms of groups ($n \geq 1$ and k -integer) such that

$$p_n^{k-1,k}q_{n,n+1}^k = q_{n,n+1}^{k-1}p_n^{k-1,k}.$$

Suppose that each sequence $\underline{G}_n = (G_n^k, p_n^{k-1,k})$ is exact and let $\underline{G}^k = (G_n^k, q_{n,n+1}^k)$ for each k .

Lemma 2.2. If $\varinjlim \underline{G}^k = *$ for each k , then the sequence

$$\dots \rightarrow \lim \underline{G}^k \rightarrow \lim \underline{G}^{k-1} \rightarrow \lim \underline{G}^{k-2} \rightarrow \dots$$

is exact.

Proof. Analogous to the corresponding result in [30], where Lemma 2.2 is proved in case where \underline{G}^k satisfy the Mittag-Leffler condition.

Lemma 2.3. If \underline{G}^i are stable for $i = 0, 1$ and $\varinjlim \underline{G}^3 = *$, then $\varinjlim \underline{G}^2 = *$.

Proof. For each n we have the following exact sequence

$$0 \rightarrow \text{im } p_n^{2,3} \rightarrow G_n^2 \rightarrow \ker p_n^{0,1} \rightarrow 0.$$

By Proposition 2.3 in [8] (p. 252) there is the following

exact sequence $0 \rightarrow \varprojlim (\text{im } p_n^{2,3}) \rightarrow \varprojlim \underline{G}^2 \rightarrow \varprojlim (\ker p_n^{0,1}) \rightarrow \varprojlim^1 (\text{im } p_n^{2,3}) \rightarrow \varprojlim^1 \underline{G}^2 \rightarrow \varprojlim^1 (\ker p_n^{0,1}) \rightarrow 0$. By Lemma 2.1 the inverse sequence $(\ker p_n^{0,1})$ is stable and $\varprojlim^1 (\text{im } p_n^{2,3}) = *$. Hence $\varprojlim^1 (\ker p_n^{0,1}) = *$ (see [8], p. 252) and consequently $\varprojlim^1 \underline{G}^2 = *$.

Lemma 2.4. *If \underline{G}^i is isomorphic to an inverse sequence of countable groups for $i = 1, 3$, then \underline{G}^2 is isomorphic in pro-Gr to an inverse sequence of countable groups.*

Proof. An inverse sequence (A_n, r_n^m) of groups is isomorphic to an inverse sequence of countable groups iff for each n there is $m > n$ such that $r_n^m(A_m)$ is countable.

Let $n \geq 1$ and take $k > m > n$ such that $q_{n,m}^i(G_m^i)$ is countable for $i = 1, 3$ and $q_{m,k}^i(G_k^i)$ is countable for $i = 1, 3$. Take elements $a_i \in G_m^2$, $i \geq 1$, such that each element in $p_m^{1,2}(G_m^2) \cap q_{m,k}^1(G_k^1)$ is equal to $p_m^{1,2}(a_i)$ for some i .

Let $a \in G_k^2$ be an arbitrary element. Then $q_{m,k}^1 p_k^{1,2}(a) = p_m^{1,2}(a_i)$ for some i . Thus $p_m^{1,2}(a_i q_{m,k}^2(a^{-1})) = 0$ and there is $b \in G_m^3$ with $p_m^{2,3}(b) = a_i q_{m,k}^2(a^{-1})$. Then $q_{n,k}^2(a) = q_{n,k}^2 p_m^{2,3}(b^{-1}) q_{n,m}^2(a_i) = p_n^{2,3} q_{n,m}^3(b^{-1}) q_{n,m}^2(a_i)$. This implies that $q_{n,k}^2(G_k^2)$ is countable because $q_{n,m}^3(G_m^3)$ is countable.

Lemma 2.5. *If $\underline{G} = (G_n, r_n^{n+1})$ is isomorphic to an inverse sequence of countable groups and $\varprojlim^1 \underline{G} = *$, then \underline{G} satisfies the Mittag-Leffler condition.*

Proof. Take an increasing sequence $(n_k)_{k=1}^\infty$ of natural numbers such that

$$r_{n_k}^{n_{k+1}}(G_{n_{k+1}})$$

is countable for each k and $n_1 = 1$.

Define $\underline{H} = (H_n, s_n^{n+1})$ as follows: $H_n = 0$ for $1 \leq n \leq n_2$ and $H_n = \text{im } r_{n_{k-1}}^n$ for $n_k < n \leq n_{k+1}$ and $k \geq 2$,

$s_n^{n+1}: H_{n+1} \rightarrow H_n$ is the inclusion homomorphism if $n_k < m < m+1 \leq n_{k+1}$ for some k and s_n^{n+1} is induced by $r_{n_{k-2}}^{n_{k-1}}$ if $n = n_k$ for some k . Let $f_n: G_n \rightarrow H_n$ be induced by $r_{n_{k-1}}^n$ if $n_k < n \leq n_{k+1}$ and $k \geq 2$ or be the zero homomorphism if $n \leq n_2$.

Then each H_n is countable and $H_n = f_n(G_n)$. By Lemma 2.1 $\varprojlim \underline{H} = *$ and R. Geoghegan [15] has proved that \underline{H} satisfies the Mittag-Leffler condition in such a case (see [16] for the Abelian case). Since \underline{G} and \underline{H} are isomorphic, then \underline{G} satisfies the Mittag-Leffler condition as well.

3. Properties of LC^n -Spaces

From now on by $H_k(X)$ we denote the reduced singular homology group of a space X and by $\check{H}_k(X)$ we denote the reduced Čech homology group of X (all groups are taken with integer coefficients).

In the sequel we shall need the following:

Theorem 3.1. (see Theorem V.2.1 and Propositions II.9.1, II.10.1 in [17]). *Let X and Y be metrizable spaces such that $X \cap Y$ is a closed subset of both X and Y . If X , Y and $X \cap Y$ are LC^n -spaces, then $X \cup Y$ is an LC^n -space. If $X \cup Y$ and $X \cap Y$ are LC^n -spaces, then X and Y are LC^n -spaces.*

The following result of W. Hurewicz [18] is basic in

our considerations.

Theorem 3.2. An LC^1 -space X is an LC^n -space ($n \geq 2$) iff for each $x \in X$ and for each neighborhood U of x in X there is a neighborhood V of x in U such that the inclusion map $i: V \rightarrow U$ induces trivial homomorphism

$$\check{H}_k(i): \check{H}_k(V) \rightarrow \check{H}_k(U)$$

for $k \leq n$.

Theorem 3.3. If X is a connected metrizable LC^n -space, then the natural morphism from $H_k(X)$ to $pro-H_k(X)$ is an isomorphism for $k \leq n$ and an epimorphism for $k = n+1$.

Proof. Take $x \in X$. Theorem 8.7 in [11] says that the natural morphism from $\pi_k(X, x)$ to $pro-\pi_k(X, x)$ is an isomorphism for $k \leq n$ and an epimorphism for $k = n+1$. Hence if $f: \text{Sin}(X, x) \rightarrow \check{C}(X, x)$ is the natural morphism from the geometric realization of the singular complex of (X, x) to the Čech system of (X, x) , then $pro-\pi_k(f)$ is an isomorphism for $k \leq n$ and an epimorphism for $k = n+1$. Now results in [24] and [28] imply that $pro-H_k(f)$ is an isomorphism for $k \leq n$ and an epimorphism for $k = n+1$ which concludes the proof.

Lemma 3.4. Let A be a closed subset of a metrizable space X . If $(A_n)_{n=1}^{\infty}$ is a decreasing sequence of subsets of X such that for any neighborhood W of A in X there is A_n with $A \subset A_n \subset W$, then

- a) $pro-H_k(A)$ is an inverse limit of $(pro-H_k(A_n), pro-H_k(i_n^{n+1}))$ in $pro-Gr$, where $i_n^{n+1}: A_{n+1} \rightarrow A_n$ is the inclusion,
- b) $\check{H}_k(A)$ is the inverse limit of $(\check{H}_k(A_n), \check{H}_k(i_n^{n+1}))$,

c) if $\text{pro-}H_k(A_n)$ is stable for each n , then $\text{pro-}H_k(A)$ is isomorphic to $(\check{H}_k(A_n), \check{H}_k(i_n^{n+1}))$ in pro-Gr .

Proof. It follows from the assumptions that A is an inverse limit of $(A_n, S(i_n^{n+1}))$ in the shape category, where S denotes the shape functor (see [21] and [22]).

A description of inverse limits in pro-categories given in [2] implies that for any functor $F: C \rightarrow D$ the corresponding functor $\text{pro-}F: \text{pro-}C \rightarrow \text{pro-}D$ is continuous i.e. preserves inverse limits. Taking $F = H_k$ we get that Condition a holds. The Condition b follows from Condition a and the fact that the inverse limit functor $\lim_{\leftarrow} : \text{pro-Gr} \rightarrow \text{Gr}$ is continuous. The Condition c is a consequence of Conditions a and b.

4. LC^n -Divisors

Definition. A continuum X is said to be an LC^n -divisor provided for each LC^n -space Y containing X the quotient space Y/X is an LC^n -space.

It is proved in [4] (see also [1]) that

Proposition 4.1. Each FAR-space X is an LC^n -divisor for all n .

We need the following to show that X is an LC^n -divisor iff Y/X is an LC^n -space for some LC^n -space Y containing X (for each space Z we denote by $C(Z)$ a cone over Z).

Lemma 4.2. If $X \subset Y \subset Q$ are subcontinua of the Hilbert cube Q such that Y and Y/X are LC^n -spaces, then Q/X is an LC^n -space.

Proof. By Theorem 3.1 $Q \cup C(Y)$ is an LC^n -space and by Proposition 4.1 $(Q \cup C(Y))/C(X)$ is an LC^n -space. Since

$(Q \cup C(Y))/C(X) = (Q/X) \cup (C(Y)/C(X))$ and $(Q/X) \cap (C(Y)/C(X)) = Y/X$, then by Theorem 3.1 Q/X is an LC^n -space. Thus the proof of Lemma 4.2 is concluded.

If $A \subset X$ are subsets of a compact space Z , then we consider $X \cup C(\bar{A})$ as a space with topology induced from $C(Z)$. Since $C(A)$ is contractible, then the inclusion from $X \cup C(A)$ into the pair $(X \cup C(A), C(A))$ induces isomorphisms of all reduced singular homology groups. By the excision property of singular homology (see [32]) we get that the inclusion from (X, A) to $(X \cup C(A), C(A))$ induces isomorphisms of all reduced singular homology groups. From the exact sequence of homology groups for the pair (X, A) we get the following exact sequence

$$\dots \rightarrow H_k(A) \rightarrow H_k(X) \rightarrow H_k(X \cup C(A)) \rightarrow H_{k-1}(A) \rightarrow \dots$$

Moreover, if $B \subset Y \subset X$ and $B \subset A$, then the diagram

$$\begin{array}{cccccccc} \dots & \rightarrow & H_k(B) & \rightarrow & H_k(Y) & \rightarrow & H_k(Y \cup C(B)) & \rightarrow & H_{k-1}(B) & \rightarrow & \dots \\ & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \\ \dots & \rightarrow & H_k(A) & \rightarrow & H_k(X) & \rightarrow & H_k(X \cup C(A)) & \rightarrow & H_{k-1}(A) & \rightarrow & \dots \end{array}$$

is commutative.

Recall that a continuum X is nearly 1-movable ([25]) provided for each neighborhood U of X in the Hilbert space Q there is a neighborhood V of X in U such that for any loop $f: S^1 \rightarrow V$ and for each neighborhood W of X in Q there is a finite disjoint collection of discs D_i in $\text{Int } \Delta^2$ and an extension of f to

$$\bar{f}: (\Delta^2 - \cup \text{Int } D_i, \cup \partial D_i) \rightarrow (U, U \cap W).$$

Theorem 4.3. If X is a nearly 1-movable continuum such

that $\text{pro-}H_k(X)$ is stable for $k < m$ and satisfies the Mittag-Leffler condition for $k = m$ ($m \geq 1$), then Y/X is an LC^m -space for each LC^m -space Y containing X .

Proof. Let $(U_n)_{n=1}^\infty$ be a decreasing sequence of open neighborhoods of X in Y such that $U_{n+1} \subset \text{cl}(U_{n+1}) \subset U_n$ and $X = \bigcap U_n$. By Theorem 3.3 and Lemma 3.4 the inverse sequence $(H_k(U_n), H_k(i_n^{n+1}))$ is stable for $k < m$ and satisfies the Mittag-Leffler condition for $k = m$.

Fix $n \geq 1$. Then for each $r > n$ there is the following exact sequence

$$0 \rightarrow A_r \rightarrow H_m(U_n) \rightarrow \dots \rightarrow H_k(U_n) \rightarrow H_k(U_n \cup C(U_r)) \rightarrow H_{k-1}(U_r) \rightarrow \dots,$$

where $A_r = \text{im } H_m(i_n^r) \subset H_m(U_n)$. Since $(H_m(U_r), H_m(i_r^S))$

satisfies the Mittag-Leffler condition, then $\varinjlim A_r = *$

by Lemma 2.1. By Lemma 2.3 we get that $\varinjlim (H_k(U_n \cup C(U_r)),$

$H_k(j_r^S)) = *$, where j_r^S is the inclusion map. By Lemma 2.2

the following sequence is exact

$$0 \rightarrow B_n \rightarrow H_m(U_n) \rightarrow \dots \rightarrow H_k(U_n) \rightarrow \check{H}_k(U_n \cup C(X)) \rightarrow \check{H}_{k-1}(X) \rightarrow \dots,$$

where $B_n = \varinjlim A_r$ is the image of $\check{H}_m(X)$ in $H_m(U_n)$. Observe

that the diagram

$$\begin{array}{cccccccc} 0 & \rightarrow & B_p & \rightarrow & H_m(U_p) & \rightarrow & \dots & \rightarrow & H_k(U_p) & \rightarrow & \check{H}_k(U_p \cup C(X)) & \rightarrow & \check{H}_{k-1}(X) & \rightarrow & \dots \\ & & \downarrow & & \downarrow & & & & \downarrow & & \downarrow & & \downarrow & & \downarrow \\ 0 & \rightarrow & B_n & \rightarrow & H_m(U_n) & \rightarrow & \dots & \rightarrow & H_k(U_n) & \rightarrow & \check{H}_k(U_n \cup C(X)) & \rightarrow & \check{H}_{k-1}(X) & \rightarrow & \dots \end{array}$$

is commutative for $p > n$.

By applying Lemma 2.3, 2.4 and 2.5 we infer that

$(\check{H}_k(U_n \cup C(X)), \check{H}_k(1_n^{n+1}))$ satisfies the Mittag-Leffler condi-

tion for $k \leq m$, where 1_n^{n+1} is the inclusion map. Since

$\varinjlim \check{H}_k(U_n \cup C(X)) = \check{H}_k(C(X)) = 0$ (see Lemma 3.4), then

$(\check{H}_k(U_n \cup C(X), \check{H}_k(i_n^{n+1}))$ is isomorphic to the trivial group in pro-Gr for $k \leq m$ by a result of J. Keesling [20] (see also [13]). Hence for each n there exists $p > n$ such that the inclusion l_n^P induces zero homomorphisms of Čech homology groups in dimensions less than or equal to m . Since the projection $U_n \cup C(X) \rightarrow U_n/X$ is a shape equivalence for each n , we get that the inclusion $U_p/X \rightarrow U_n/X$ induces zero homomorphisms on Čech homology groups up to dimension m . By a result in [12] the space Y/X is an LC^1 -space and by the result of W. Hurewicz [18] the space Y/X is an LC^m -space.

Lemma 4.4. *Let X be a subcontinuum of the Hilbert cube Q . If Q/X is an LC^m -space ($m \geq 1$), then X is nearly 1-movable and $\text{pro-}H_k(X)$ is stable for $k < m$ and satisfies the Mittag-Leffler condition for $k = m$.*

Proof. By a result of N. Shrikanhande [31] X is nearly 1-movable (see also [12]).

Take a decreasing sequence $(A_n)_{n=1}^\infty$ of ANR's in Q such that $X = \bigcap A_n$. Then we have the following exact sequence for each n

$$\rightarrow H_{k+1}(Q) \rightarrow H_{k+1}(Q \cup C(A_n)) \rightarrow H_k(A_n) \rightarrow H_k(Q) \rightarrow \dots$$

i.e. the homomorphism from $H_{k+1}(Q \cup C(A_n))$ into $H_k(A_n)$ is an isomorphism for each k . Thus $\text{pro-}H_k(X)$ is isomorphic to $(H_{k+1}(Q \cup C(A_n)), H_{k+1}(i_n^{n+1}))$, where i_n^{n+1} is the inclusion map.

Since the projections $Q \cup C(A_n) \rightarrow Q/A_n$ are homotopy equivalences, $\text{pro-}H_k(X)$ is isomorphic to $(H_{k+1}(Q/A_n), H_{k+1}(p_n^{n+1}))$, where $p_n^{n+1}: Q/A_{n+1} \rightarrow Q/A_n$ is the natural projection.

Now $Q/X = \varprojlim (Q/A_n, p_n^{n+1})$ and therefore $\text{pro-}H_k(X)$ is

isomorphic to $\text{pro-H}_{k+1}(Q/X)$. Since $\text{pro-H}_{k+1}(Q/X)$ is stable for $k < m$ and satisfies the Mittag-Leffler condition for $k = m$ (see Theorem 3.3), the proof of Lemma 4.4 is finished.

As an immediate consequence from Theorem 4.3, Lemma 4.2 and Lemma 4.4 we get

Theorem 4.5. For a continuum X the following conditions are equivalent for $n \geq 1$:

- a) *X is an LC^n -divisor,*
- b) *Y/X is an LC^n -space for some LC^n -space Y containing X,*
- c) *X is nearly 1-movable and $\text{pro-H}_k(X)$ is stable for $k < n$ and satisfies the Mittag-Leffler condition for $k = n$.*

Corollary 4.6. Being an LC^n -divisor is a hereditary shape invariant.

Proof. If $\text{Sh}(Y) \leq \text{Sh}(X)$ and X is an LC^n -divisor, then by Theorem 4.5 X is nearly 1-movable and $\text{pro-H}_k(X)$ is stable for $k < n$ and satisfies the Mittag-Leffler condition for $k = n$. Since $\text{pro-H}_k(Y)$ is dominated by $\text{pro-H}_k(X)$ in pro-Gr for each k, then $\text{pro-H}_k(Y)$ is stable for $k < n$ and satisfies the Mittag-Leffler condition for $k = n$. Now Corollary 4.6 follows from Theorem 4.5 and the fact that being nearly 1-movable continuum is a hereditary shape invariant (see [25]).

Theorem 4.7. Let X be a continuum such that $\text{Fd}(X) = n < +\infty$. Then the following conditions are equivalent:

- a) *X is an ANR-divisor,*
- b) *X is an LC^{n+1} -divisor,*
- c) *X is nearly 1-movable and $\text{pro-H}_k(X)$ is stable for $k \leq n$.*

Proof. It suffices to prove Theorem 4.7 for the case $\dim X = n$ (in view of [29] and Theorem 4.5).

a) \rightarrow b) It follows from Lemma 4.2.

b) \rightarrow a) By a result of Bothe [7] there is an ANR-space Y containing X such that $\dim Y \leq n+1$. Then Y/X is an LC^{n+1} -space and $\dim(Y/X) \leq n+1$. Hence Y/X is an ANR (see [5], p. 122) and by results of Hyman X is an ANR-divisor.

b) \leftrightarrow c) It follows from Theorem 4.5 in view of the fact that $\text{pro-}H_{n+1}(X) = 0$.

Corollary 4.8. In the class of continua of finite fundamental dimension the property of being an ANR-divisor is a hereditary shape invariant. In particular each FANR-space is an ANR-divisor.

Proof. Analogous to the proof of Corollary 4.6.

Example 4.9. We construct an ANR-divisor X whose fundamental dimension is not finite.

For each n let $f_n: S^1 \vee S^n \rightarrow S^1 \vee S^n$ be a map such that $f_n/S^1 = \text{id}$ and $f_n/S^n: S^n \rightarrow S^1 \vee S^n$ is the composition of maps $g_n: S^n \rightarrow \bigvee_{i=1}^{\infty} S_i^n$ and $e_n: \bigvee_{i=1}^{\infty} S_i^n \rightarrow S^1 \vee S^n$, where $\pi_n(e_n)$ is an isomorphism and g_n represents the difference $[S_1^n] - [S_2^n]$ of two generators of $\pi_n(\bigvee_{i=1}^{\infty} S_i^n)$.

Then $H_k(f_n) = 0$ for each k and the induced map

$f'_n: (S^1 \vee S^n)/S^1 \rightarrow (S^1 \vee S^n)/S^1$ is homotopically trivial.

Let $X_n = \bigvee_{k=1}^n S^k$ and let $h_n^{n+1}: X_{n+1} \rightarrow X_n$ be defined by $h_n^{n+1}(x) = f_k(x)$ for $x \in S^k$, $k \leq n$, and S^{n+1} is mapped onto the base point.

Let $X = \varprojlim (X_n, h_n^{n+1})$. Then $S^1 \subset X$ and X/S^1 is an FAR. Since FAR's are ANR-divisors we infer by a result of Hyman [19] that X is an ANR-divisor.

Observe that $\text{Fd}(X)$ is not finite because finiteness of $\text{Fd}(X)$ would imply triviality of $\pi_n(f_n)^k$ for $n = \text{Fd}(X) + 1$ and some k .

Remark. Example 4.9 is constructed in the spirit of an example in [12].

Analogous to the corresponding results for ANR-divisors in [19] one can prove the following

Theorem 4.10. Let X and Y be continua. If X , Y and $X \cap Y$ are LC^n -divisors, then $X \cup Y$ is an LC^n -divisor. If $X \cup Y$ and $X \cap Y$ are LC^n -divisors, then X and Y are LC^n -divisors. If $X \subset Y$ and X and Y/X are LC^n -divisors, then Y is an LC^n -divisor.

The author is grateful to Jack Segal for his help during the preparation of this paper.

References

- [1] S. Armentrout, *Decompositions and absolute neighborhood retracts*, Lecture Notes in Math. 438, Springer, New York, 1975.
- [2] M. Artin and B. Mazur, *Etale homotopy theory*, Lecture Notes in Math. 100, Springer, 1969.
- [3] M. F. Atiyah and G. Segal, *Equivariant K-theory and completion*, J. Diff. Geometry 3 (1969), 1-18.
- [4] S. Bogatyĭ, *On a Vietoris theorem in the category of homotopies and a problem of Borsuk*, Fund. Math. 84 (1974), 209-228.

- [5] K. Borsuk, *Theory of retracts*, Monografie Matematyczne 44, Warszawa, 1967.
- [6] _____, *Theory of shape*, Monografie Matematyczne 59, Warszawa, 1975.
- [7] H. Bothe, *Eine Einbettung m -dimensionaler Mengen in einen $(m+1)$ -dimensionalen absoluten Retrakt*, Fund. Math. 51 (1962), 209-224.
- [8] A. K. Bousfield and D. M. Kan, *Homotopy limits, completions and localizations*, Lecture Notes in Math. 304 (1972), Springer, New York.
- [9] Z. Čerin, *C_p -movable at infinity spaces, compact ANR-divisors and property UVW^n* , Publ. de l' Institut Math. 23 (1978), 53-65.
- [10] T. A. Chapman, *On some applications of infinite-dimensional manifolds to the theory of shape*, Fund. Math. 76 (1972), 181-193.
- [11] J. Dydak, *The Whitehead and Smale theorems in shape theory*, Dissertationes Mathematicae 156 (1979), 1-50.
- [12] _____, *Some properties of nearly 1-movable continua*, Bull. Ac. Pol. Sci. 25 (1977), 685-689.
- [13] _____, *An algebraic condition characterizing FANR-spaces*, Bull. Ac. Pol. Sci. 24 (1976), 501-503.
- [14] D. A. Edwards and H. M. Hastings, *Čech and Steenrod homotopy theories with applications to geometric topology*, Lecture Notes in Math. 542, Springer, New York, 1976.
- [15] R. Geoghegan, *A note on the vanishing of \lim^1* , (preprint).
- [16] B. I. Gray, *Spaces of the same n -type for all n* , Topology 5 (1966), 241-243.
- [17] S.-T. Hu, *Theory of retracts*, Wayne University Press, Detroit, 1965.
- [18] W. Hurewicz, *Homotopie, Homologie, und lokaler Zusammenhang*, Fund. Math. 25 (1935), 467-485.
- [19] D. M. Hyman, *ANR-divisors and absolute neighborhood contractibility*, Fund. Math. 62 (1968), 61-73.
- [20] J. E. Keesling, *On the Whitehead theorem in shape theory*, Fund. Math. 92 (1976), 247-253.

- [21] G. Kozłowski and J. Segal, *Local behavior and the Vietoris and Whitehead theorems in shape theory*, *Fund. Math.* 99 (1978), 210-219.
- [22] S. Mardešić, *Shapes for topological spaces*, *Gen. Top. Appl.* 3 (1973), 265-282.
- [23] _____ and J. Segal, *Shape of compacta and ANR-systems*, *Fund. Math.* 72 (1971), 41-59.
- [24] S. Mardešić, *On the Whitehead theorem in shape theory II*, *Fund. Math.* 91 (1976), 93-103.
- [25] D. R. McMillan, *One-dimensional shape properties and three manifolds*, *Studies in Topology*, Academic Press, 1975, pp. 367-381.
- [26] J. Milnor, *On axiomatic homology theory*, *Pacific J. Math.* 12 (1966), 337-341.
- [27] K. Morita, *On shapes of topological spaces*, *Fund. Math.* 86 (1975), 251-259.
- [28] _____, *The Hurewicz and the Whitehead theorems in shape theory*, *Sci. Rep. of the Tokyo Kyoiku Daigaku, Sec. A*, 12 (1974), 246-258.
- [29] S. Nowak, *Some properties of the fundamental dimension*, *Fund. Math.* 85 (1974), 211-227.
- [30] R. H. Overton, *Čech homology for movable compacta*, *Fund. Math.* 77 (1973), 241-251.
- [31] N. Shrikhande, *Homotopy properties of decomposition spaces*, *Abstract 75T-638*, *Notices AMS*, Apr. 1975.
- [32] E. H. Spanier, *Algebraic topology*, McGraw-Hill, New York, 1966.

University of Washington

Seattle, Washington 98195

and

Institute of Mathematics of the Polish Academy of Sciences

Warsaw, ul. Śniadeckich 8, Poland