
TOPOLOGY PROCEEDINGS



Volume 6, 1981

Pages 329–334

<http://topology.auburn.edu/tp/>

INSERTION OF A CONTINUOUS FUNCTION AND $X \times I$

by

ERNEST P. LANE

Topology Proceedings

Web: <http://topology.auburn.edu/tp/>

Mail: Topology Proceedings
Department of Mathematics & Statistics
Auburn University, Alabama 36849, USA

E-mail: topolog@auburn.edu

ISSN: 0146-4124

COPYRIGHT © by Topology Proceedings. All rights reserved.

INSERTION OF A CONTINUOUS FUNCTION AND $X \times I$

Ernest P. Lane¹

If g and f are real-valued functions defined on a topological space X such that $g < f$ (i.e., $g(x) < f(x)$ for each x in X), consider the problem of finding necessary and sufficient conditions in order for there to be a continuous function h defined on X such that $g < h < f$. Such conditions seem to have characteristics that keep them from being applicable in some cases. For example, the condition given in Theorem 3.11 of [2] requires that $-g$ must possess the same property as f and that the classes to which g and f belong are closed under sum, supremum, and infimum. The condition given in Theorem 2.1 of [5] places a restriction on $f - g$. This paper gives a necessary and sufficient condition that avoids these problems, but it is in terms of subsets of $X \times I$.

All functions considered are assumed to map into the interval $(0,1)$; I denotes the interval $[0,1]$. (In most situations there is no loss in generality in replacing \mathbb{R} with $(0,1)$.) It is therefore convenient to let $C(X)$ represent the set of all continuous functions from X into $(0,1)$. The characterization that is given below is in terms of the following subsets of $X \times I$: If k maps X into $(0,1)$, let

¹This work was supported by a released teaching load research grant from Appalachian State University.

$$L_k = \{(x,t) \in X \times I: t \leq k(x)\}$$

and

$$U_k = \{(x,t) \in X \times I: t \geq k(x)\}.$$

It is observed in [6] that if k is continuous then L_k and U_k are zero sets in $X \times I$; this will be used below in a proof. It is convenient to use the following terminology. If k is a real-valued function then k is *z-lower semicontinuous* (zlsc) [respectively, *z-upper semicontinuous* (zusc)] if for each real number r , $\{x \in X: k(x) \leq r\}$ [respectively, $\{x \in X: k(x) \geq r\}$] is a zero set. (These functions have been used in [1] and in [8].)

Proposition 1. If g and f are functions from X into $(0,1)$ such that $g < f$ and such that L_g and U_f are completely separated in $X \times I$, then there is a continuous function h on X such that $g < h < f$.

Proof. If g and f are functions from X into $(0,1)$ such that $g < f$ and such that L_g and U_f are completely separated, let F be a continuous function from $X \times I$ into I such that if (x,t) is in L_g then $F(x,t) = 0$ and if (x,t) is in U_f then $F(x,t) = 1$. Define functions f_1 and g_1 from X into $(0,1)$ as follows:

$$f_1(x) = \inf\{t \in I: (\{x\} \times [t,1]) \subset F^{-1}(\{1\})\}$$

and

$$g_1(x) = \sup\{t \in I: (\{x\} \times [0,t]) \subset F^{-1}(\{0\})\}.$$

Since $L_g \subset F^{-1}(\{0\})$ and $U_f \subset F^{-1}(\{1\})$, $g \leq g_1$ and $f_1 \leq f$.

Since F is continuous it is easy to show that $g_1 < f_1$.

Also, g_1 is zusc and f_1 is zlsc; the outline of an argument to show that f_1 is zlsc follows: If r is any real number

in $(0,1)$, define a function F_r from X into I by

$$F_r(x) = \inf\{F(x,t) : r \leq t \leq 1\}.$$

Then F_r is a continuous function from X into I . Next observe that

$$\{x \in X : f_1(x) \leq r\} = \{x \in X : F_r(x) = 1\}:$$

For if $f_1(x) \leq r$ then $\{x\} \times [r,1] \subset F^{-1}(\{1\})$ and hence $F_r(x) = 1$. Conversely, if $F_r(x) = 1$ then $F(x,t) = 1$ for all t such that $r \leq t \leq 1$ and consequently $\{x\} \times [r,1] \subset F^{-1}(\{1\})$; thus $f_1(x) \leq r$. Since F_r is continuous, $\{x \in X : F_r(x) = 1\}$ is a zero set. Since $\{x \in X : f_1(x) \leq r\}$ is a zero set for any r in $(0,1)$, it follows from the definition that f_1 is zlsc. A similar argument shows that g_1 is zusc. Thus $g \leq g_1 \leq f_1 \leq f$, g_1 is zusc and f_1 is zlsc; by Proposition 6.1 of [1], there is a continuous function h defined on X such that $g_1 < h < f_1$. Hence $g < h < f$ and this concludes the proof.

Remark. In the above proof the zero set $F^{-1}(\{1\})$ in $X \times I$ is used to define a zlsc function on X ; in the following it is observed that a zlsc function on X corresponds to a zero set in $X \times I$. Tong [9] proves that a space X is perfectly normal if and only if each lower semicontinuous function on X is a pointwise limit of an increasing sequence of continuous functions. His proof is inductive and involves the complete separation of certain subsets of X . It is straightforward to observe that this proof yields the following result: If f is a zlsc function on any topological space X , there is an increasing sequence of continuous functions on X whose pointwise limit is f . (The converse of

this is immediate from the definition of zlsc.) Consequently, if f is zlsc then U_f is a countable intersection of zero sets, and hence U_f is a zero set in $X \times I$.

The main result of the paper follows from Proposition 1.

Theorem 1. Let $L(X)$ and $U(X)$ be classes of functions from X into $(0,1)$ such that $C(X) \subset L(X)$ and $C(X) \subset U(X)$.

The following are equivalent:

(i) For any g in $U(X)$ and any f in $L(X)$ such that $g < f$ there is an h in $C(X)$ such that $g < h < f$.

(ii) For any g in $U(X)$ and any f in $L(X)$ such that $g < f$ then L_g and U_f are completely separated in $X \times I$.

Proof. That (ii) implies (i) follows from Proposition 1 (and does not require the hypothesis that $C(X) \subset L(X)$ and $C(X) \subset U(X)$). Conversely, suppose that (i) is satisfied. If $g \in U(X)$, $f \in L(X)$ and $g < f$, then by (i) there exists h in $C(X)$ such that $g < h < f$. Since $C(X) \subset L(X)$, apply (i) to g and h ; there exists k in $C(X)$ such that $g < k < h$. Since k and h are continuous L_k and U_h are zero sets; since $k < h$ it follows that L_k and U_h are disjoint. Since $L_g \subset L_k$ and $U_f \subset U_h$, the sets L_g and U_f are completely separated.

An application of this theorem is given below that involves normal semicontinuous functions. If f_* [respectively, f^*] denotes the lower [respectively, upper] limit function of f then f is normal lower semicontinuous (nlsc) in case $f = (f^*)_*$ and f is normal upper semicontinuous (nusc) in case $f = (f_*)^*$. Since any continuous function is nlsc and nusc, the following is immediate from Theorem 1.

Corollary 1. Let $L(X)$ [respectively, $U(X)$] denote the nlsc [respectively, nusc] functions from X into $(0,1)$. The following are equivalent:

(i) For any g in $U(X)$ and any f in $L(X)$ such that $g < f$ there is an h in $C(X)$ such that $g < h < f$.

(ii) For any g in $U(X)$ and any f in $L(X)$ such that $g < f$ then L_g and U_f are completely separated in $X \times I$.

The proof of Theorem 3.2 of [6] shows that if X is any space such that g nusc, f nlsc and $g < f$ imply that L_g and U_f are completely separated in $X \times I$, then there is a continuous function h on X such that $g < h < f$. The proof of this uses the result that for g nusc and f nlsc then L_g and U_f are regular closed sets in $X \times I$. The general converse of Theorem 3.2 of [6] remains open: Does condition (i) of Corollary 1 (for g nusc and f nlsc) imply that arbitrary disjoint regular closed subsets of $X \times I$ are completely separated?

If $X \times I$ satisfies the equivalent conditions of Theorem 1 for various classes of functions, the corresponding characterization of the space X is known. If $L(X)$ (respectively, $U(X)$) is the class of lower (respectively, upper) semicontinuous functions on X , the conditions of Theorem 1 are equivalent to X is normal and countably paracompact. (This follows from Theorem 4 of [3] or from Theorem 2 of [4].) If $L(X)$ (respectively, $U(X)$) is the class of upper (respectively, lower) semicontinuous functions on X , then the conditions of Theorem 1 are equivalent to X is an extremally disconnected P -space that satisfies Baire's condition. (See Proposition 6.11 of [1].) For an

extensive list of corresponding characterizations of X for various cases of the classes $L(X)$ and $U(X)$, see Theorem 4.2 of [7].

References

1. J. Blatter and G. L. Seever, *Interposition of semi-continuous functions by continuous functions*, *Analyse Fonctionnelle et Applications (Comptes Rendus due Colloque d'Analyse, Rio de Janeiro 1972)*, Hermann, Paris, 1975, 27-51.
2. _____, *Interposition and lattice cones of functions*, *Trans. Amer. Math. Soc.* 222 (1976), 65-96.
3. C. H. Dowker, *On countably paracompact spaces*, *Can. J. Math.* 3 (1951), 219-224.
4. M. Katětov, *On real-valued functions in topological spaces*, *Fund. Math.* 38 (1951), 85-91.
5. E. P. Lane, *Insertion of a continuous function*, *Pacific J. Math.* 66 (1976), 181-190.
6. _____, *PM-normality and the insertion of a continuous function*, *Pacific J. Math.* 82 (1979), 155-162.
7. _____, *Insertion of a continuous function*, *Topology Proc.* 4 (1979), 463-478.
8. M. H. Stone, *Boundedness properties in function lattices*, *Can. J. Math.* 1 (1949), 176-186.
9. H. Tong, *Some characterizations of normal and perfectly normal spaces*, *Duke Math. J.* 19 (1952), 289-292.

Appalachian State University

Boone, NC 28608