

---

# TOPOLOGY PROCEEDINGS



Volume 8, 1983

Pages 45–50

---

<http://topology.auburn.edu/tp/>

## G-SYSTEMS

by

GORDON G. JOHNSON

---

### Topology Proceedings

**Web:** <http://topology.auburn.edu/tp/>

**Mail:** Topology Proceedings  
Department of Mathematics & Statistics  
Auburn University, Alabama 36849, USA

**E-mail:** [topolog@auburn.edu](mailto:topolog@auburn.edu)

**ISSN:** 0146-4124

COPYRIGHT © by Topology Proceedings. All rights reserved.

## G-SYSTEMS

Gordon G. Johnson

It is well known [1], that if  $X$  is a real finite dimensional inner product space and  $S$  is a closed subset of  $X$ , then  $S$  is convex only in case every point in  $X$  has a unique nearest point in  $S$ .

The following definition was motivated by the search for the answer to the question: Is there a real inner product space  $X$  containing a closed nonconvex subset  $S$ , such that each point in  $X$  has a unique nearest point in  $S$ ?

In all that follows it is to be understood that  $X$  is a real inner product space.

*Definition.* By a  $G$ -system for  $X$  we mean an ordered pair  $(M, S)$  where  $M$  is pairwise disjoint, closed and convex set collection filling up  $X$  and  $S$  is a subset of  $X$  such that each set in  $M$  contains exactly one point of  $S$ , and if  $m_1$  and  $m_2$  are two sets in  $M$  with points  $s_1$  and  $s_2$  from  $S$  respectively, then each point  $p$  in  $m_1$  is closer to  $s_1$  than  $p$  is to  $s_2$ .

Suppose  $S$  is a closed and convex subset of the finite dimensional space  $X$  and  $s$  is a point in  $S$ . Let  $K$  be the set to which  $x$  belongs only in case  $x$  is in  $X$  and  $s$  is the nearest point in  $S$  to  $x$ . Then  $K$  is a closed and convex set, and the collection of all such  $K$  is a decomposition of  $X$  into pairwise disjoint, closed and convex sets, each containing exactly one point of  $S$ .

Let  $X$  be the euclidean plane,  $M$  the set of all vertical lines and  $S$  the  $x$ -axis. Then  $(M,S)$  is a  $G$ -system for  $X$ . If  $S'$  is any horizontal line, then  $(M,S')$  is a  $G$ -system for  $X$ . However, if  $L$  is a non-horizontal line, then  $(M,L)$  is not a  $G$ -system for  $X$ .

*Question 1.* If each of  $(M,S)$  and  $(M,S')$  is a  $G$ -system for  $X$ , then how are  $S$  and  $S'$  related?

*Theorem 1.* If  $(M,S)$  is a  $G$ -system for  $X$ , then  $S$  is closed.

*Proof.* Suppose  $p$  is a point in  $X$ . Then there is a unique set  $m$  in  $M$  which contains  $p$ . Let  $s$  be that point in  $S$  which is in  $m$ . If  $p \neq s$ , then every point in  $S$  different from  $s$  is farther from  $p$ , than  $p$  is from  $s$ , and hence  $p$  is not a limit point of  $S$ . Hence  $S$  is closed.

*Theorem 2.* If  $(M,S)$  is a  $G$ -system for  $X$  and  $S$  is non-degenerate, then  $S$  is uncountable.

*Proof.* Suppose  $S$  is nondegenerate and  $u$  and  $v$  are two points in  $S$ . Let  $[u,v]$  denote the line interval with end-points  $u$  and  $v$ . For each point  $p$  in  $[u,v]$  there is a set  $m_p$  in  $M$  which contains  $p$ . Notice that  $m_p \cap [u,v]$  is a closed set and if  $p$  and  $q$  are two points in  $[u,v]$ , then either  $m_p = m_q$  or  $m_p$  does not intersect  $m_q$ . We also have that  $\bigcup_{p \in [u,v]} [m_p \cap [u,v]] = [u,v]$ .

Therefore  $\{m_p : p \in [u,v]\}$  is uncountable, because no interval is the union of countably many pairwise disjoint closed sets. Hence  $S$  is uncountable.

If  $X$  is separable, then except for at most a countable subset of  $S$ , each point of  $S$  is a limit point of  $S$ , and hence is a boundary point of that set in  $M$  to which it belongs. This then leads to the following theorem.

*Theorem 2.* *If  $(M,S)$  is a  $G$ -system for  $X$  and  $S$  is nondegenerate, then each point in  $S$  is a boundary point of that set in  $M$  to which it belongs, and hence is a limit point of  $S$ .*

*Proof.* Suppose  $S$  is nondegenerate and that there is a point  $s$  in  $S$  which is not a boundary point of the set  $m$  in  $M$  to which it belongs. Then  $s$  is an interior point of  $m$  and therefore there is a number  $d > 0$  such that  $S_d(s) = \{x: ||x - s|| \leq d\}$  is a subset of  $m$  and if  $\epsilon > 0$ , then there is a point  $q$  not in  $m$  such that  $||s - q|| \leq d + \epsilon$ .

Let  $b$  be a boundary point of  $m$  and  $C$  be the union of all line intervals having one endpoint  $b$  and the other endpoint in  $S_d(s)$ . Since  $m$  is convex,  $C$  is a subset of  $m$ .

Let  $C^+ = \bigcup_{\substack{c \in C \\ c \neq b}} S_{||c-s||}(c)$  and  $p$  be a point on the boundary

of  $S_{||b-s||}(b)$ , different from  $2b - s$  and not in  $m$ . Let  $w$  be the point common to the boundary of  $S_{||b-s||}(b)$  and boundary of  $S_d(s)$  lying in the plane determined by  $s$ ,  $b$  and  $p$ , and closest to  $p$ .

The line interval  $[w,b]$  is a subset of  $C$  and hence for each  $t$ ,  $0 \leq t \leq 1$ ,  $(1 - t)w + tb = r$  is in  $C$ . There is a number  $t$  in  $[0,1]$  such that  $S_{||r-s||}(r)$  contains  $p$ , and hence  $p$  is in  $C^+$ . Notice then that  $S_{||b-s||}(b) - \{2b - s\}$  is a subset of  $C^+$ ,  $2b - s$  is not in  $C^+$ , and no point of  $S$  different from  $s$  is in  $C^+$ .

Let  $R = \{(1 - t)s + tb : t \geq 0\}$  and let  $\{q_i\}_{i=1}^{\infty}$  be a sequence such that

1. for each  $i = 1, 2, \dots$   $q_i$  is in  $R - m$
2.  $\lim_{i \rightarrow \infty} q_i = b$ , and
3. if for each  $i$ ,  $m_i$  is that set in  $M$  containing  $q_i$ ,

then  $m_i \neq m_j$  if  $i \neq j$ .

For each  $i$ , let  $s_i$  be that point in  $S$  that is in  $m_i$ .

Then

$$\lim_{i \rightarrow \infty} \|s_i - q_i\| = \|s - b\|,$$

$$\lim_{i \rightarrow \infty} \|q_i - b\| = 0,$$

and if  $E > 0$ , then there is  $N > 0$  such that if  $n > N$ , then  $\|s_n - s\| \leq 2\|s - b\| + E$ .

Hence the point sequence  $\{s_i\}_{i=1}^{\infty}$  must tend to the boundary of  $S_{\|s-b\|}(b)$ . Recall that no point of  $S$  different from  $s$  is in  $C^+$ . Therefore,  $\lim_{i \rightarrow \infty} s_i = 2b - s$  and since  $S$  is closed,  $2b - s$  is in  $S$ . We then have a contradiction since  $\|s - b\| = \|(2b - s) - b\|$  and  $2b - s \neq s$ .

It is easy to show that if  $(M, S)$  is a  $G$ -system for  $X$  and  $S$  is nondegenerate and connected, then each point in  $S$  is a boundary point of that set in  $M$  to which it belongs. This then raises the following question.

*Question 2.* If  $(M, S)$  is a  $G$ -system for  $X$ , then is  $S$  connected?

The proof for theorem 2 also raises the following question.

*Question 3.* If  $(M,S)$  is a G-system for  $X$ ,  $m$  is a set in  $M$  which contains two points,  $s$  is that point in  $S$  which is in  $m$  and  $p$  is a point in  $m$  different from  $s$ , then is  $\{(1 - t)s + tp: t \geq 0\}$  a subset of  $m$ ?

*Theorem 4.* Suppose  $(M,S)$  is a G-system for  $X$ ,  $\{s_i\}_{i=1}^\infty$  is a convergent sequence from  $S$ , and  $\{w_i\}_{i=1}^\infty$  is a convergent sequence such that for each  $i$ ,  $w_i$  and  $s_i$  belong to the same set in  $M$ . If  $\lim_{i \rightarrow \infty} s_i = s$  and  $\lim_{i \rightarrow \infty} w_i = w$ , then  $w$  and  $s$  belong to the same set in  $M$ .

*Proof.* Let  $m$  be that set in  $M$  which contains  $s$ . And  $\bar{m}$  that set in  $M$  which contains  $w$ . Suppose  $\bar{m} \neq m$ , and let  $\bar{s}$  be that point in  $S$  which is in  $\bar{m}$ . Then

$$d = ||s - w|| - ||s - \bar{w}|| > 0.$$

Now

$$\begin{aligned} ||w_i - \bar{s}|| &\leq ||w_i - w|| + ||w - \bar{s}|| = \\ &||w_i - w|| + ||s - w|| - d < ||w_i - w|| \\ &+ ||s_i - s|| + ||s_i - w|| - d. \end{aligned}$$

Choose  $N > 0$  such that if  $n > N$ , then  $||w_n - w|| < d/4$  and  $||s_n - s|| < d/4$ .

Hence

$$\begin{aligned} ||w_n - \bar{s}|| &< d/4 + d/4 + ||s_n - w|| - d \\ &< ||s_n - w_n|| + ||w_n - w|| - d/2 \\ &< ||s_n - w_n|| - d/4 \end{aligned}$$

Therefore  $||w_n - \bar{s}|| < ||s_n - w_n||$ , which is a contradiction.

*Question 4.* If  $(M,S)$  is a G-system for  $X$  and  $\{w_i\}_{i=1}^\infty$  is a convergent sequence in  $X$  must  $\{s_i\}_{i=1}^\infty$  be a convergent sequence where for each  $i$ ,  $s_i$  and  $w_i$  belong to the same set in  $M$ ?

*Theorem 5.* Suppose  $(M,S)$  is a  $G$ -system for  $X$  such that if  $m_1$  and  $m_2$  are two sets in  $M$  with points  $s_1$  and  $s_2$  from  $S$  respectively, then the unique nearest point for  $s_2$  in  $m_1$  is  $s_1$  and the unique nearest point for  $s_1$  in  $m_2$  is  $s_2$ . Then  $S$  is convex.

*Proof.* Let  $s_1$  and  $s_2$  be two points in  $S$  and  $q$  a point between  $s_1$  and  $s_2$  on the line interval  $[s_1, s_2]$ . If  $s_1$  is in  $m_1$  and  $s_2$  is in  $m_2$ , then  $q$  is not in  $m_1$  or  $m_2$  since  $q$  is closer to  $s_1$  than  $s_2$  is to  $s_1$  and  $q$  is closer to  $s_2$  than  $s_1$  is to  $s_2$ . Let  $m$  be that set in  $M$  which contains  $q$  and  $s$  that point in  $S$  which is in  $m$ . Then  $||s_1 - s|| + ||s_1 - s_2|| \geq ||s_1 - q|| + ||q - s_2||$  and equality holds only in case  $s$  is in  $[s_1, s_2]$ . Hence  $s = q_1$  and therefore  $S$  is convex.

Notice that in order to construct a closed nonconvex set have the unique nearest point property, the hypothesis of theorem 5 must not hold.

### References

1. F. A. Valentine, *Convex sets*, McGraw-Hill, New York, 1969.
2. R. L. Moore, *Foundations of point set theory*, Am. Math. Soc. Colloq. Publs., Vol. VIII, 1932.

University of Houston

Houston, Texas 77004

and

Emory University

Atlanta, Georgia 30322