

---

# TOPOLOGY PROCEEDINGS



Volume 8, 1983

Pages 303–314

---

<http://topology.auburn.edu/tp/>

## AN ADMISSIBLE CONDITION FOR CONTRACTIBLE HYPERSPACES

by

CHOON JAI RHEE AND TOGO NISHIURA

---

### Topology Proceedings

**Web:** <http://topology.auburn.edu/tp/>

**Mail:** Topology Proceedings  
Department of Mathematics & Statistics  
Auburn University, Alabama 36849, USA

**E-mail:** [topolog@auburn.edu](mailto:topolog@auburn.edu)

**ISSN:** 0146-4124

COPYRIGHT © by Topology Proceedings. All rights reserved.

## AN ADMISSIBLE CONDITION FOR CONTRACTIBLE HYPERSPACES

Choon Jai Rhee and Togo Nishiura

Let  $X$  be a nonvoid metric continuum. Denote by  $2^X$  and  $C(X)$  the hyperspaces of nonempty closed subsets and subcontinua of  $X$  respectively and endow each with the Hausdorff metric  $H$ .

In 1938 Wojkyslawski proved that  $2^X$  is contractible if  $X$  is locally connected [11]. In 1942 Kelley [1] proved that the contractibility of  $2^X$  is equivalent to the contractibility of  $C(X)$ . Furthermore, he introduced a sufficient condition, namely property (3.2), for the contractibility of the hyperspaces of metric continua. In [5], a necessary condition, namely admissibility, is given for a space whose hyperspace is contractible. It was also proven that the contractibility of the hyperspace  $C(X)$  is equivalent to the existence of a continuous fiber map on  $X$  into the hyperspace  $C^2(X)$  of subcontinua of  $C(X)$  for the class of metric continua with property  $c$  (abbreviated as  $c$ -space). In the present paper we show that if  $f: X \rightarrow Y$  and  $g: Y \rightarrow X$  are continuous such that  $f \circ g$  is homotopic to the identity map  $\text{id}_Y$  on  $Y$ , and if  $X$  is a  $c$ -space then  $Y$  is also a  $c$ -space. Hence if  $X$  and  $Y$  are homotopically equivalent, then  $X$  is a  $c$ -space if and only if  $Y$  is. We also show that the product space  $X \times Y$  is a  $c$ -space if and only if both  $X$  and  $Y$  are  $c$ -spaces. Many corollaries to the above results are also given which are generalizations of results in [4].

Throughout the paper, the symbols  $I$  and  $\mu$  will be reserved for the closed interval and a Whitney map [10] with  $\mu(X) = 1$  respectively. Note that  $\mu(X) = 1$  necessarily requires  $X$  to be nondegenerate, a condition which we will assume whenever required without explicitly stating so.

### 1. Preliminaries

We collect in this section some definitions and known facts and prove a new lemma. Let  $X$  be a nonvoid metric continuum.

A map  $H: X \times I \rightarrow C(X)$  is *increasing* if  $h(x,t) \subset h(x,t')$  for  $t \leq t'$  and  $x \in X$ . A *contraction of  $X$  in  $C(X)$*  is a continuous homotopy  $h: X \times I \rightarrow C(X)$  such that, for each  $x \in X$ ,  $h(x,0) = \{x\}$  and  $h(x,1) = A$ . A contraction of  $X$  in  $2^X$  is analogously defined.

*Theorem 1.1 [1]. The following statements are equivalent.*

1. *A contraction of  $X$  in  $C(X)$  exists.*
2.  *$2^X$  is contractible.*
3.  *$C(X)$  is contractible.*

*Theorem 1.2 [1]. If  $C(X)$  is contractible then an increasing contraction of  $X$  in  $C(X)$  exists.*

The contractibility of  $C(X)$  implies the contractibility of  $C^2(X)$ , by Theorem 1.1. Also, since the union map from  $C^2(X)$  onto  $C(X)$  is a retraction, the contractibility of  $C^2(X)$  implies the contractibility of  $C(X)$ . Thus we have the following.

*Theorem 1.3 [1].*  $C(X)$  is contractible if and only if the hyperspace  $C^2(X)$  of subcontinua of  $C(X)$  is contractible.

We recall the definition of the Hausdorff metric  $H$  on  $2^X$ . For  $A, B \in 2^X$ ,

$$H(A, B) = \max\{\max_{a \in A} d(a, B), \max_{b \in B} d(b, A)\},$$

where  $d(x, A)$  is the distance from  $x$  to  $A$ .

*Lemma 1.4.* If  $A, B, C, D \in 2^X$  then

$$H(A \cup B, C \cup D) \leq \max\{H(A, C), H(B, D)\}.$$

*Proof.* Let  $\eta > \max\{H(A, C), H(B, D)\}$ . Then  $C \cup D \subset \{x \mid d(x, A) < \eta\} \cup \{x \mid d(x, B) < \eta\} = \{x \mid d(x, A \cup B) < \eta\}$ . Also  $A \cup B \subset \{x \mid d(x, C \cup D) < \eta\}$ .

Let  $X$  be a nonvoid continuum. We now define an admissibility condition [5] and prove some propositions. For  $x \in X$ , let  $F(x) = \{A \in C(X) \mid x \in A\}$ , and for  $(x, t) \in X \times I$ ,  $F_t(x) = F(x) \cap \mu^{-1}(t)$ . An element  $A \in F(x)$  is said to be *admissible at  $x$*  if, for each  $\varepsilon > 0$ , there is  $\delta > 0$  such that each  $y$  in the  $\delta$ -neighborhood of  $x$  has an element  $B \in F(y)$  such that  $H(A, B) < \varepsilon$ . For each  $x \in X$ , the collection  $A(x) = \{A \in F(x) \mid A \text{ is admissible at } x\}$  is called the *admissible fiber at  $x$* . We say that  $X$  is *admissible* if  $A_t(x) = A(x) \cap \mu^{-1}(t)$  is nonempty for each  $(x, t) \in X \times I$ .

*Proposition 1.5.* If  $A \in A(\xi)$  and  $B \in A(x)$  and  $\xi \in A \cap B$  then  $A \cup B \in A(x)$ . Hence, if  $A_i \in A(x)$ ,  $i = 1, 2, \dots, n$ , then  $\bigcup_{i=1}^n A_i \in A(x)$ .

*Proof.* Let  $\varepsilon > 0$ . Since  $A \in A(\xi)$ , there is  $\tau < \varepsilon$  where each point  $y$  of the  $\tau$ -neighborhood  $V$  of  $\xi$  has an element  $C \in F(y)$  such that  $H(A, C) < \varepsilon$ . Since  $B \in A(x)$  there is

$\delta > 0$  such that each point  $z$  of the  $\delta$ -neighborhood  $W$  of  $x$  has an element  $D \in F(z)$  such that  $H(B, D) < \tau$ . One sees that  $\xi \in B$  and  $H(B, D) < \tau$  imply  $V \cap D \neq \emptyset$ . Hence, for each  $z \in W$  there are  $D \in F(z)$ ,  $y \in V \cap D$  and  $C \in F(y)$  such that  $H(A, C) < \varepsilon$ ,  $H(B, D) < \varepsilon$  and  $C \cup D \in F(z)$ . By Lemma 1.4, we have  $H(A \cup B, C \cup D) \leq \max\{H(A, C), H(B, D)\} < \varepsilon$ , and the proposition is proved.

*Proposition 1.6.* For each  $x \in X$  its admissible fiber  $A(x)$  is closed in  $C(X)$ ,  $\{x\} \in A(x)$  and  $X \in A(x)$ .

*Proof.* Suppose  $A_n$ ,  $n = 1, 2, \dots$ , is a sequence in  $A(x)$  which converges to  $A$  in  $C(X)$ . Obviously,  $A \in F(x)$ . Let  $\varepsilon > 0$ . There is a positive integer  $N$  such that  $H(A, A_N) < \varepsilon/2$ . Since  $A_N \in A(x)$ , there is a  $\delta$ -neighborhood  $V$  of  $x$  such that each point  $y$  of  $V$  has an element  $B \in F(y)$  such that  $H(A_N, B) < \varepsilon/2$ . From  $H(A, B) \leq H(A, A_N) + H(A_N, B) < \varepsilon$ , we have  $A \in A(x)$  and hence  $A(x)$  is closed. The remaining parts of the proposition are obvious.

We note that since  $C(X)$  is compact [1],  $A(x)$  is compact.

*Proposition 1.7.* Let  $B \in F(x)$  and  $C = \cup\{A \in A(x) \mid A \subset B\}$  then  $C \in A(x)$ .

*Proof.* First we prove  $C$  is a subcontinuum of  $X$ . Clearly  $C$  is connected and  $x \in C$ . Let  $x_n$ ,  $n = 1, 2, \dots$ , be a sequence in  $C$  converging to  $x_0$ . For each  $n \geq 1$  choose  $A_n \in A(x)$  such that  $x_n \in A_n \subset B$ . Since  $A(x)$  is compact in  $C(X)$ , we may assume that the sequence  $A_n$ ,  $n = 1, 2, \dots$ , also converges to an element  $A_0 \in A(x)$ . Obviously,  $x_0 \in A_0 \subset B$ . Hence  $x_0 \in A_0 \subset C$ . We conclude that  $C$  is closed in  $X$ .

Now suppose  $\varepsilon > 0$ . Since  $C$  is compact in  $X$ , there are points  $c_1, c_2, \dots, c_n$  in  $C$  such that  $C$  is contained in the  $\varepsilon$ -neighborhood of the finite set  $\{c_1, c_2, \dots, c_n\}$ . For each  $i$ , let  $A_i \in \mathcal{A}(x)$  such that  $c_i \in A_i \subset B$  and let  $B_0 = \bigcup_{i=1}^n A_i$ . Since  $C \supset B_0 \supset \{c_1, c_2, \dots, c_n\}$ , we have  $H(C, B_0) < \varepsilon$ . By Proposition 1.5,  $B_0 \in \mathcal{A}(x)$ . Since  $\mathcal{A}(x)$  is compact in  $C(X)$  we have  $C \in \mathcal{A}(x)$ .

*Proposition 1.8 [7]. If  $h: X \times I \rightarrow C(X)$  is a continuous increasing map such that  $x \in h(x, 0)$  for  $x \in X$  then  $h(x, t) \in \mathcal{A}(x)$  for  $(x, t) \in X \times I$ .*

*Theorem 1.9 [7]. If  $X$  is a nondegenerate metric continuum and  $C(X)$  is contractible, then  $X$  is an admissible space.*

## 2. Fiber Maps

In [5] it was shown that the contractibility of  $C(X)$  is equivalent to an existence of a set-valued map  $\alpha: X \rightarrow C(X)$  possessing a certain property. In this section we prove that this property is preserved by the homotopy equivalence relation. Hence, we obtain generalizations of many of the results in [4] and [8].

*Definition 2.1 [5]. A set-valued map  $\alpha: X \rightarrow C(X)$  is said to be a c-map if, for each  $x \in X$ ,  $\alpha(x)$  is a closed subset of the admissible fiber  $\mathcal{A}(x)$  such that*

(1)  $\{x\}, x \in \alpha(x)$ .

(2) For each pair  $A_0, A_1$  in  $\alpha(x)$  with  $A_0 \subset A_1$ , there is an ordered segment  $[2, p. 57]$  in  $\alpha(x)$  from  $A_0$  to  $A_1$ .

(3) For each  $A \in \alpha(x)$ , and  $\varepsilon > 0$ , there is a neighborhood  $W$  of  $x$  such that each point  $y$  of  $W$  has an element  $B \in \alpha(y)$  such that  $H(A, B) < \varepsilon$ .

We say that the space  $X$  is a *c-space* if there is a set-valued *c-map*  $\alpha: X \rightarrow C(X)$ . Clearly every *c-space* is an admissible space.

*Proposition 2.2* [5]. Every set-valued *c-map*  $\alpha: X \rightarrow C(X)$  is lower semicontinuous. Furthermore, if  $\hat{\alpha}(x, t) = \alpha(x) \cap \mu^{-1}(t)$ , then  $\hat{\alpha}$  is lower semicontinuous on  $X \times I$ .

*Theorem 2.3* [5]. Let  $X$  be a metric continuum. Then  $C(X)$  is contractible if and only if there is continuous set-valued *c-map* on  $X$  into  $C(X)$ .

In [1] Kelley defined a property (subsequently named property  $K$  in Nadler [2]) and proved that the hyperspaces of a space having property  $K$  are always contractible. The class of metric continua having property  $K$  includes locally connected continua and the hereditarily indecomposable continua. We now restate the result of Kelley.

*Proposition 2.4* [1]. If  $X$  has property  $K$ , then there is a continuous *c-map*  $\alpha: X \rightarrow C(X)$ .

*Proof.* Since  $X$  has property  $K$ ,  $F(x) = \mathcal{A}(x)$  by [5, Proposition 2.4] and  $F: X \rightarrow C(X)$  is continuous by [9, Theorem 2.2]. The existence of ordered segments in  $F(x)$  for every pair  $A_0 \subset A_1$  is given in [1, p. 24]. Hence the admissible fiber map  $\mathcal{A}$  is a continuous set-valued *c-map*.

Let  $X$  be a metric continuum. A function  $\alpha: X \rightarrow C^2(X)$  is called *admissible* if, for each  $x \in X$ ,

(1)'  $\{x\} \in \alpha(x)$ ,

(2)'  $\alpha(x) \subset A(x)$  and  $\alpha(x)$  is closed in  $A(x)$ ,

(3)'  $\alpha(x)$  contains a maximal element  $A_x$ , i.e.,  $A \subset A_x$  for all  $A \in \alpha(x)$ ,

(4)'  $\alpha(x)$  is segmentwise connected, i.e., for each pair  $A_0, A_1$  in  $\alpha(x)$  with  $A_0 \subset A_1$ , there is an ordered segment [2, p. 57] in  $\alpha(x)$  from  $A_0$  to  $A_1$ .

Let  $N_\alpha = \{A_x | A_x \text{ is a maximal element in } \alpha(x), x \in X\}$  and  $N_\alpha^2 = \{\{A_x\} | A_x \in N_\alpha\} \subset C^2(X)$ .

*Proposition 2.5. The following statements are equivalent.*

(1)  $C(X)$  is contractible.

(2) There is a continuous admissible function  $\alpha: X \rightarrow C^2(X)$  such that  $\cap N_\alpha \neq \emptyset$ .

(3) There is a continuous admissible function  $\alpha: X \rightarrow C^2(X)$  such that the set  $N_\alpha^2$  is contractible in  $C^2(X)$ .

*Proof.* (1)  $\Rightarrow$  (2). Suppose  $h: X \times I \rightarrow C(X)$  is an increasing contraction. Let  $\alpha(x) = \{h(x, t) | t \in I\}$ . Then the continuity of  $h$  provides the continuity of  $\alpha$  and it is obvious that  $\alpha$  satisfies the admissible conditions (1')-(4') with  $A_x = h(x, 1) = \cap N_\alpha$ .

(2)  $\Rightarrow$  (3). Suppose  $\alpha: X \rightarrow C^2(X)$  is a continuous admissible function such that  $\cap N_\alpha \neq \emptyset$ . Let  $x_0 \in \cap N_\alpha$  and let  $\gamma: I \rightarrow C(X)$  be an ordered segment from  $\{x_0\}$  to  $X$ . Define  $\beta: N_\alpha^2 \times I \rightarrow C^2(X)$  by

$$\beta(\{A_x\}, t) = \{A_x \cup \gamma(t)\}.$$



Then  $\beta$  is continuous and  $\beta(\{A_x\}, 1) = \{X\}$  for each  $\{A_x\} \in \mathcal{N}_\alpha^2$ .

(3)  $\Rightarrow$  (1). Suppose  $\alpha: X \rightarrow C^2(X)$  is a continuous admissible function and  $\beta: \mathcal{N}_\alpha^2 \times I \rightarrow C^2(X)$  is a contraction. We may assume  $\beta$  is increasing. Let  $\sigma: C^2(X) \rightarrow C(X)$  be the function defined by  $\sigma(T) = UT$ . Then  $\sigma$  is continuous.

We now observe that for each  $x \in X$  the maximal element  $A_x$  of  $\alpha(x)$  is unique and  $A_x = \sigma \circ \alpha(x)$ . Therefore the function  $x \rightarrow \{A_x\}$  is continuous from  $X$  to  $C^2(X)$ . Hence we define a function  $\tau: X \rightarrow C^2(X)$  by  $\tau(x) = \alpha(x) \cup \{\sigma\beta(\{A_x\}, t) \mid t \in I\}$ . Then  $\tau$  is continuous. Since  $\sigma\beta(\{A_x\}, 1) = A$ , for some  $A$ , and for all  $x \in X$ , we may define an ordered segment  $\gamma: I \rightarrow C(X)$  from  $A$  to  $X$  and join it to  $\tau$ , that is  $\phi(x) = \tau(x) \cup \{\gamma(t) \mid t \in I\}$ . Then it is not difficult to check that  $\phi$  satisfies the definition of a set-valued c-map and  $\phi$  is continuous. Hence by Theorem 2.3,  $C(X)$  is contractible.

Suppose  $X$  and  $Y$  are metric continua.

*Theorem 2.6.* Suppose  $f: X \rightarrow Y$  and  $g: Y \rightarrow X$  are continuous functions such that  $f \circ g: Y \rightarrow Y$  is homotopic to the identity  $\text{id}_Y$ . If  $X$  is a c-space then  $Y$  is a c-space.

*Proof.* Let  $h: Y \times I \rightarrow Y$  be a homotopy such that  $h(y, 0) = y$  and  $h(y, 1) = f \circ g(y)$  for each  $y \in Y$ . Let  $\bar{h}(y, t) = U\{h(y, s) \mid 0 \leq s \leq t\}$ . Then  $\bar{h}: Y \times I \rightarrow C(Y)$  is a continuous homotopy such that  $\bar{h}(y, t) \subset \bar{h}(y, t')$  whenever  $t \leq t'$ . Let  $\beta_1(y) = \{\bar{h}(y, t) \mid t \in I\}$ . Then the continuity of  $\bar{h}$  implies the continuity of  $\beta_1: Y \rightarrow C^2(Y)$  and each element  $\bar{h}(y, t)$  of the set  $\beta_1(y)$  is an admissible element at  $y$  such

that for each pair  $B_0, B_1$  in  $\beta_1(y)$  with  $B_0 \subset B_1$ , there is an ordered segment in  $\beta_1(y)$  from  $B_0$  to  $B_1$ .

Let  $\alpha: X \rightarrow C(X)$  be a set-valued c-map. For  $y \in Y$ , let  $x = g(y)$  and  $\beta_2(y) = \{\bar{h}(y, 1) \cup f(A) \mid A \in \alpha(x)\}$ . Since  $f(x) = f \circ g(y) \in \bar{h}(y, 1) \cap f(A)$ , we have  $\bar{h}(y, 1) \cup f(A) \in C(Y)$ . Now we will show that  $\beta_2: Y \rightarrow C^2(Y)$  is lower semicontinuous.

Let  $\varepsilon > 0$ . Since  $f$  is continuous, there is  $\varepsilon' > 0$  such that if  $A, A' \in C(X)$  such that  $A$  and  $A'$  are less than  $\varepsilon'$  apart, then  $H(f(A), f(A')) < \varepsilon$ . Since  $\alpha$  is lower semicontinuous and  $\alpha(x)$  is compact, there exists  $\delta > 0$  such that if  $d(x, x') < \delta$  and  $A \in \alpha(x)$ , there is an element  $A' \in \alpha(x')$  such that  $A$  and  $A'$  are less than  $\varepsilon'$  apart. Now the continuity of  $g$  implies that there is  $\delta_0 > 0$  such that if  $d(y, y') < \delta_0$  then  $d(g(y), g(y')) < \delta$ . Also, by the continuity of  $\bar{h}$ , we choose  $\delta_1 > 0$  such that if  $d(y, y') < \delta_1$ , then  $H(\bar{h}(y, 1), \bar{h}(y', 1)) < \varepsilon'$ . Let  $\bar{\delta} = \min\{\delta_0, \delta_1\}$ ,  $x = g(y)$ ,  $x' = g(y')$ . Then if  $d(y, y') < \bar{\delta}$  and  $\bar{h}(y, 1) \cup f(A) \in \beta_2(y)$  then there is  $\bar{h}(y', 1) \cup f(A') \in \beta_2(y')$  such that  $H(\bar{h}(y, 1) \cup f(A), \bar{h}(y', 1) \cup f(A')) \leq \max\{H(\bar{h}(y, 1), \bar{h}(y', 1)), H(f(A), f(A'))\} < \varepsilon$  by Lemma 1.4. Hence the elements of  $\beta_2(y)$  are admissible at  $y$  and  $\beta_2$  is lower semicontinuous. Since  $f$  preserves ordered segments, we see that  $\beta_2$  satisfies the condition of ordered segment. Now let  $\gamma: I \rightarrow C(Y)$  be an ordered segment from  $f(X)$  to  $Y$  and let  $\beta_3(y) = \{\gamma(t) \cup \bar{h}(y, 1) \mid t \in I\}$  and  $\beta(y) = \beta_1(y) \cup \beta_2(y) \cup \beta_3(y)$ . The maximal element of  $\beta_1(y)$  is  $\bar{h}_1(y, 1)$  which is also the minimal element of  $\beta_2(y)$ , and the maximal element of  $\beta_2(y)$  is  $\bar{h}(y, 1) \cup f(X)$ , and the minimal element of  $\beta_3(y)$  is  $f(X) \cup \bar{h}(y, 1)$ . So the continuity of  $\beta_1$

and  $\beta_3$  together with the lower semicontinuity of  $\beta_2$  provide the lower semicontinuity of  $\beta$ , and hence (3) is verified for  $\beta$ . It is easy to verify that  $\beta$  also satisfies the condition (1) and (2) of Definition 2.1.

*Corollary 2.7.* Suppose  $X$  and  $Y$  are homotopically equivalent metric continua. Then  $X$  is a  $c$ -space if and only if  $Y$  is.

Let  $\text{id}_Y$  denote the identity map of  $Y$  onto itself and  $[f]$  the homotopy class of continuous maps of  $Y$  into itself which contains  $f$ .

*Theorem 2.8.* A metric continuum  $Y$  is a  $c$ -space if and only if for some  $g$  in  $[\text{id}_Y]$  it is true that  $g(Y)$  is a  $c$ -space.

*Proof.* If  $Y$  is a  $c$ -space then let  $g = \text{id}_Y$ . Conversely, suppose for some  $g \in [\text{id}_Y]$ ,  $g(Y)$  is a  $c$ -space. Let  $X = g(Y)$  and  $f: X \rightarrow Y$  be the inclusion map. Then  $f \circ g = g \in [\text{id}_Y]$ . So Theorem 2.6 gives the conclusion.

*Corollary 2.9.* Suppose  $X$  is a deformation retract of  $Y$ . If  $X$  is a  $c$ -space, so is  $Y$ .

*Proof.* Let  $\gamma: Y \rightarrow X$  be a retraction which is homotopic to the identity map  $\text{id}_Y$ . Then Theorem 2.8 provides the conclusion.

*Corollary 2.10.* If  $Y$  is a retract of  $X$  and  $X$  is a  $c$ -space, then so is  $Y$ .

*Proof.* Let  $f: X \rightarrow Y$  be a retraction and  $g: Y \rightarrow X$  the inclusion map. Then  $f \circ g = \text{id}_Y$ . So by Theorem 2.6,  $Y$  is a  $c$ -space.

*Theorem 2.11.* Let  $X = X_1 \cup X_2$  where  $X$  and  $X_1$  are subcontinua and  $X_2$  is a closed subset such that  $X_1 \cap X_2$  is a strong deformation retract of  $X_2$ . If  $X_1$  is a  $c$ -space so is  $X$ .

*Proof.*  $X_1$  is a deformation retract of  $X$ .

*Theorem 2.12.* The product space  $X \times Y$  is a  $c$ -space if and only if both  $X$  and  $Y$  are  $c$ -spaces.

*Proof.* Each factor space is a retract of  $X \times Y$ . Therefore by Corollary 2.10,  $X \times Y$  is a  $c$ -space.

Conversely, if  $\alpha_X: X \rightarrow C(X)$  and  $\alpha_Y: Y \rightarrow C(Y)$  are set-valued  $c$ -maps, then  $\alpha_X \times \alpha_Y: X \times Y \rightarrow C(X \times Y)$  defined by  $\alpha_X \times \alpha_Y(x, y) = \alpha_X(x) \times \alpha_Y(y) = \{A \times B \mid A \in \alpha_X(x), B \in \alpha_Y(y)\}$  is a set-valued  $c$ -map.

## References

- [1] J. L. Kelley, *Hyperspaces of a metric continuum*, Trans. Amer. Math. Soc. 52 (1942), 22-36.
- [2] S. B. Nadler, Jr., *Hyperspaces of sets*, Marcel Dekker, Inc., 1978.
- [3] T. Nishiura and C. J. Rhee, *Cut points of  $X$  and hyperspace of subcontinua*, Proc. Amer. Math. Soc. 82 (1982), 149-154.
- [4] \_\_\_\_\_, *Contractibility of hyperspace of subcontinua*, Houston J. Math. 8 (1982), 119-127.
- [5] C. J. Rhee, *On a contractible hyperspace condition*, Topology Proc. 7 (1982), 147-155.
- [6] \_\_\_\_\_, *On dimension and hyperspace of a metric continuum*, Bull. de la Soc. Royale des Sciences de Liege 38 (1969), 602-604.

- [7] C. J. Rhee and T. Nishiura, *Contractible hyperspaces of subcontinua* (submitted).
- [8] \_\_\_\_\_, *On contraction of hyperspaces of metric continua* (preprint).
- [9] R. Wardle, *On a property of J. L. Kelley*, Houston J. Math. 3 (1977), 291-299.
- [10] H. Whitney, *Regular families of curves*, Annals of Math. 34 (1933), 244-270.
- [11] M. Wojdyslawski, *Sur la Contractibilité des hyperspaces des continus localement connexes*, Fund. Math. 30 (1938), 247-252.

Wayne State University  
Detroit, Michigan 48202