

---

# TOPOLOGY PROCEEDINGS



Volume 9, 1984

Pages 149–163

---

<http://topology.auburn.edu/tp/>

## SPACES DOMINATED BY METRIC SUBSETS

by

YOSHIO TANAKA AND ZHOU HAO-XUAN

---

### Topology Proceedings

**Web:** <http://topology.auburn.edu/tp/>

**Mail:** Topology Proceedings  
Department of Mathematics & Statistics  
Auburn University, Alabama 36849, USA

**E-mail:** [topolog@auburn.edu](mailto:topolog@auburn.edu)

**ISSN:** 0146-4124

COPYRIGHT © by Topology Proceedings. All rights reserved.

## SPACES DOMINATED BY METRIC SUBSETS

Yoshio Tanaka and Zhou Hao-xuan

### Introduction

Let  $X$  be a space and  $\mathcal{J}$  be a closed cover of  $X$ . Then  $X$  is determined by  $\mathcal{J}^1$ , if  $A \subset X$  is closed in  $X$  whenever  $A \cap F$  is relatively closed in  $F$  for each  $F \in \mathcal{J}$ . A space  $X$  is a  $k$ -space (resp. sequential space) if it is determined by the cover of all compact subsets (resp. compact metric subsets). Recall that  $X$  is dominated by  $\mathcal{J}^2$ , if the union of any subcollection  $\mathcal{J}'$  of  $\mathcal{J}$  is closed in  $X$  and the union is determined by  $\mathcal{J}'$ . We remark that if  $X$  is dominated by  $\mathcal{J}$ , then it is determined by  $\mathcal{J}$ , but the converse does not hold. In case that the closed cover  $\mathcal{J}$  is increasing and countable, the converse holds. Every CW-complex, more generally every chunk-complex [2] is dominated by a cover of compact metric subsets.

Now, let  $X$  be a regular space determined by a point-countable closed cover of separable metric subsets. In [14] and [15], the first author showed respectively that  $X$  is Fréchet if and only if it is Lašnev, and that  $X$  is metric if and only if it contains no closed copy of a sequential fan  $S_\omega$  and no Arens' space  $S_2$ . Recall that a space  $X$  is Fréchet, if for every  $A \subset X$  and every  $x \in \bar{A}$  there

---

<sup>1</sup>Following [7], we use " $X$  is determined by  $\mathcal{J}$ " instead of the usual " $X$  has the weak topology with respect to (or determined by)  $\mathcal{J}$ ".

<sup>2</sup>In some literature, " $X$  has the weak (hereditarily weak; or Whitehead weak) topology with respect to  $\mathcal{J}$ " is used instead of " $X$  is dominated by  $\mathcal{J}$ ".

exists a sequence in  $A$  converging to the point  $x$ . Also, a space  $X$  is *Lašnev* if  $X$  is the closed image of a metric space. The space  $S_\omega$  is the quotient space obtained from the topological sum of countably many convergent sequences by identifying all the limit points. As for the space  $S_2$ , e.g., see [4; Example 1.6.19].

In this paper, we shall give some analogous results  
 1 spaces are dominated by metric subsets, and investigate  
 chunk-complexes and CW-complexes as spaces dominated by  
 compact metric subsets. In Section 1, we show that every  
 Fréchet space dominated by metric subsets is *Lašnev*, and  
 that every space dominated by metric subsets is metric if  
 and only if it contains no closed copy of  $S_\omega$  and no  $S_2$ .  
 The former is an affirmative answer to a question in [14;  
 Problem 2.4], and the latter was proved in [5] for every  
 space dominated by countably many compact metric subsets.  
 In Section 2, we show that every Fréchet space dominated by  
 compact metric subsets is equivalent to a Fréchet chunk-  
 complex, and such a complex is characterized as the closed  
 image of a locally compact metric space<sup>3</sup>. In Section 3, we  
 show that every Fréchet CW-complex is especially the closed  
 image of the topological sum of Euclidean simplexes (hence,  
 the closed image of a locally compact metric CW-complex).

We assume that all spaces are Hausdorff and all maps  
 are continuous and onto.

---

<sup>3</sup>After preparing this paper, M. Ito independently  
 proved that every Fréchet chunk-complex is *Lašnev*.

# 1. Spaces Dominated by Metric Subsets and Lasnev Spaces

*Lemma 1.1.* Suppose that  $X$  is a Fréchet space dominated by a closed cover  $\{X_\alpha; \alpha < \beta\}$ . Let  $Y_\alpha = \overline{X_\alpha - \bigcup_{\delta < \alpha} X_\delta}$ , and  $Y_\alpha^* = Y_\alpha \times \{\alpha\}$ . Let  $X^*$  be the topological sum  $\sum_\alpha Y_\alpha^*$ . Then the obvious map  $f: X^* \rightarrow X$  is closed.

*Proof.* Since  $f$  is clearly continuous, to prove that  $f$  is closed, it is sufficient to show that  $\overline{f(F)} \subset f(\overline{F})$  for any  $F \subset X^*$ . Let  $x \in \overline{f(F)} - f(F)$ . Since  $X$  is Fréchet, there exist distinct points  $x_i \in f(F)$  with  $x_i \rightarrow x$ . Choose a point  $p_i \in f^{-1}(x_i) \cap F$  for each  $i \in \mathbb{N}$ , and let  $P = \{p_i; i \in \mathbb{N}\}$ . Suppose that  $P$  is not contained in any finite union of  $Y_\alpha^*$ 's. Let  $\alpha_1 = \min\{\alpha; Y_\alpha^* \cap P \neq \emptyset\}$  and choose a point  $p_{i_1} \in Y_{\alpha_1}^*$ . By induction, we can choose countably many ordinals  $\alpha_n = \min\{\alpha; Y_\alpha^* \cap \{p_i; i > i_{n-1}\} \neq \emptyset, \alpha \neq \alpha_1, \dots, \alpha_{n-1}\}$  and countably many points  $p_{i_n} \in Y_{\alpha_n}^*$  with  $i_n < i_{n+1}$ . Then for each  $n \in \mathbb{N}$ ,  $\alpha_n < \alpha_{n+1}$  and the subset  $Y_{\alpha_n}$  of  $X$  contains  $x_{i_n} = f(p_{i_n})$ . Since  $X$  is Fréchet, for each  $n \in \mathbb{N}$  there exist  $y_{nk} \in X_{\alpha_n} - \bigcup_{\delta < \alpha_n} X_\delta$  with  $y_{nk} \rightarrow x_{i_n}$ . Let  $A = \{y_{nk}; n, k \in \mathbb{N}\} - \{x\}$ . Since  $x \in \overline{A}$ , there exists a sequence  $B$  in  $A$  converging to the point  $x$ . Let  $C = \bigcup\{X_{\alpha_n}; n \in \mathbb{N}\}$ . Then  $B \subset C$  and  $B \cap X_{\alpha_n}$  is finite for each  $n \in \mathbb{N}$ . But the closed subset  $C$  of  $X$  is determined by  $\{X_{\alpha_n}; n \in \mathbb{N}\}$ . Then  $B$  is closed in  $C$ , hence in  $X$ . Thus,  $x \in B$ , a contradiction. Hence the subset  $P$  of  $X^*$  is contained in a finite union of  $Y_\alpha^*$ 's. So, infinitely many points  $p_{n_j}$  are in some  $Y_{\alpha_0}^*$ . But  $f|_{Y_{\alpha_0}^*}$  is a homeomorphism and  $x_{n_j} = f(p_{n_j}) \in Y_{\alpha_0}$ .

Thus, since  $x_{n_j} \rightarrow x$ ,  $p_{n_j} \rightarrow y$  for some  $y \in Y_{\alpha_0}^*$  with  $f(y) = x$ .

Then, since  $p_{n_j} \in f^{-1}(x_{n_j}) \cap F$ ,  $y \in \bar{F}$ , so that  $x \in f(\bar{F})$ .

Thus,  $\overline{f(F)} \subset f(\bar{F})$ . That completes the proof.

By Lemma 1.1, we have the following affirmative answer to the question whether every Fréchet space dominated by compact metric subsets (e.g., Fréchet CW-complex) is Lašnev; see [14; Problem 2.4].

*Theorem 1.2. Every Fréchet space dominated by a closed cover of metric (resp. locally compact metric) subsets is a Lašnev space (resp. the closed image of a locally compact metric space).*

*Theorem 1.3. Let  $X$  be dominated by a closed cover of Lašnev subsets. Then  $X$  is Lašnev if and only if it is Fréchet.*

*Proof.* The "only if" part follows from the easy fact that every Lašnev space is Fréchet. The subsets  $X_\alpha$  of  $X$  in Lemma 1.1 are Lašnev, so are the subsets  $Y_\alpha^*$  of  $X^*$ . Thus the "if" part is routinely verified by Lemma 1.1.

Not every countable CW-complex is decomposed into a metric subset and a  $\sigma$ -discrete subset; see [14; p. 284]. But, among Fréchet spaces, we have the following decomposition theorem by Theorem 1.2 and [8; Theorem 2] (resp. [11; Theorem 4]).

*Theorem 1.4.\* Every Fréchet space dominated by a closed cover of metric (resp. locally compact metric) subsets is*

---

\* See note added in proof.

*decomposed into a metric (resp. locally compact metric) subset and a  $\sigma$ -discrete (resp. closed discrete) subset.*

In [5], the metrizability of a space dominated by countably many compact metric subsets is characterized by whether or not it contains two copies of the spaces  $S_\omega$  and  $S_2$ . As for the metrizability of a space dominated by metric subsets, analogously we have

*Theorem 1.5. Let  $X$  be dominated by a closed cover of metric (resp. locally compact metric) subsets. Then  $X$  is metric (resp. locally compact metric) if and only if it contains no closed copy of  $S_\omega$  and no  $S_2$ .*

*Proof.* The "only if" part is obvious, so we will prove the "if" part. Since  $X$  is dominated by metric subsets, by [10; Theorem 3]  $X$  is a regular space in which every point is  $G_\delta$ . Now,  $X$  is a sequential space which contains no closed copy of  $S_\omega$  and no  $S_2$ . Thus, by [15; Theorem 3.1],  $X$  is strongly Fréchet; that is, if whenever  $x \in \overline{A_n}$  with  $A_{n+1} \subset A_n$ , then there exists  $x_n \in A_n$  with  $x_n \rightarrow x$ . Since  $X$  is Fréchet, by Theorem 1.2,  $X$  is the closed image of a metric (resp. locally compact metric) space. Then, since  $X$  is strongly Fréchet,  $X$  is metric (resp. locally compact metric) by [9; Corollary 9.10].

In concluding this section, let us consider the products of spaces dominated by metric subsets. For  $c = 2^\omega$ , the quotient space  $S_c$  is similarly defined as the space  $S_\omega$ .

*Theorem 1.6. Let  $X$  be dominated by a closed cover of metric subsets. Then the following are equivalent.*

- (1)  $X$  is a locally compact metric space.
- (2)  $X \times S_C$  is a  $k$ -space.
- (3)  $X \times S_C$  is dominated by a closed cover of metric subsets.

*Proof.* (3)  $\rightarrow$  (2) is easily proved.

(1)  $\rightarrow$  (3). Let  $\mathcal{J}$  be a locally finite cover of compact metric subsets of  $X$ , and  $\mathcal{C}$  be the obvious cover of the infinite convergent sequences in  $S_C$ . Since  $S_C$  is dominated by  $\mathcal{C}$ , from [10; Theorem 1],  $X \times S_C$  is dominated by a cover  $\mathcal{J} \times \mathcal{C}$  of compact metric subsets.

(2)  $\rightarrow$  (1). Suppose that  $X$  contains a closed copy  $P$  of  $S_\omega$  or  $S_2$ . Note that the space  $S_2$  is the perfect pre-image of  $S_\omega$ . Thus,  $P \times S_C$  is the perfect pre-image of  $S_\omega \times S_C$ . Since  $P \times S_C$  is a  $k$ -space, so is  $S_\omega \times S_C$ . But, the example [3; p. 563] implicates that  $S_\omega \times S_C$  is not a  $k$ -space. This contradiction implies that  $X$  contains no closed copy of  $S_\omega$  and no  $S_2$ . Then  $X$  is metric by Theorem 1.5. Thus each point of  $X \times S_C$  is  $G_\delta$ . Then a  $k$ -space  $X \times S_C$  is sequential by [9; Theorem 7.3]. Thus a metric space  $X$  is locally compact by [12; Theorem 1.1].

*Theorem 1.7. Let  $X$  be dominated by a closed cover of metric subsets. Then (1) and (2) below hold.*

(1) *Let  $X$  be a Fréchet space. Then  $X^2$  is a  $k$ -space if and only if  $X$  is metric, or  $X$  is dominated by a countable closed cover of locally compact metric subsets.*

(2)  $X^\omega$  is a  $k$ -space if and only if  $X$  is metric.

*Proof.* (1) Since  $X$  is Lašnev by Theorem 1.2, (1) follows from [6; Theorem 2.15].

(2) Let  $X^\omega$  be a  $k$ -space.  $(S_\omega)^\omega$  is not a  $k$ -space by [12; Proposition 4.2]. While,  $(S_2)^\omega$  is the perfect pre-image of  $(S_\omega)^\omega$ . Then  $X$  contains no closed copy of  $S_\omega$  and no  $S_2$ . Hence  $X$  is metric by Theorem 1.5.

## 2. Chunk-Complexes and Closed Images of Locally Compact Metric Spaces

Recall that a *chunk-complex* [2] is a space determined by a cover  $\mathcal{H}$  of compact metric subsets, called *chunks*, such that for  $S, T \in \mathcal{H}$ , either  $S \cap T = \emptyset$  or  $S \cap T \in \mathcal{H}$ , and for  $S \in \mathcal{H}$ ,  $\{T \in \mathcal{H}; T \subset S\}$  is finite. A chunk-complex is an  $M_1$ -space [2] dominated by its chunks. Every CW-complex and every locally compact metric space is a chunk-complex. Let us call a chunk-complex *locally countable*, if it has a locally countable cover of chunks.

*Lemma 2.1.* *Every space determined by a locally countable cover  $\mathcal{J}$  of compact metric subsets is a locally countable chunk-complex.*

*Proof.* For any  $A \in \mathcal{J}$ ,  $\{F \in \mathcal{J}; A \cap F \neq \emptyset\}$  is countable. Then, since  $X$  is determined by  $\mathcal{J}$ , by the proof of (a)  $\rightarrow$  (c) of [13; Theorem 1],  $X$  is the topological sum of spaces  $X_\beta$  ( $\beta \in B$ ) determined by countably many compact metric subsets  $X_{\beta n}$  ( $n \in \mathbb{N}$ ). We can assume that  $X_{\beta n} \subset X_{\beta n+1}$  for  $\beta \in B$  and  $n \in \mathbb{N}$ . Then  $X$  is a locally countable chunk-complex with chunks  $X_{\beta n}$ .



The regular, separable, non-Lindelöf space  $Y$  of [7; Example 9.3] shows that every space determined by a point-finite cover of compact metric subsets is not Lašnev, nor a chunk-complex. But, among Fréchet spaces, we have

*Theorem 2.2. Let  $X$  be a regular Fréchet space. Then the following are equivalent.*

(1)  $X$  is determined by a point-countable cover of compact metric subsets.

(2)  $X$  is a locally countable chunk-complex.

(3)  $X$  is the quotient  $s$ -image of a locally compact metric space.

(4)  $X$  is the closed  $s$ -image of a locally compact metric space.

*Proof.* The equivalence  $(1) \leftrightarrow (3) \leftrightarrow (4)$  follows from [14; Theorem 2.2] and [1; Theorem 4]. By Lemma 1.1, we have  $(2) \rightarrow (4)$ .

$(4) \rightarrow (2)$ . Let  $f: L \rightarrow X$  be a closed  $s$ -map with  $L$  locally compact metric. Since  $L$  is determined by a locally finite cover  $\mathcal{J}$  of compact metric subsets,  $X$  is determined by a locally countable cover  $f(\mathcal{J})$  of compact metric subsets. Thus by Lemma 2.1, we have  $(2)$ .

Every chunk-complex is a space dominated by a cover of compact metric subsets. However, the authors do not know whether the converse holds. So, we shall pose the following question.

*Question 2.3.* Is every space dominated by a cover of compact metric subsets a chunk-complex?

Concerning the above question, if the cover is point-countable (equivalently, locally countable), then the answer is affirmative by Lemma 2.1. Among Fréchet spaces, the answer is also affirmative without the point-countability, and in addition, every Fréchet chunk-complex is characterized as the closed image of a locally compact metric space. To prove this, we need the following lemma.

*Lemma 2.4.* Let  $X$  be a Fréchet space dominated by a cover  $\{X_\alpha; \alpha < \beta\}$  of compact subsets, and let  $Y_\alpha = \overline{X_\alpha - \bigcup_{\delta < \alpha} X_\delta}$ . Then for each  $\alpha$ , there exists a finite subset  $I$  such that  $\bigcup \{Y_\delta \cap Y_\alpha; \delta \notin I\}$  is finite.

*Proof.* For some  $\alpha_0$ , suppose that  $\bigcup \{Y_\alpha \cap Y_{\alpha_0}; \alpha \notin I\}$  is infinite for any finite subset  $I$ . Then, by induction we can choose countable distinct points  $y_n \in X$  and countably many ordinals  $\alpha_n$  with  $\alpha_n < \alpha_{n+1}$  such that  $y_n \in Y_{\alpha_n} \cap Y_{\alpha_0}$  for each  $n \in \mathbb{N}$ . Since  $y_n \in Y_{\alpha_n}$ , there exists a sequence  $L_n$  in  $X_{\alpha_n} - \bigcup_{\delta < \alpha_n} X_\delta$  converging to the point  $y_n$ . Since the points  $y_n$  are contained in a compact subset  $X_{\alpha_0}$ , there exists a sequence  $\{y_{n_i}; i \in \mathbb{N}\}$  accumulating to a point  $y_0 \in X_{\alpha_0}$  with  $y_{n_i} \neq y_0$ . Then,  $y_0 \in \overline{\bigcup \{L_{n_i}; i \in \mathbb{N}\}}$ . Thus there exists a sequence  $P = \{p_j; j \in \mathbb{N}\}$  converging to the point  $y_0$  such that  $p_j \in L_{n_i(j)}$  and  $p_j \neq y_0$ . But,  $P \cap X_{\alpha_{n_i(j)}}$  is finite for each  $j \in \mathbb{N}$ , so that  $P$  is closed in  $X$ . Hence,  $y_0 \in P$ , a contradiction.

*Theorem 2.5. The following are equivalent.*

- (1) *X is a Fréchet space dominated by a cover of compact metric subsets.*
- (2) *X is a Fréchet chunk-complex.*
- (3) *X is the closed image of a locally compact metric space.*

*Proof.* (2)  $\rightarrow$  (3) follows from Theorem 1.2.

(3)  $\rightarrow$  (1). The closed image  $X$  of a locally compact metric space has a hereditarily closure-preserving cover<sup>4</sup> of compact metric subsets. Then  $X$  is dominated by this cover.

(1)  $\rightarrow$  (2). To show that  $X$  is a chunk-complex, let  $\mathcal{J} = \{Y_\alpha; \alpha < \beta\}$  be a collection of the subsets  $Y_\alpha$  in Lemma 2.4; here the  $Y_\alpha$  are compact metric. Let  $\mathcal{H}$  be the cover consisting of all finite intersections of members of  $\mathcal{J}$ . We will prove that  $\mathcal{H}$  is a cover of chunks for  $X$ . Since  $X$  is Fréchet, in view of Lemma 1.1,  $\mathcal{J}$  is a hereditarily closure-preserving cover of  $X$ . Thus  $X$  is determined by  $\mathcal{J}$ . Since  $\mathcal{J} \subset \mathcal{H}$ ,  $X$  is determined by  $\mathcal{H}$ . To show that for each  $S \in \mathcal{H}$ ,  $\{T \in \mathcal{H}; T \subset S\}$  is finite, take any  $Y_{\alpha_0} \in \mathcal{J}$  with  $S \subset Y_{\alpha_0}$ . By Lemma 2.4, there exists a finite subset  $I_0$  such that  $F_0 = \cup\{Y_\alpha \cap Y_{\alpha_0}; \alpha \notin I_0\}$  is finite. For any  $T \subset S$ , let  $T = \cap\{T_{\alpha_i}; i = 1, 2, \dots, n\}$ . Then all  $\alpha_i \in I_0$ , otherwise some  $\alpha_{i_0} \notin I_0$ . The latter case implies that  $T \subset Y_{\alpha_{i_0}} \cap Y_{\alpha_0} \subset F_0$ , so that  $T$  is a subset of the finite

---

<sup>4</sup>A cover  $\mathcal{J} = \{F_\alpha; \alpha \in A\}$  of a space is hereditarily closure-preserving if  $\cup\{A_\alpha; \alpha \in A\} = \cup\{A_\alpha; \alpha \in A\}$  for any  $A_\alpha \subset F_\alpha$ .

set  $F_0$ . Hence,  $\{T \in \#; T \subset S\}$  is finite, because there exist only finitely many subsets of the finite set  $I_0$  and only finitely many subsets of the finite set  $F_0$ .

Let  $f: X \rightarrow Y$  be a closed map such that  $X$  is dominated by a closed cover  $\mathcal{J}$ . Then it is easy to show that  $Y$  is dominated by  $f(\mathcal{J})$ . Thus we have the following by Theorem 2.5.

*Corollary 2.6. Every Fréchet space which is the closed image of a chunk-complex is a chunk-complex.*

*Corollary 2.7. Every Fréchet space dominated by a closed cover of chunk-complexes is a chunk-complex.*

*Proof.* Let  $X$  be a Fréchet space dominated by a closed cover  $\{X_\alpha; \alpha < \beta\}$  of chunk-complexes, and let  $Y_\alpha = \overline{X_\alpha - \bigcup_{\delta < \alpha} X_\delta}$ . Then by Lemma 1.1,  $X$  is the closed image of the topological sum of  $Y_\alpha$ 's. But each closed subset  $Y_\alpha$  of a chunk-complex  $X_\alpha$  is dominated by compact metric subsets. Since  $Y_\alpha$  is Fréchet, by Theorem 1.2,  $Y_\alpha$  is the closed image of a locally compact metric space, hence so is  $X$ . Thus  $X$  is a chunk-complex by Theorem 2.5.

### 3. CW-Complexes

Let  $X$  be a complex. A subcomplex  $L$  of  $X$  is the union of a subset of the cells of  $X$ , which are the cells of  $L$ , such that, if  $e \in L$  then  $\bar{e} \subset L$ . Recall that  $X$  is a CW-complex if it is dominated by the cover of all finite subcomplexes, and each cell is contained in a finite subcomplex. Note that every CW-complex is determined by the

cover of all closed cells  $\bar{e}$ , but not always dominated by this cover.

Now, every CW-complex is a chunk-complex. Thus, by Theorem 2.5, every Fréchet CW-complex is the closed image of a locally compact metric space. But this conclusion can be refined by writing "CW-complex" instead of "space." To show this, we need the following lemma. The proof is essentially the same as in the proof of Lemma 2.4.

*Lemma 3.1. Let  $X$  be a CW-complex with cells  $\{e\}$ . If  $X$  is Fréchet, then for each  $e \in X$  there is a finite collection  $E$  of cells such that  $\cup\{\bar{\sigma} \cap e; \sigma \in X - E\}$  is finite.*

*Theorem 3.2. Let  $X$  be a CW-complex. Then the following are equivalent.*

(1)  $X$  is a Fréchet space (resp. Fréchet, locally countable CW-complex<sup>5</sup>).

(2)  $X$  is the closed image (resp. closed  $s$ -image) of a locally compact metric CW-complex. (The domain is actually the topological sum of Euclidean simplexes.)

*Proof.* Since every Lašnev space is Fréchet, (2)  $\rightarrow$  (1) is obvious. For the parenthetic part, since  $X$  is a locally separable space, the CW-complex  $X$  is locally countable.

To prove (1)  $\rightarrow$  (2), let  $X$  be a Fréchet CW-complex with cells  $\{e\}$ . Let  $X^*$  be the topological sum  $\sum \bar{e}^*$  of  $\bar{e}^*$ 's. Since each  $\bar{e}^*$  is the closed image of a Euclidean simplex  $\sigma$ ,

---

<sup>5</sup>A CW-complex  $X$  is locally countable, if for each  $x \in X$  there exists a countable subcomplex  $A$  of  $X$  such that  $x \in \text{int } A$ ; equivalently, the cover of all closed cells of  $X$  is locally countable.

$X^*$  is the closed image of  $\sum \sigma$ . Thus it suffices to prove that the obvious map  $f: X^* \rightarrow X$  is closed. Let  $A$  be a closed subset of  $X^*$ . Then we will prove that  $f(A) \cap \bar{d}$  is closed in  $X$  for each cell  $d \in X$ , so that  $f(A)$  is closed in  $X$ . For  $d \in X$ , let  $D$  be a finite subcomplex of  $X$  containing  $d$ . Let  $A_e = f(A \cap \bar{e}^*) \cap \bar{d}$  for each  $e \in X$ . Then,  

$$f(A) \cap \bar{d} = \bigcup \{A_e; e \in X\} = (\bigcup \{A_e; e \in D\}) \cup (\bigcup \{A_e; e \notin D\}).$$
Since each  $A_e$  is closed in  $X$ , so is  $\bigcup \{A_e; e \in D\}$ . Thus it is sufficient to prove that  $B = \bigcup \{A_e; e \notin D\}$  is closed in  $X$ . By Lemma 3.1 there is a finite collection  $F$  of cells such that  $C = \bigcup \{\bar{e} \cap d'; d' \in D, e \in X - F\}$  is finite. Since  $B = \bigcup \{A_e \cap d'; d' \in D, e \notin D\}$ ,  $B = (\bigcup \{A_e \cap d'; d' \in D, e \in F - D\}) \cup (\bigcup \{A_e \cap d'; d' \in D, e \in X - (F \cup D)\})$ . Since  $\bigcup \{A_e \cap d'; d' \in D\} = A_e$  is closed in  $X$ , so is the first union. The second union is contained in a finite subset  $C \cap \bar{d} = \bigcup \{(\bar{e} \cap d') \cap \bar{d}; d' \in D, e \in X - F\}$ . Thus the second union is finite, hence is closed in  $X$ . Thus  $B$  is closed in  $X$ . Hence  $f$  is a closed map. That completes the proof.

The proof of the previous theorem suggests the "only if" part of the following. The "if" part is obvious.

*Corollary 3.3. Let  $X$  be a CW-complex with cells  $\{e\}$ . Then  $X$  is Fréchet if and only if  $\{\bar{e}\}$  is a hereditarily closure-preserving cover of  $X$ .*

*Note Added in Proof.* The authors have recently shown that every Fréchet space dominated by a closed cover of metric subsets is decomposed into a metric subset and a closed discrete subset.

## References

- [1] A. V. Arhangel'skii, *Some types of factor mappings and the relations between classes of topological spaces*, Soviet Math. Dokl. 4 (1963), 1726-1729.
- [2] J. G. Ceder, *Some generalizations of metric spaces*, Pacific J. Math. 4 (1961), 105-126.
- [3] C. H. Dowker, *Topology of metric complexes*, Amer. J. Math. 74 (1952), 557-577.
- [4] R. Engelking, *General Topology*, Polish Scientific Publishers, Warszawa, 1977.
- [5] S. P. Franklin and B. V. Smith Thomas, *On the metrizable-ability of  $k_\omega$ -spaces*, Pacific J. Math. 72 (1977), 399-402.
- [6] G. Gruenhage and Y. Tanaka, *Products of  $k$ -spaces and spaces of countable tightness*, Transactions of Amer. Math. Soc. 273 (1982), 299-308.
- [7] G. Gruenhage, E. Michael and Y. Tanaka, *Spaces determined by point-countable covers*, Pacific J. Math. 113 (1984), 303-332.
- [8] N. Lašnev, *Continuous decompositions and closed mappings of metric spaces*, Soviet Math. Dokl. 165 (1965), 1504-1506.
- [9] E. Michael, *A quintuple quotient quest*, General Topology and Appl. 2 (1972), 91-138.
- [10] K. Morita, *On spaces having the weak topology with respect to closed coverings*, Proc. Japan Acad. 29 (1954), 537-543.
- [11] \_\_\_\_\_, *On closed mappings*, Proc. Japan Acad. 32 (1956), 539-543.
- [12] Y. Tanaka, *Products of sequential spaces*, Proc. Amer. Math. Soc. 54 (1976), 371-375.
- [13] \_\_\_\_\_, *Point-countable  $k$ -systems and products of  $k$ -spaces*, Pacific J. Math. 101 (1982), 199-208.
- [14] \_\_\_\_\_, *Closed images of locally compact spaces and Fréchet spaces*, Topology Proceedings 7 (1982), 279-292.
- [15] \_\_\_\_\_, *Metrizability of certain quotient spaces*, Fund. Math. 119 (1983), 157-168.

- [16] J. H. C. Whitehead, *Combinatorial homotopy. I*,  
Bull. Amer. Math. Soc. 55 (1949), 213-245.

Tokyo Gakugei University  
Koganei-shi, Tokyo, Japan  
and

Sichuan University  
Chengdu, China (P.R.C.)