
TOPOLOGY PROCEEDINGS



Volume 9, 1984

Pages 201–215

<http://topology.auburn.edu/tp/>

COMPACTIFICATIONS OF THE RAY WITH THE CLOSED ARC AS REMAINDER

by

MARWAN M. AWARTANI

Topology Proceedings

Web: <http://topology.auburn.edu/tp/>

Mail: Topology Proceedings
Department of Mathematics & Statistics
Auburn University, Alabama 36849, USA

E-mail: topolog@auburn.edu

ISSN: 0146-4124

COPYRIGHT © by Topology Proceedings. All rights reserved.

COMPACTIFICATIONS OF THE RAY WITH THE CLOSED ARC AS REMAINDER

Marwan M. Awartani*

0. Introduction and Summary

Let J denote the ray $(0,1]$ and let I denote the closed interval $[0,1]$. If $f: J \rightarrow I$ is a continuous function, let Jf denote the graph of f . Let $\alpha_f J$ denote the closure, \overline{Jf} , of Jf in $I \times I$. Then $\alpha_f J$ is a compactification of J , since the function $h: J \rightarrow \alpha_f J$ given by $h(t) = (t, f(t))$ is a dense embedding of J into $\alpha_f J$. Let $\hat{J}f$ denote the remainder $\alpha_f J \setminus Jf$. In (1) a procedure similar to the above is used to obtain compactifications for a large class of non-compact, locally compact spaces. Techniques using the closure of the graph of a function f are used in (2) to obtain various topological extensions of f .

It is readily seen that if $f: J \rightarrow I$ is continuous and is continuously extendible to I , then $\alpha_f J$ is homeomorphic (\cong) to I (the one point compactification of J). Let F denote the class of all functions $f: J \rightarrow I$, which are continuous but are not continuously extendible to I . If $f \in F$, then $\hat{J}f$ is a closed subinterval of I and $\alpha_f J$ is non-locally connected because Jf is forced to oscillate as it approaches $\hat{J}f$. In (3) and (4), the author and S. Khabbaz develop invariants to study the homeomorphism and the

* This paper was accomplished while the author was a visiting professor at the Department of Mathematics, Cornell University. The author wishes to extend to David Henderson deep appreciation and gratitude for his support.

homotopy types of the spaces $Jf \cup \{(0,0)\}$, where $f \in F$.

Our purpose here is to study the compactifications $\alpha_f J$, where $f \in F$. For related work see (5) and (6).

In Section 1 we associate with each $\alpha_f J$, $f \in F$, a closed ordered subset E_f of $\hat{J}f$, where the order is that induced by the natural order on $\hat{J}f$. E_f is called the type of the compactification $\alpha_f J$ and consists of those points of $\hat{J}f$ arbitrarily close to which Jf makes significant turns. In theorem 1.4, E_f is proved to be a topological invariant of $\alpha_f J$. In Section 2 we prove a reduction theorem that associates with each $\alpha_f J$, another compactification $\alpha_g J$, homeomorphic to $\alpha_f J$, where g is piecewise linear over a sequence V in J converging to 0, and where each $v \in V$, is a local extremum of g . Moreover $\alpha_g J$ has the nice property that $E_g = \overline{Vg} \setminus Vg$, where $Vg = \{(v, g(v)) : v \in V\}$. Hence Vg enjoys some sort of minimality in the sense that it contains no subsequences converging to any point of $\hat{J}g \setminus E_g$.

Finally, in Section 3, we prove that for each closed subset T of I , there exists continuum many nonhomeomorphic compactifications of the ray, all of which have type T .

1. The Invariant E_f

Definition 1.1. Let p, q be two points in Jf . Then $[p, q]_f$ denotes the closed arc in Jf joining p and q . $[p, q]_f$ is called a *wedge* (respectively a *spike*) if the lowest (highest) points of $[p, q]_f$ are all interior points. Such a wedge or spike is called *symmetric* if $\pi_2(p) = \pi_2(q)$,

where π_2 is the projection onto the y-coordinate. Finally if $p, q \in I \times I$, then $[p, q]$ denotes the straight line segment in $I \times I$ joining p and q .

Definition 1.2. [See (7) or (8)]. Let $\{A_i\}$ be a sequence of nonempty closed subsets of $\alpha_f J$. Then define:

(a) $\text{Lim inf}\{A_i\} = \{x \in \alpha_f J: \text{if } U \text{ is an open neighborhood of } x \text{ in } \alpha_f J, \text{ then } U \cap A_i \neq \emptyset \text{ for all but finitely many } i\}$.

(b) $\text{Lim sup}\{A_i\} = \{x \in \alpha_f J: \text{if } U \text{ is an open neighborhood of } x \text{ in } \alpha_f J, \text{ then } U \cap A_i \neq \emptyset \text{ for infinitely many } i\}$.

If $\text{lim inf}\{A_i\} = A = \text{lim sup}\{A_i\}$, then we say that the sequence $\{A_i\}$ converges to A , or $\text{lim}\{A_i\} = A$.

The above definition of convergence is equivalent to convergence with respect to the Hausdorff metric on the set of all nonempty closed subsets of $\alpha_f J$. See for example (8).

Definition 1.3. Let $s \in \hat{J}f$. Then s is called *essential* in $\alpha_f J$, if it satisfies one of the following two conditions:

(i) There exists a sequence $\{[p_i, q_i]_f\}$ of wedges (spikes) in Jf and a positive number ϵ , such that $\text{lim}\{[p_i, q_i]_f\} = [s, s+\epsilon]([s, s-\epsilon])$, and $\text{lim}\{p_i\} = \text{lim}\{q_i\} = s + \epsilon(s - \epsilon)$.

(ii) s is the limit of a sequence of points in $\hat{J}f$ satisfying condition (i). Otherwise s is called *inessential* in $\alpha_f J$. Let Ef denote the set of essential points of $\alpha_f J$, ordered by the natural order on Jf . (In figure 1 $Ef = \{0, \frac{1}{3}, 1\}$.)




Figure 1

Theorem 1.4. Let $f, g \in F$ and let $h: \alpha_f J \rightarrow \alpha_g J$ be a homeomorphism. Then $h|_{E_f}$ is a monotone homeomorphism onto E_g .

The proof of this theorem follows from Lemma 1.6.

First we need the following:

Convention 1.5.

(i) If a statement P is made about the elements of a sequence S , such that all but finitely many elements of S satisfy P , then we say S almost satisfies P , or almost each element of S satisfies P .

(ii) Let $X \subseteq J_f$, then $C(X)$ denotes the set of path components of X ordered from right to left. And J_f is always assumed to have the natural order of the reals. A point of $\alpha_f J$ contained in \hat{J}_f will be referred to by its Y -coordinate.

Lemma 1.6. Let $h: \alpha_f J \rightarrow \alpha_g J$ be a homeomorphism. Then the following hold:

(i) $h|_{\hat{J}_f}$ is a monotone homeomorphism onto \hat{J}_g .

(ii) If $\{[p_i, q_i]_f\}$ is a sequence of wedges converging to $[s, s+\epsilon] \subseteq \hat{J}_f$ with $\lim\{p_i\} = \lim\{q_i\} = s + \epsilon$, and if

$h|_{\hat{J}f}$ is order preserving (reversing), then $\{[h(p_i), h(q_i)]_g\}$ is almost a sequence of wedges (spikes).

(iii) If $\{[p_i, q_i]_f\}$ is a sequence of spikes converging $[s, s-\epsilon] \subseteq \hat{J}f$, with $\lim\{p_i\} = \lim\{q_i\} = s - \epsilon$, and if $h|_{\hat{J}f}$ is order preserving (reversing), then $\{[h(p_i), h(q_i)]_g\}$ is almost a sequence of spikes (wedges).

Proof.

(i) is immediate.

(ii) Since h is a homeomorphism, $h[p_i, q_i]_f = [h(p_i), h(q_i)]_g$ and the sequence $\{[h(p_i), h(q_i)]_g\}$ converges to $[h(s), h(s+\epsilon)]$. For each i , let m_i be a lowest point in $[h(p_i), h(q_i)]_g$, then

$$\lim\{h(p_i)\} = \lim\{h(q_i)\} = h(s+\epsilon) > h(s) = \lim\{m_i\}.$$

Hence almost each m_i is an interior point of $[h(p_i), h(q_i)]_g$. Since m_i was an arbitrary lowest point of $[h(p_i), h(q_i)]_g$, this implies that in almost each $[h(p_i), h(q_i)]_g$ the lowest points are interior points. Hence the desired result follows. The case when $h|_{\hat{J}f}$ is order reversing is handled similarly. The proof of (iii) is similar to that of (ii).

2. A Reduction Theorem

Definition 2.1. Let V be a decreasing sequence of points in J converging to 0, and let $f \in F$. Then f is called *piecewise linear* over $V_f = \{(v, f(v)) : v \in V\}$ if $f|_{[v, v']}$ is linear for each pair, v, v' , of consecutive elements of V . If no ambiguity arises, f is called simply P.L. The set V_f is called the *set of vertices of f* . Moreover if f is P.L. and every $P \in V_f$ is a local extremum, then f is called *sawtooth*. Finally let m_f and M_f denote,

respectively, the local minima and local maxima of f .

Remark 2.2. Let $f \in F$ be P.L. over Vf , then $V = \pi_x(Vf)$ is a copy of the integers (π_x is the projection on the x -coordinate). Hence $f|_V: V \rightarrow I$ also yields a compactification of the integers, $\alpha_f V = \widehat{Vf}$. \widehat{Vf} denotes the remainder $\alpha_f V \setminus Vf$.

In this section we prove the following:

Reduction Theorem 2.3. For each $f \in F$, there exists $g \in F$ having the following properties:

- (i) g is sawtooth
- (ii) $\alpha_f J \cong \alpha_g J$ and $\widehat{Jf} = \widehat{Jg}$
- (iii) $Ef = Eg = \widehat{Vg}$

A particularly nice property of the above sawtooth function g is that $Eg = \widehat{Vg}$. Although it can be predicted from 1.3 and the proof of 1.6 that $Eg \subseteq \widehat{Vg}$, equality is not in general true. In fact, Eg may consist of just two points, whereas \widehat{Vg} may be all of I . So the above theorem implies some sort of minimality about the vertices Vg , in the sense that if a subsequence of Vg converges to a point s , the $s \in Eg$.

Lemma 2.4. Let $f, g \in F$ such that $\lim_{x \rightarrow 0} |f(x) - g(x)| = 0$. Then the function $h: \alpha_f J \rightarrow \alpha_g J$ given by

- (i) $h|_{\widehat{Jf}} = \text{id}$
- (ii) $h(x, f(x)) = (x, g(x))$

is a homeomorphism.

Lemma 2.5. Let $f \in F$ be P.L. Then there exists a sawtooth function $g \in F$, such that $\alpha_f J \cong \alpha_g J$.

Proof. Let V_f be the set of vertices of f , and let $\pi_x(V_f) = V$. Choose $a_1 = (1, f(1))$, and let $a_1 > a_2 > a_3 > \dots$ be a sequence of points in V , such that for each i , f is monotone over $[a_i, a_{i+1}]$ and is not monotone over any subinterval of J properly containing $[a_i, a_{i+1}]$. Clearly $\cup_{i=1}^{\infty} [a_i, a_{i+1}] = J$. Let g be the P.L. function over $\{(a_i, f(a_i))\}$. Then g is sawtooth, since each a_i is a local extremum of g . In order to prove that $\alpha_f J \cong \alpha_g J$, we construct another function $g_1 \in F$ as follows:

(i) $g_1(a_i) = f(a_i) = g(a_i)$ for each i .

(ii) $g_1|_{(a_i, a_{i+1})}$ is a strictly monotone function such that $|f(x) - g_1(x)| < \epsilon_i$, where the sequence $\{\epsilon_i\}$ is a decreasing sequence of positive numbers converging to 0. This is possible because $f|_{(a_i, a_{i+1})}$ is monotone for each i . Since $\lim\{\epsilon_i\} = 0$, it follows that $\lim_{x \rightarrow 0} |f(x) - g_1(x)| = 0$. By Lemma 2.4, $\alpha_f J \cong \alpha_{g_1} J$. Finally define a function

$h: \alpha_{g_1} J \rightarrow \alpha_g J$ as follows:

(i) $h|_{Jg_1} = \text{id}$, since $\hat{J}g_1 = \hat{J}g$.

(ii) Let $P_i = (a_i, g_1(a_i)) = (a_i, g(a_i))$ for each i .

Then h maps $[P_i, P_{i+1}]_{g_1}$ onto $[P_i, P_{i+1}]_g$ by the horizontal projection: $h(x, g_1(x)) = (x', g(x'))$ where $g_1(x) = g(x')$. h is 1-1 on each $[P_i, P_{i+1}]_{g_1}$ because both g_1 and g are strictly monotone over $[a_i, a_{i+1}]$. One readily verifies that h is a homeomorphism.

Proof of Theorem 2.3. We break the proof into steps:

Step 1. Let $g_1 \in F$ be a P.L. function such that $\alpha_f J$ is homeomorphic to $\alpha_{g_1} J$ and where $\pi_X(Vg_1)$ converges to 0. By Lemma 2.5, we may assume that g_1 is sawtooth. It follows from Definition 1.3 that Eg_1 is closed in $\hat{J}g_1$. Hence $\hat{J}g_1 \setminus Eg_1$ is the countable (possibly finite) union of disjoint open intervals (t_i, s_i) , $t_i < s_i$. For each i , choose positive numbers ℓ_i, k_i, r_i , so that $s_i > \ell_i > k_i > r_i > t_i$, and let $U_i = ((t_i, s_i) \times I) \cap Jg_1$.

Step 2. A new function $g_2 \in F$ is obtained from g_1 by altering Jg_1 over each U_i separately. Any portion of Jg_1 which is not altered is assumed to stay as part of Jg_2 . We alter a typical U_i by considering each $K \in C(U_i)$ separately. If $K \in C(U_i)$ lies totally in one of the strips $k_i \leq y \leq s_i$; $t_i \leq y \leq k_i$, then K is left intact. Otherwise, removing the line $y = k_i$ splits K into at least two components. The closure of a typical such component $[p_0, q_0]_{g_1}$ is one of the following types:

(i) a wedge (spike) contained in the strip $k_i \geq y \geq r_i$ ($k_i \leq y \leq \ell_i$). In this case, replace $[p_0, q_0]_{g_1}$ by $[p_0, q_0]$.

(ii) $[p_0, q_0]_{g_1}$ is neither a wedge nor a spike, and is either totally contained in the strip $k_i \leq y \leq s_i$ or the strip $t_i \leq y \leq k_i$. We treat the two cases simultaneously, the alternative choices for the second case being included in parenthesis. We may assume that $\pi_2(p_0) > \pi_2(q_0)$.

Before we alter $[p_0, q_0]_{g_1}$, we obtain a sequence

$\{p_1, \dots, p_j\}(\{q_1, \dots, q_j\})$ of vertices in $[p_0, q_0]_{g_1}$ such

that for each $k, 1 \leq k \leq j, p_k (q_k)$ is a highest (lowest) vertex in the interior of $[p_{k-1}, q_0]_{g_1} ([q_{k-1}, p_0]_{g_1})$ and such that $[p_j, q_0]_{g_1} ([q_j, p_0]_{g_1})$ contains no vertices of g_1 . Also, for each $k, 1 \leq k \leq j, [p_k, Q_k]_{g_1} ([q_k, Q_k]_{g_1})$ is chosen to be the largest symmetric wedge (spike) in $[p_k, p_0]_{g_1} ([q_k, q_0]_{g_1})$ having $p_k (q_k)$ as one endpoint. Now $[p_0, q_0]_{g_1}$ is altered by replacing each $[p_k, Q_k]_{g_1} ([q_k, Q_k]_{g_1})$ by $[p_k, Q_k] ([q_k, Q_k])$.

(iii) Finally if $[p_0, q_0]_{g_1}$ is a wedge (respectively, spike) that extends below the line $y = r_i$ (above the line $y = \ell_i$), then choose a lowest (highest) vertex M in $[p_0, q_0]_{g_1}$. The arcs $[p_0, M]_{g_1}, [M, q_0]_{g_1}$ are both the types discussed in (ii) and are altered accordingly.

Step 3. $\lim_{x \rightarrow 0} |g_1(x) - g_2(x)| = 0$. To establish this, let $\{[p_i, q_i]_{g_1}\}$ be an enumeration from right to left of all the wedges in Jg_1 that have been altered in a particular U_j . And let $\{[p'_i, q'_i]_{g_1}\}$ be the similar sequence of all the spikes. Our claim is equivalent to showing that each of $\{[p_i, q_i]_{g_1}\}$ and $\{[p'_i, q'_i]_{g_1}\}$ converges to a point. Suppose $\{[p_i, q_i]_{g_1}\}$ does not converge to a point, then one deduces that a point of $[k_j, s_j)$ is essential. Similarly if $\{[p'_i, q'_i]_{g_1}\}$ does not converge to a point, then a point of $(t_j, k_j]$ is essential. Both of these conclusions contradict the assumption that (t_j, s_j) consists of inessential points.

Step 4. Applying Lemma 2.5 to g_2 we obtain the desired sawtooth function g . To see this let $h_1: \alpha_{f^J} \rightarrow \alpha_{g_1^J}$, $h_3: \alpha_{g_2^J} \rightarrow \alpha_g^J$ be the homeomorphisms guaranteed by Lemma 2.5, and let $h_2: \alpha_{g_1^J} \rightarrow \alpha_{g_2^J}$ be the homeomorphism guaranteed by Lemma 2.4. Then the composition $h_3 \cdot h_2 \cdot h_1: \alpha_{f^J} \rightarrow \alpha_g^J$ is a homeomorphism. Also $h|_{\hat{J}f} = \text{id}$. This and Theorem 1.3 establish (i), (ii) and half of (iii) of Theorem 2.3.

Finally to prove that $\hat{V}g = Eg$, we notice that the sequence $Vg \cap ([k_j, s_j] \times I)$ converges to s_j and the sequence $Vg \cap ((t_j, k_j] \times I)$ converges to t_j , otherwise a point of $[k_j, s_j]$ or $(t_j, k_j]$ would be essential.

3. Homeomorphism Classes of Compactifications of a Given Type

The type Ef of a compactification α_{f^J} is not a complete invariant. In this section, we prove the following:

Theorem 3.1. Let T be a closed subset of I containing more than two points. Then there exists continuum many nonhomeomorphic compactifications of the ray all of which are of type T .

In the case where the type Ef consists of two points, using a procedure similar to that of Theorem 2.3, one obtains a reduction that "straightens" Jf enough so that the resulting graph looks like that in figure 2. Hence all compactifications of J whose type consists of two points are homeomorphic to $\alpha_{\sin \frac{1}{x}}^J$.




Figure 2

The strategy for proving the above theorem involves associating (in 3.3) with each nonempty set A of positive integers, a function $f \in F$ where $\alpha_f J$ is of type T , and then proving that functions associated with different sets of positive integers yield nonhomeomorphic compactifications, all of which are of type T .

Before we proceed with the proof of Theorem 3.1, we need the following:

Definition 3.2. Let S be a totally ordered sequence. A *block* in S is a finite set of consecutive elements of S . If the cardinality of b , $|b| = n$, then b is an n -*block* in S . The *boundary of b in S* , $bd_S(b)$, is a subset of $S \setminus b$ consisting of two elements: the element preceding the first element of b , and that succeeding the last element of b . A subsequence S_1 of S is called an n -*subsequence* of S , for a positive integer n , if $S_1 = \cup_{i=1}^{\infty} b_i$, such that almost each b_i is an n -block in S , with $bd_S(b_i) \subset S \setminus S_1$. The *boundary of S_1 in S* , $bd_S(S_1) = \cup_{i=1}^{\infty} bd_S(b_i)$.

Observe that if S_1 is an n -subsequence of S and $m \neq n$, then S_1 is not an m -subsequence of S .

Construction 3.3. Let T be the set specified in Theorem 3.1 and let t, t' be respectively the smallest and largest elements of T , and let D be a countable dense subset of $T \setminus \{t'\}$, containing t . Given a nonempty set A of positive integers we associate with A a sawtooth function $f \in F$ whose vertices V_f satisfy the following conditions:

- (i) All elements of M_f are at height t' .
- (ii) For each $d \in D$, and each $a \in A$, the minima of f at height d , denoted by m_d , contain an a -subsequence S of m_f with $bd_{m_f}(S) \subseteq mt$.
- (iii) Moreover, if b is a block in m_f contained in m_d , with $bd_{m_f}(b) \subseteq mt$, then $|b| \in A$.

To illustrate the above construction, let $T = \{0, \frac{1}{3}, \frac{2}{3}, 1\}$: $A = \{1, 2\}$. The function in figure 3 satisfies the conditions of Construction 3.3




Figure 3

Lemma 3.4. Let f be the function constructed in 3.3, and let S be an n -subsequence of m_f which converges to $d \in D \setminus \{t\}$, and whose boundary, $bd_{m_f}(S)$, converges to t . Then $n \in A$.

Proof. Since S is an n -subsequence of m_f , $S = \bigcup_{i=1}^{\infty} b_i$, where almost each b_i is an n -block in m_f and $bd_{m_f}(b_i) \subseteq m_f \setminus S$.

Since $\lim S \neq t$, almost each $b_i \subseteq mf \setminus mt$. It follows then from part (ii) of construction 3.3, that an element of $bd_{mf}(b_i)$ is either at the same height as elements of b_i , or is contained in mt . Since $bd_{mf}(S)$ converges to $t \neq d$, it follows that almost each $bd_{mf}(b_i)$ is contained in mt . Hence part (iii) of Construction 3.3 implies that $n \in A$.

Proof of Theorem 3.1. Let A and B be two sets of positive integers, and let f, g be the function associated with A, B respectively. Observe that $\hat{J}f = \hat{J}g = [t, t']$. Suppose that $A \neq B$, and suppose that $h: \alpha_f J \rightarrow \alpha_g J$ is a homeomorphism. We may assume that there exists an $a \in A \setminus B$. Choose $d \in D \setminus \{t\}$, and let $U = ([t, t'] \times I) \cap Jf$. Then by part (ii) of Construction 3.3, there exists a sequence S in $C(U)$ having the following properties:

- (i) S is an a -subsequence of $C(U)$ converging to $[d, t']$.
- (ii) $bd_{C(U)}(S)$ converges to $[t, t']$.
- (iii) each $K \in C(U)$ contains exactly one element of mf .

Since h is a homeomorphism, it is an order isomorphism taking $C(U)$ onto $C(h(U))$. Now we verify the following:

a) $h|_{\hat{J}f}$ is order preserving. Suppose the contrary and consider the above sequence $S \subseteq C(U)$. Since $h|_{\hat{J}f}$ is order reversing, part (iii) of Lemma 1.6 implies that almost each $L \in h(S)$ is a spike. Consequently almost each L contains at least one element of Mg . Since h is a homeomorphism and since $h(t') = t, h(S)$ must converge to $[t, h(d)]$. But this contradicts the fact that $t' \in \lim h(S)$ since almost each $L \in h(S)$ contains at least one vertex of Mg , and Mg converges to t' .

b) It follows from (a) that $h(S)$ is an a -subsequence of $C(h(U))$ converging to $[h(d), t']$ and $\text{bd}_{C(h(U))}(h(S))$ converges to $[t, t']$.

c) Almost each element L of $h(S)$ or $\text{bd}_{C(h(U))}(h(S))$ contains exactly one element of mg . First, it follows from part (ii) of Lemma 1.6 and (a) above that almost each L is a wedge, and hence contains at least one element of mg . Suppose that S_1 is a subsequence of $h(S)$ such that each element of S_1 contains at least two elements of mg . Then each such element L must contain at least one element M of Mg , which when deleted from L splits L into two arcs whose closures are both wedges. Whereas if $h^{-1}(M)$ is deleted from $h^{-1}(L) \in S$, then clearly $h^{-1}(L)$ breaks up into two arcs such that the closure of at most one of them is a wedge. This contradicts part (ii) of Lemma 1.6. Similarly, we prove that almost each $L \in \text{bd}_{C(h(U))}(h(S))$ contains exactly one element of mg .

d) It follows from (b) and (c) above that the sequence $h(S) \cap mg$ is an a -subsequence of mg which converges to $h(d)$ and whose boundary in mg converges to t . Hence by Lemma 3.4, $a \in B$, contradicting the fact that a was chosen in $A \setminus B$, and hence proving that α_f^J and α_g^J are nonhomeomorphic.

References

- (1) A. K. Steiner and E. F. Steiner, *Compactifications as closures of graphs*, *Fundamenta Mathematicae* (1968), 221-223.
- (2) F. A. Delahan and G. E. Strecker, *Graphic extensions of mappings*, *Quaestiones Mathematicae* 2 (1977), 401-417.

- (3) M. M. Awartani and S. A. Khabbaz, *On almost continuous functions*, Proceedings of the Texas Topology Symposium (1980), 221-228.
- (4) _____, *On the topology of graphs of discontinuous functions on the unit interval* (to appear).
- (5) Sam B. Nadler, Jr., *Arc continua*, Canadian Mathematical Bulletin 14 (2) (1971), 183-189.
- (6) David P. Bellamy, *An uncountable collection of chainable continua*, Transactions of the American Mathematical Society 16 (October 1971), 297-303.
- (7) Gordon Thomas Whyburn, *Analytic topology*, American Mathematical Society Colloquium Publications 28 (1942).
- (8) Sam B. Nadler, Jr., *Hyperspaces of sets*, Marcel Dekker, Inc. (1978).

Birzeit University, Birzeit

West Bank

via Israel