


Topology Proceedings



Web: <http://topology.auburn.edu/tp/>

Mail: Topology Proceedings

Department of Mathematics & Statistics

Auburn University, Alabama 36849, USA

E-mail: topolog@auburn.edu

ISSN: 0146-4124

COPYRIGHT © by Topology Proceedings. All rights reserved.

A Note on Property III in Generalized Ordered Spaces

Harold R Bennett and David J. Lutzer

Abstract

A topological space X has Property III provided there are subsets $U(k)$ and $D(k)$ of X for each $k \geq 1$ such that:

- 1) $U(k)$ is open in X and $D(k)$ is relatively closed in $U(k)$;
- 2) $D(k)$ is discrete in itself;
- 3) if p is a point of an open set G , then there is a $k \geq 1$ such that $p \in U(k)$ and $D(k) \cap G \neq \emptyset$.

We show that Property III is exactly what is needed, in a generalized ordered space, to transform a $\delta\theta$ -base in the sense of Aull into a σ -disjoint base (equivalently, into a quasi-development). We also give an example of a linearly ordered topological space that has a $\delta\theta$ -base, but not a point-countable base.

1 Introduction

A topological space X has Property III provided there are subsets $U(k)$ and $D(k)$ of X for each $k \geq 1$ such that:

- 1) $U(k)$ is open in X and $D(k)$ is relatively closed in $U(k)$;
- 2) $D(k)$ is discrete in itself, i.e., when endowed with the relative topology from X , the space $D(k)$ is discrete;
- 3) if p is a point of an open set G , then there is a $k \geq 1$ such that $p \in U(k)$ and $D(k) \cap G \neq \emptyset$.

In [BL] we showed that Property III plays an important bridging role in the theory of generalized ordered spaces by proving that a generalized ordered space X has a σ -disjoint base if and only if it has a point-countable base and has Property III. In this paper we extend the main theorem of [BL] by showing that Property III plays an even larger bridging role in ordered space theory. We show that for generalized ordered spaces, Property III is the solution of the equation “ σ -disjoint base = (?) + $\delta\theta$ -base” by adding (e) to the list of equivalent conditions in the following theorem:

1.1) Theorem *The following properties of a generalized ordered space X are equivalent:*

- a) X is quasi-developable;
- b) X has a σ -disjoint base;
- c) X has a σ -point finite base;
- d) X has a point countable base and has Property III;
- e) X has a $\delta\theta$ -base and has Property III.

Recall that a generalized ordered space or GO-space is a triple $(X, \mathcal{T}, <)$ such that $(X, <)$ is a linearly ordered set and where \mathcal{T} is a Hausdorff topology on X that has a base of order-convex open sets. If \mathcal{T} is the open interval topology of $<$, then

then X is a linearly ordered topological space or LOTS. The class of GO-spaces is exactly the class of subspaces of LOTS.

Quasi-developments, σ -point finite bases, and σ -disjoint bases are familiar objects: see [B1], [BL] and [L1] for general background material. The notion of a $\delta\theta$ -base was introduced by Aull in [Au]: A collection $\mathcal{B} = \bigcup\{\mathcal{B}(n) : n \geq 1\}$ is a $\delta\theta$ -base for X provided that if G is open and $p \in G$, then for some n , some $B \in \mathcal{B}(n)$ has $p \in B \subset G$, and the order of p in $\mathcal{B}(n)$ is countable. (By the order of a point p in a collection \mathcal{C} we mean the cardinality of the collection $\{C \in \mathcal{C} : p \in C\}$. We will denote the order of p in \mathcal{C} by $\text{ord}(p, \mathcal{C})$.) Clearly, any point-countable base is a $\delta\theta$ -base, but the converse is false, even among GO-spaces, as can be seen from the example constructed in Section 3.

Throughout this paper, we must carefully distinguish between the statements “ E is a discrete subspace of X ” and “ E is a closed discrete subspace of X . We will do this, for example, by writing “Suppose X has a σ -closed discrete dense subset” even though that makes our terminology more cumbersome. For example, it is easy to see that the usual space of countable ordinals has a σ -discrete dense subspace but no dense subspace that is σ -closed-discrete.

2 Property III, $\delta\theta$ -bases, and perfect GO spaces

We begin by recalling two technical results from [BL] and a characterization of hereditary paracompactness that follows immediately from [Fa].

2.1) Lemma *Let X be a GO-space with Property III. Then:*

- a) *Every subspace of X also has Property III;*
- b) *X is hereditarily paracompact.*

2.2) Lemma Suppose $\{C(\alpha) : \alpha \in A\}$ is a pairwise disjoint collection of open subsets of a first-countable GO-space X . For each α , suppose that $E(\alpha)$ is a discrete-in-itself subspace of $C(\alpha)$. Then there is a σ -disjoint collection \mathcal{B} of open subsets of X that contains a base at each point of $\bigcup\{E(\alpha) : \alpha \in A\}$.

2.3) Lemma A GO-space X is hereditarily paracompact if and only if for each subspace $C \subset X$ there are sets E and E' such that:

- a) E and E' are relatively closed subsets of C that are discrete;
- b) E is well-ordered by the given ordering of X , and E' is reverse-well-ordered by the given ordering of X ;
- c) if $p \in C$ then $p \in [e, e']$ for some $e \in E$ and $e' \in E'$.

2.4) Proposition Suppose that X is a GO-space with a $\delta\theta$ -base. If X has Property III, then X has a σ -disjoint base.

Proof: Let $\mathcal{B} = \bigcup\{\mathcal{B}(n) : n \geq 1\}$ be a $\delta\theta$ -base for X . We may assume that members of \mathcal{B} are convex subsets of X . Then X is certainly first countable. For each n let $X(n) = \{p \in X : \text{ord}(p, \mathcal{B}(n)) \leq \omega\}$. Because X is first countable, it is easy to show that the set $X(n)$ is a relatively closed subset of the open set $\bigcup \mathcal{B}(n)$. For $k \geq 1$, let $U(k)$ and $D(k)$ be the sets given by Property III. Let $O(n, k) = U(k) \cap (\bigcup \mathcal{B}(n))$. Then $O(n, k) - D(k)$ is open in $U(k)$ and hence also in X , so that $O(n, k) - D(k)$ is open in X . Let $\{C(n, k, \alpha) : \alpha \in A(n, k)\}$ be the family of all convex components of $O(n, k) - D(k)$. In addition, notice that $X(n) \cap O(n, k)$ is a relatively closed subset of $O(n, k)$ so that for each $\alpha \in A(n, k)$, the set $X(n) \cap C(n, k, \alpha)$ is a relatively closed subset of $C(n, k, \alpha)$. Let $A^*(n, k) = \{\alpha \in A(n, k) : X(n) \cap C(n, k, \alpha) \neq \emptyset\}$ and for each $\alpha \in A^*(n, k)$ use (2.3) to

find discrete, relatively closed subsets $E(n, k, \alpha), E'(n, k, \alpha)$ of $X(n) \cap C(n, k, \alpha)$ as described in (2.3).

For each $\alpha \in A^*(n, k)$ let $\mathcal{H}(n, k, \alpha)$ be the collection of all convex components of the open set $C(n, k, \alpha) - E(n, k, \alpha)$ that have at least one endpoint in the set $E(n, k, \alpha)$. Define $\mathcal{H}'(n, k, \alpha)$ in an analogous way, using $E'(n, k, \alpha)$ in place of $E(n, k, \alpha)$. For each $H \in \mathcal{H}(n, k, \alpha)$, define $\mathcal{G}(n, k, \alpha, H)$ to be the collection of all sets $H \cap B$ where $B \in \mathcal{B}(n)$ and B contains at least one of the endpoints of H that belongs to $E(n, k, \alpha)$. Define collections $\mathcal{G}(n, k, \alpha, H')$ using the sets $E'(n, k, \alpha)$ in place of $E(n, k, \alpha)$ and sets $H' \in \mathcal{H}'(n, k, \alpha)$.

For a fixed (n, k, α, H) , if $B \cap H \in \mathcal{G}(n, k, \alpha, H)$ then B contains at least one end point of H that belongs to $E(n, k, \alpha) \subset X(n)$, and $B \in \mathcal{B}(n)$. Because there are at most two such endpoints, the collection $\mathcal{G}(n, k, \alpha, H)$ is countable. Index the family $\mathcal{G}(n, k, \alpha, H)$ as $\{G(n, k, \alpha, H, j) : j \geq 1\}$. Similarly, index $\mathcal{G}'(n, k, \alpha, H')$ as $\{G'(n, k, \alpha, H', j) : j \geq 1\}$

Let $\mathcal{U}(n, k, j) = \{G(n, k, \alpha, H, j) : \alpha \in A^*(n, k) \text{ and } H \in \mathcal{H}(n, k, \alpha)\}$. If α and β are distinct members of $A^*(n, k)$ then $G(n, k, \alpha, H_1, j)$ and $G(n, k, \beta, H_2, j)$ are, respectively, subsets of the disjoint convex components $C(n, k, \alpha)$ and $C(n, k, \beta)$, for any $H_1 \in \mathcal{H}(n, k, \alpha)$ and $H_2 \in \mathcal{H}(n, k, \beta)$. And if $H_1 \neq H_2$ belong to $\mathcal{H}(n, k, \alpha)$, then $G(n, k, \alpha, H_i, j) \subset H_i$ so that $G(n, k, \alpha, H_1, j) \cap G(n, k, \alpha, H_2, j) = \emptyset$. Therefore each $\mathcal{U}(n, k, j)$ is a pairwise disjoint collection. Analogously, $\mathcal{U}'(n, k, j)$, the collection of all sets $G'(n, k, \alpha, H', j)$ where $\alpha \in A^*(n, k)$ and $H' \in \mathcal{H}'(n, k, \alpha)$, is a pairwise disjoint collection. Let $\mathcal{U} = \bigcup \{\mathcal{U}(n, k, j) : n, k, j \geq 1\}$ and define $\mathcal{U}' = \bigcup \{\mathcal{U}'(n, k, j) : n, k, j \geq 1\}$. Then \mathcal{U} and \mathcal{U}' are σ -disjoint collections of open subsets of X .

Let $Y = (\bigcup \{D(k) : k \geq 1\}) \cup (\bigcup \{E(n, k, \alpha) \cup E'(n, k, \alpha) : n, k \geq 1 \text{ and } \alpha \in A^*(n, k)\})$ Repeated application of (2.2) yields a σ -disjoint collection \mathcal{E} of open subsets of X that contains a base at each point $y \in Y$. Thus it will be enough to show that

the collection $\mathcal{U} \cup \mathcal{U}'$ defined above contains a base at each point of $X - Y$.

To that end, suppose $p \in X - Y$ and $p \in G$, where G is open in X . Find n such that $\text{ord}(p, \mathcal{B}(n))$ is countable, and such that for some $B \in \mathcal{B}(n)$, we have $p \in B \subset G$. Then $p \in X(n)$. Next find some k such that $p \in U(k)$ and some point $d \in D(k) \cap B$. Because $d \in Y$ while $p \notin Y$, we have $p \neq d$. There are two cases to consider. If $p < d$, then we will show that \mathcal{U} contains a set that contains p and is a subset of $B \subset G$. If $d < p$ then an analogous argument (which we leave to the reader) shows that \mathcal{U}' contains the required set.

Because $p \in U(k) \cap (\bigcup \mathcal{B}(n)) = O(n, k)$ and $p \notin D(k)$ there is a unique $\alpha \in A(n, k)$ such that $p \in C(n, k, \alpha)$. Because $p \in C(n, k, \alpha) \cap X(n)$, it follows that $\alpha \in A^*(n, k)$ so that $E(n, k, \alpha)$ is defined. From $d \notin C(n, k, \alpha)$ it follows that either $C(n, k, \alpha) \subset]\leftarrow, d[$ or else $C(n, k, \alpha) \subset]d, \rightarrow[$. Because $p \in C(n, k, \alpha)$ and $p < d$, we have $C(n, k, \alpha) \subset]\leftarrow, d[$. If p is the largest element of the set $X(n) \cap C(n, k, \alpha)$, then $\{p\} = E(n, k, \alpha) \subset Y$, contrary to $p \notin Y$. Therefore some $q \in E(n, k, \alpha)$ has $p < q$. We may assume that q is the first such point. Then q is the right endpoint of the unique convex component H of $C(n, k, \alpha) - E(n, k, \alpha)$ that contains p . Therefore $H \in \mathcal{H}(n, k, \alpha)$.

Because $B \in \mathcal{B}(n)$ is a convex set containing both p and d , and because $p < q$ and $q \in E(n, k, \alpha) \subset C(n, k, \alpha) \subset]\leftarrow, d[$, we know that B contains q . Therefore, $B \cap H \in \mathcal{G}(n, k, \alpha, H)$ so that for some j we have $B \cap H \in \mathcal{U}(n, k, j) \subset \mathcal{U}$, and $p \in B \cap H \subset B \subset G$ as required. The case where $d < p$ being analogous, we now know that X has a σ -disjoint base. \square

3 Examples

In this section we construct a LOTS Z with a $\delta\theta$ -base that does not have a point-countable base. In the light of (2.4), Z

cannot have Property III. The space Z is an extension of an example in [B] of a LOTS with a point-countable base, but not a σ -disjoint base.

3.1) Construction Let P , Q , and R denote, respectively, the sets of irrational, rational, and real numbers. For each limit ordinal $\mu \leq \omega_1$, let $X(\mu)$ be the set of all functions $f : [0, \mu] \rightarrow R$ such that $f(\alpha) \in P$ whenever $\alpha < \mu$ and $f(\mu) \in Q$. Let $X = \bigcup\{X(\mu) : \mu < \omega_1\}$ and let $Y = X(\omega_1)$. Let $Z = X \cup Y$ and order Z lexicographically, i.e., for distinct $f, g \in Z$ define $f \prec g$ if $f(\alpha) < g(\alpha)$ where α is the first ordinal such that $f(\alpha) \neq g(\alpha)$. Let Z have the lexicographic order topology of \prec . For each $f \in Z$ let $\lambda(f)$ be the largest member of the domain of f . For each $f \in Z$ and each $n \geq 1$ define $B(f, n)$ to be the set of all $g \in Z$ such that $\lambda(g) \geq \lambda(f)$ and

$$g(\alpha) = f(\alpha) \text{ for each } \alpha < \lambda(f) \text{ and } |g(\lambda(f)) - f(\lambda(f))| < 1/n.$$

3.2) Lemma *With notation as above,*

- a) *for each $f \in Z$, $\{B(f, n) : n \geq 1\}$ is a countable base of open neighborhoods for f in the space Z ;*
- b) *the collection $\mathcal{B} = \{B(f, n) : f \in X \text{ and } n \geq 1\}$ is point-countable at each point $g \in X$;*
- c) *the collection $\mathcal{C} = \{B(f, n) : f \in Y \text{ and } n \geq 1\}$ is point-countable at each $g \in Z$;*
- d) *Z has a $\delta\theta$ -base.*

Proof: See [B2] for a proof of a) and b). As for c), suppose $g \in Z$. If $\lambda(g) < \omega_1$, then $g \notin \bigcup \mathcal{C}$. If $\lambda(g) = \omega_1$ and if $g \in B(f, n) \in \mathcal{C}$, then $f(\alpha) = g(\alpha)$ for each $\alpha < \omega_1$, and $f(\omega_1)$ is one of the countably many numbers in Q . Hence g lies in at

most countably many members of \mathcal{C} . Combining assertions a), b), and c) gives d). \square

3.3) Lemma *Suppose $f \in X$ and $n \geq 1$ are given. Then there is a function $g \in X$ such that:*

- a) $\lambda(g) = \lambda(f) + \omega$
- b) $B(g, 1)$ is a proper subset of $B(f, n)$.

Proof: Write $\lambda = \lambda(f)$ and choose an irrational number p such that $|f(\lambda) - p| < 1/n$. Define g on $[0, \lambda + \omega]$ by:

- 1) $g(\alpha) = f(\alpha)$ if $\alpha < \lambda$;
- 2) $g(\lambda) = p$;
- 3) $g(\alpha) = \pi$ if $\lambda < \alpha < \lambda + \omega$;
- 4) $g(\lambda + \omega) = 3.14$.

Then $B(g, 1) \subset B(f, n)$ as required, and the inclusion is proper because $f \in B(f, n) - B(g, 1)$. \square

3.4) Lemma *The space Z does not have a point-countable base.*

Proof: For contradiction, suppose that \mathcal{D} is a point- countable base of open sets for Z . Let $\{\mu_\alpha : \alpha < \omega_1\}$ be an increasing well-ordering of all limit ordinals less than ω_1 . Define a function $f_0 : [0, \omega] \rightarrow R$ by the rule that $f_0(\alpha) = \pi$ for each $\alpha < \omega$ and $f_0(\omega) = 3.14$. Then $f_0 \in X$ has $\lambda(f_0) = \omega = \mu_0$. Choose any $D(0) \in \mathcal{D}$ with $f_0 \in D(0)$. Find n_0 such that $B(f_0, n_0) \subset D(0)$ and then use (3.3) to find a function $g_0 : [0, \lambda(f_0) + \omega] \rightarrow R$ such that $B(g_0, 1)$ is a proper subset of $B(f_0, n_0)$.

For induction hypothesis, suppose that $\alpha < \omega_1$ and that for each $\beta < \alpha$ we have defined functions f_β , g_β , an integer $m_\beta \geq 1$, and a set $D(\beta) \in \mathcal{D}$ such that:

- 1) $f_\beta, g_\beta \in X$ and $\lambda(g_\beta) = \lambda(f_\beta) + \omega > \lambda(f_\beta) \geq \mu_\beta$;
- 2) $N(g_\beta, 1) \subset N(f_\beta, m_\beta) \subset D(\beta)$, with each inclusion being proper;
- 3) if $\gamma < \beta < \alpha$ then $D(\beta) \subset N(g_\gamma, 1)$.

There are two cases to consider.

Case 1: Suppose that α is not a limit ordinal, say $\alpha = \beta + 1$. Then g_β is defined and we let $f_\alpha = g_\beta$. Then $f_\alpha \in N(g_\beta, 1)$ so that there is some $D(\alpha) \in \mathcal{D}$ such that $f_\alpha \in D(\alpha) \subset N(g_\beta, 1)$ with the last inclusion being proper. Choose m_α so that $N(f_\alpha, m_\alpha) \subset D(\alpha)$. Use Lemma (3.3) to find g_α with $\lambda(g_\alpha) = \lambda(f_\alpha) + \omega$ and $N(g_\alpha, 1) \subset N(f_\alpha, m_\alpha)$. Observe that $\lambda(g_\alpha) > \lambda(f_\alpha) = \lambda(g_\beta) = \mu_\beta + \omega = \mu_\alpha$. Thus, the induction continues across non-limit ordinals.

Case 2: Suppose α is a limit ordinal less than ω_1 . Let $\lambda = \sup\{\lambda(f_\beta) : \beta < \alpha\}$. Then λ is a limit ordinal with $\lambda \geq \lambda(f_\beta) \geq \mu_\beta$ for each $\beta < \alpha$ so that $\lambda \geq \mu_\alpha$. Because $[0, \lambda] = \{\lambda\} \cup (\bigcup\{[0, \lambda(f_\beta)] : \beta < \alpha\})$, we may define $f_\alpha \in X$ by

- a) $f_\alpha(\delta) = f_\beta(\delta)$ if $\delta < \lambda(f_\beta)$ for some $\beta < \alpha$;
- b) $f_\alpha(\lambda) = 3.14$.

Observe that for each $\beta < \alpha$ we have $f_\alpha \in N(f_\alpha, 1) \subset N(g_\beta, 1) \subset D(\beta)$, with both inclusions being proper. Now find $D(\alpha) \in \mathcal{D}$ such that $f_\alpha \in D(\alpha) \subset N(f_\alpha, 1)$ and then an integer m_α such that $N(f_\alpha, m_\alpha) \subset D(\alpha)$. Use (3.3) to find $g_\alpha \in X$ such that $\lambda(g_\alpha) = \lambda(f_\alpha) + \omega$ and $N(g_\alpha, 1) \subset N(f_\alpha, m_\alpha) \subset D(\alpha)$. Thus, the induction continues across limit ordinals.

The above induction produces a subset $\{f_\alpha : \alpha < \omega_1\}$ of X and distinct sets $D(\alpha) \in \mathcal{D}$ for $\alpha < \omega_1$. Define $h : [0, \omega_1] \rightarrow R$ by

$$g(\delta) = f_\alpha(\delta) \text{ whenever } \delta < \lambda(f_\alpha) \text{ and } \alpha < \omega_1; g(\omega_1) = 3.14.$$

Then $g \in Z$ and $g \in D(\alpha)$ for each $\alpha < \omega_1$. Because the sets $D(\alpha)$ are distinct members of \mathcal{D} , it follows that \mathcal{D} cannot be point-countable in Z . \square

References

- [Au] Aull, C., *Quasi-developments and $\delta\theta$ -bases*, J London Math. Soc., **32** (1974), 197-204.
- [B1] Bennett, H., *Point-countability in linearly ordered spaces*, Proc. Amer. Math. Soc., **28** (1971), 598-606.
- [B2] Bennett, H., *On quasi-developable spaces*, General Topology and its Applications, **1** (1971), 253-262.
- [BL] Bennett, H., and Lutzer, D., *Point countability in generalized ordered spaces*, to appear in Topology and its Applications.
- [Fa] Faber, M., *Metrizability in generalized ordered spaces*, MC Tracts, no. 53, Mathematical Center, Amsterdam, 1974.
- [L] Lutzer, David J., *On generalized ordered spaces*, Dissertationes Math., **89** (1971), 1-42.
- [vW] van Wouwe, J., *GO-spaces and generalizations of metrizability*, MC Tract no. 104, Mathematical Centre, Amsterdam, 1979.

Texas Tech University
Lubbock, TX79409

College of William and Mary
Williamsburg, VA 23187