

TOPOLOGY PROCEEDINGS



Volume 1, 1976

Pages 17–28

<http://topology.auburn.edu/tp/>

ON MONOTONE RETRACTS, ACCESSIBILITY, AND SMOOTHNESS IN CONTINUA

by

G. R. GORDH, JR. AND LEWIS LUM

Topology Proceedings

Web: <http://topology.auburn.edu/tp/>

Mail: Topology Proceedings
Department of Mathematics & Statistics
Auburn University, Alabama 36849, USA

E-mail: topolog@auburn.edu

ISSN: 0146-4124

COPYRIGHT © by Topology Proceedings. All rights reserved.

ON MONOTONE RETRACTS, ACCESSIBILITY, AND SMOOTHNESS IN CONTINUA

G. R. Gordh, Jr. and Lewis Lum

1. Introduction

Consider the following conditions which a continuum M may satisfy.

- (*) Each subcontinuum of M is a monotone retract of M .
- (**) (Pointed version) Each subcontinuum of M which contains a fixed point p is a monotone retract of M .

It is easy to verify that dendrites satisfy both conditions (see [10], Theorem 2.1). The second author has proved that if M is a dendroid, then each of (*) and (**) implies that M is a dendrite ([10], Theorem 2.3, and [12], Theorem 3). More recently, the authors have obtained the same conclusion for arbitrary metric continua satisfying (*), and for arcwise connected metric continua satisfying (**) [6].

In particular, (*) and (**) are equivalent for arcwise connected continua. However, they are not equivalent in general since the familiar "sin $1/x$ curve" satisfies (**).

Thus it is natural to ask for a characterization of continua satisfying (**). The main purpose of this paper is to provide such a characterization.

*Theorem. A continuum M satisfies (**) if and only if*

- (a) M is smooth at p , and
- (b) for each subcontinuum N of M containing p , N is accessible and the components of $M - N$ form a null family.

In this result "smoothness" refers to the concept introduced by the first author in [4]. A more general definition

of "smoothness" has been studied by T. Maćkowiak [14].

It is also shown that a metric continuum M satisfying (**) becomes a dendrite under the canonical monotone decomposition \mathcal{D} of smooth continua defined in [4]. Thus condition (*) is recovered in the decomposition space M/\mathcal{D} .

2. Definitions and Preliminary Remarks

A *continuum* is a compact connected Hausdorff space. The reader is referred to [7] for basic properties of continua and undefined terms.

A subcontinuum N of a continuum M is called a *monotone retract* of M if there exists a mapping $r: M \rightarrow N$ which is both monotone and a retraction.

Let X be a subset of a continuum M . A point $x \in X$ is said to be *accessible* from a point $y \in M - X$ if there exists a subcontinuum H such that $y \in H$ and $H \cap X = \{x\}$. If some point of X is accessible from some point of $M - X$, then X is called *accessible*.

A collection \mathcal{C} of subsets of a continuum M will be called a *null family* if each convergent net C_n of elements of \mathcal{C} which is not eventually constant has a degenerate limit.

A continuum M is *irreducible* from the point p to the point q if no proper subcontinuum of M contains p and q . If, in addition, no proper connected subset of M contains p and q , then M is called an *arc* (sometimes generalized arc or ordered continuum).

The continuum M is *hereditarily unicoherent at p* if for each pair of subcontinua H and K containing p , $H \cap K$ is connected; or equivalently, if for each q in $M - \{p\}$, there is a unique subcontinuum, denoted by pq , which is irreducible from p to q . If M is hereditarily unicoherent at p and for each convergent net q_n , $\lim q_n = q$ implies that the net of subcontinua pq_n converges to pq , then M is said to be *smooth at p* [4].

A *tree (dendrite)* is a locally connected, hereditarily unicoherent (metric) continuum. A *generalized tree (smooth dendroid)* is an arcwise connected, hereditarily unicoherent, smooth (metric) continuum.

Let M be a continuum which is hereditarily unicoherent at the point p . The *weak cutpoint order* on M with respect to p will be denoted by \leq (i.e., $x \leq y$ if $px \subseteq py$). For each $x \in M$ the set $D(x) = \{y \in M: py = px\}$ is the *level set* of x relative to \leq . The collection \mathcal{D} of all level sets forms a decomposition (not necessarily upper semicontinuous) of M . Let $\phi: M \rightarrow M/\mathcal{D}$ denote the natural mapping where M/\mathcal{D} is given the quotient topology. Observe that for any subcontinuum N of M which contains p , $\phi^{-1}(\phi(N)) = N$. We now list, for reference, some of the basic facts concerning the decomposition \mathcal{D} .

- (i) For each $x \in M$, $D(x)$ is connected (see [9], Theorem 3, p. 210 for metric continua, and [2], Theorem 1.2 for the general case).
- (ii) For each $x \in M$, $D(x)$ has void interior in px (see [7], Theorem 3-44).
- (iii) If M is smooth at p , then \mathcal{D} is a monotone upper semicontinuous decomposition and M/\mathcal{D} is a generalized tree which is smooth at $D(p)$ (see [4], Theorem 5.2 and Theorem 4.1).
- (iv) If M/\mathcal{D} is a continuum which is smooth at $D(p)$, then M is smooth at p (see [13], Theorem 3.1 for metric continua, and [11], Theorem 6.3 for the general case).

3. The Main Results

Throughout this section M will denote a continuum containing a fixed point p .

We shall prove

Theorem 1. Each subcontinuum of M which contains p is a

monotone retract of M (i.e., M satisfies (**)) if and only if

- (a) M is smooth at p , and
- (b) for each subcontinuum N of M containing p , N is accessible and the components of $M - N$ form a null family.

Furthermore, if (**) holds, then M/\mathfrak{D} is a tree.

We shall need several lemmas.

Lemma 1. Let M be hereditarily unicoherent at p , and let N and P be subcontinua of M such that $p \in N \subseteq P$. If $r: M \rightarrow N$ is a monotone retraction, then $r|_P$ is a monotone retraction.

Proof. It suffices to show that $r^{-1}(x) \cap P$ is connected for each $x \in N$. If not, there exist disjoint closed sets A and B such that $r^{-1}(x) \cap P = A \cup B$ and $x \in A$. But this contradicts hereditary unicoherence at p since $(r^{-1}(x) \cup N) \cap P = (A \cup N) \cup B$ and $(A \cup N) \cap B = \emptyset$.

Lemma 2. Let M be irreducible from p to q . If each subcontinuum of M which contains p is a monotone retract of M , then M is smooth at p .

Proof. According to the Lemma of [6], M is hereditarily unicoherent at p . Thus, by (iv) of Section 2, it suffices to show that M/\mathfrak{D} is a continuum which is smooth at $D(p)$. We begin by showing that $D(z)$ is closed for each z in M . First suppose that x and y belong to $\text{cl}(D(z)) - D(z)$. By the hypothesis and Lemma 1, there is a monotone retraction $r: pz \rightarrow px \cup py$. By irreducibility $pz = (px \cup py) \cup r^{-1}(r(z))$. Thus $\{x, y\} \subseteq r(D(z)) = r(z)$ and $x = y$. In particular, $\text{cl}(D(z)) - D(z) = \{x\}$. Since $D(z)$ is connected (by (i) of Section 2) and pz is irreducible, $pz = px \cup D(z)$. But this implies that $D(z)$ has nonvoid interior in pz , contradicting (ii) of Section 2. Thus $D(z)$ is closed. We now show that each element $D(z)$ of \mathfrak{D} distinct from $D(p)$ and $D(q)$ separates $D(p)$

from $D(q)$ in M/\mathcal{D} . Let $z \in M - (D(p) \cup D(q))$ and let $r:M \rightarrow pz$ be a monotone retraction. By irreducibility $M = pz \cup r^{-1}(r(q))$ and $r(q) \in D(z)$. Since $r(q)$ separates p from q in M , $D(z)$ separates $D(p)$ from $D(q)$ in M/\mathcal{D} . It follows that M/\mathcal{D} is an arc (e.g., [3], Theorem 2.1). Consequently M/\mathcal{D} is a continuum which is smooth at $D(p)$.

A quite different (and somewhat longer) proof of Lemma 2 can be obtained by applying the characterization of smoothness for irreducible continua given by J. J. Charatonik in [1].

Example 2 in Section 4 shows that the converse of Lemma 2 is false.

Lemma 3. Let M be hereditarily unicoherent at p and assume that M/\mathcal{D} is a tree. Let N be a subcontinuum of M containing p and let C be a component of $M - N$. Then

- (a) *C is open and continuumwise connected.*
- (b) *At most one point of N is accessible from any point of C .*
- (c) *If $r:M \rightarrow N$ is a monotone retraction, then $r(C)$ is degenerate.*

Proof. Note that M is smooth at p by (iv) of Section 2.

- (a) Using the facts that $\phi:M \rightarrow M/\mathcal{D}$ is monotone and $\phi^{-1}(\phi(N)) = N$ (see (iii) of Section 2), it is easy to verify that $\phi^{-1}(\phi(C)) = C$. It follows that $\phi(C)$ is a component of $M/\mathcal{D} - \phi(N)$. As a component of an open subset of a tree, $\phi(C)$ is open and arcwise connected. Thus $C = \phi^{-1}(\phi(C))$ is open and continuumwise connected.
- (b) Suppose that x and y are distinct points of N which are accessible from points in C . Then there exist subcontinua X and Y of M such that $X \cap C \neq \emptyset \neq Y \cap C$, $X \cap N = \{x\}$, and $Y \cap N = \{y\}$. Applying (a), there exists a subcontinuum $K \subseteq C$ such that $X \cap K \neq \emptyset \neq Y \cap K$.

But then $(N \cup X \cup K) \cap (N \cup Y \cup K) = N \cup (K \cup (X \cap Y))$
 which is a separation, contradicting hereditary uni-
 coherence at p .

(c) If $x, y \in r(C)$, then $r^{-1}(x) \cap C \neq \emptyset \neq r^{-1}(y) \cap C$. Thus
 $x = y$ by (b).

We shall need the notion of aposyndesis due to F. B. Jones (see [8] for a discussion of the history of this concept). A continuum M is said to be *aposyndetic at x with respect to y* if there exists a subcontinuum K of M such that $x \in \text{int}(K) \subseteq K \subseteq M - \{y\}$. If for each pair of distinct points x and y of M , M is aposyndetic at x with respect to y (either one of the points with respect to the other), then M is said to be *aposyndetic (semi-aposyndetic)*.

In the next lemma we shall use the facts that every generalized tree is semi-aposyndetic, and that every aposyndetic generalized tree is a tree ([5], Theorem 3.5 and Corollary 2.1).

Lemma 4. *If M is smooth at p and for each subcontinuum N of M containing p the components of $M - N$ form a null family, then M/\mathcal{D} is a tree.*

Proof. Applying the hypothesis and the properties of ϕ discussed in Section 2, it is easy to verify that for each subcontinuum K of M/\mathcal{D} which contains $D(p)$, the components of $M/\mathcal{D} - K$ form a null family. Thus it suffices to assume that M is a generalized tree (i.e., $M = M/\mathcal{D}$), and prove that M is aposyndetic. Let x and y be distinct points of M . Since M is semi-aposyndetic, we can assume that there is a subcontinuum H of M such that $y \in \text{int}(H) \subseteq H \subseteq M - \{x\}$. If $x \leq y$, then M is aposyndetic at x with respect to y ([5], Corollary 3.6), and the proof is complete. Otherwise, $x \notin py \cup H$. Let \mathcal{C} denote the components of $M - (py \cup H)$, and let C denote the member of \mathcal{C} containing x . If $x \notin \text{int}(C)$, then there is a net x_n in

$M - (p \cup H \cup C)$ such that $\lim x_n = x$. Let C_n be the corresponding net in \mathcal{C} (i.e., $x_n \in C_n$), and assume without loss of generality that C_n converges. Since $x_n \notin C$ for each n , C_n is not eventually constant. But $x \in \lim C_n$ and $(\lim C_n) \cap (p \cup H) \neq \emptyset$, which contradicts the assumption that \mathcal{C} is a null family. Consequently, $x \in \text{int}(C) \subseteq \text{cl}(C) \subseteq M - \{y\}$, and M is aposyndetic.

Lemma 5. Let M be smooth at p and assume that for each subcontinuum N of M containing p , N is accessible and the components of $M - N$ form a null family. If N is a subcontinuum of M containing p and C is a component of $M - N$, then $N \cap \text{cl}(C)$ is degenerate.

Proof. Suppose that N and C are as in the hypothesis and that $N \cap \text{cl}(C)$ is nondegenerate. According to Lemma 4, M/\mathcal{D} is a tree and thus Lemma 3 applies. Consequently C is open and $M - C$ is a subcontinuum of M containing p . By hypothesis and Lemma 3(b), there is a unique point $x \in M - C$ which is accessible from each point of C . Observe that $x \in N \cap \text{cl}(C)$. Let H be a nondegenerate subcontinuum of M such that $N \cap H = \{x\}$ and $H - \{x\} \subseteq C$. Let $y \in N \cap \text{cl}(C)$ such that $y \neq x$, and let y_n be a net in $C - H$ converging to y . Arguing as above, we conclude that for each n there is a unique point $z_n \in N \cup H$ which is accessible from y_n . Since C is continuumwise connected (Lemma 3(a)) and M is hereditarily unicoherent at p it follows easily that $z_n \in H - N$ for each n . Let C_n denote the component of $M - (H \cup N)$ which contains y_n . Passing to a subnet if necessary, assume that C_n converges to a continuum C_0 . Since $y \in C_0$ and $C_0 \cap H \neq \emptyset$, the net C_n must be eventually constant. It follows that z_n is eventually constant, say $z_n = z_0$ for sufficiently large n . Thus $z_0 \in p y_n$ for sufficiently large n ; and by smoothness $z_0 \in p y \subseteq N$. Thus $z_0 \in N$ and $z_0 \in H - N$ which is a contradiction.

Proof of Theorem 1. (Only if) By the Lemma in [6], M is hereditarily unicoherent at p ; and by Lemma 1 and 2, each irreducible subcontinuum of the form px is smooth at p . To show that M is smooth at p it suffices to prove that \leq is closed in $M \times M$ ([5], Theorem 3.1). Let (x_n, y_n) be a net in \leq converging to (x, y) . Let $r: M \rightarrow px \cup py$ be a monotone retraction. By [4], Theorem 4.1, r preserves order, and hence $r(x_n) \leq r(y_n)$ for each n . Since px and py are smooth at p , so is $px \cup py$; and consequently $x = r(x) \leq r(y) = y$. Thus (x, y) belongs to \leq and M is smooth at p .

Let N be any subcontinuum of M containing p , and let $r: M \rightarrow N$ be a monotone retraction. If $x \in M - N$, then $r^{-1}(r(x)) \cap N = \{r(x)\}$, so N is accessible.

We next show that M/\mathcal{D} is a tree. By [12], Theorem 3, it suffices to show that each arc of the form $D(p)D(x)$ in M/\mathcal{D} is a monotone retract of M/\mathcal{D} . Let $r: M \rightarrow px$ be a monotone retraction. Since r preserves order, it is easy to verify that the induced map $r^*: M/\mathcal{D} \rightarrow D(p)D(x)$ defined by $r^*(D(y)) = D(r(y))$ for each $y \in M$ is a monotone retraction.

Finally, let N be a subcontinuum of M containing p and let \mathcal{C} denote the components of $M - N$. Assume that C_n is a net of elements of \mathcal{C} which is not eventually constant and converges to a subcontinuum C . Since each C_n is open by Lemma 3, it follows that $C \subseteq N$. Let $r: M \rightarrow N$ be a monotone retraction. Then, by Lemma 3, $r(C_n)$ is degenerate for each n . Hence $C = r(C) = \lim r(C_n)$ is degenerate; i.e., \mathcal{C} forms a null family.

(If) Let N be a subcontinuum of M which contains p . We must define a monotone retraction $r: M \rightarrow N$. For each $x \in M - N$, denote by $C(x)$ the component of $M - N$ containing x . Define $r: M \rightarrow N$ to be the unique retraction such that for each $x \in M - N$, $\{r(x)\} = N \cap \text{cl}(C(x))$. Note that r is a well-defined function by Lemma 5. Since point inverses of r are clearly connected, it

remains only to show that r is continuous. If not, there exists an open set U in the relative topology on N such that $r^{-1}(U)$ is not open in M . Let $z \in r^{-1}(U) - \text{int}(r^{-1}(U))$. Applying Lemma 4 and Lemma 3, it follows that $C(x)$ is open for each x , and thus $z \in U \subseteq N$. Consequently, there is a net z_n in $M - N$ such that $z_n \notin r^{-1}(U)$ for each n , $\lim z_n = z$, and $(N \cap \text{cl}(C(z_n))) \cap U = \emptyset$ for each n . Without loss of generality, assume that the net $C(z_n)$ converges to a continuum C . The net $C(z_n)$ is not eventually constant; for otherwise $z_n \in r^{-1}(z) \subseteq r^{-1}(U)$ for sufficiently large n . But C contains z and meets $N - U$, contradicting the assumption that the components of $M - N$ form a null family. Thus r is continuous.

Corollary 1. Let M be a generalized tree which is smooth at p . Then M is a tree if and only if for each subcontinuum N of M containing p , the components of $M - N$ form a null family.

Proof. (Only if) If M is a tree then each subcontinuum of M is a monotone retract of M ([10], Theorem 2.1), and Theorem 1 applies.

(If) By Lemma 4, $M/\mathcal{D} = M$ is a tree.

Corollary 2. Let M be a continuum which is irreducible about a finite set. Each subcontinuum of M which contains p is a monotone retract of M (i.e., M satisfies (**)) if and only if

- (a) M is smooth at p , and
- (b) each subcontinuum of M containing p is accessible.

Furthermore, if (**) holds, then M/\mathcal{D} is a finite tree.

Proof. If N is a subcontinuum of M which contains p , then the components of $M - N$ form a finite, hence null, family. Now apply Theorem 1.

4. Examples

Corollary 2 shows that the "null family" condition in

Theorem 1 is superfluous for continua irreducible about finitely many points. The following example shows that this condition cannot be omitted in general, even if M/\mathcal{D} is known to be a tree.

Example 1. Let M be the plane continuum defined by:

$$\begin{aligned} M = \{ & (x,y) : y = 1 + \sin 1/x \text{ for } -1 \leq x < 0 \} \\ & \cup \{ (x,y) : x = 0 \text{ and } 0 \leq y \leq 2 \} \\ & \cup \left(\bigcup_{n=0}^{\infty} \{ (x,y) : y = nx \text{ for } 0 \leq y \leq 2 \} \right). \end{aligned}$$

Note that M is the union of a "simple harmonic fan" and a "sin $1/x$ curve." The "sin $1/x$ curve" is not a monotone retract of M ; and M/\mathcal{D} is a locally connected fan (i.e., a dendrite with only one ramification point).

The next example shows that the "accessibility" condition in Theorem 1 cannot be omitted even for irreducible continua.

Example 2. Let M be the plane continuum defined by:

$$\begin{aligned} M = \{ & (x,y) : y = \sin 1/x \text{ for } -1 \leq x < 0 \text{ and } 0 < x \leq 1 \} \\ & \cup \{ (x,y) : x = 0 \text{ and } -1 \leq y \leq 1 \}. \end{aligned}$$

Note that M is the union of two "sin $1/x$ curves" with a common limit segment. Neither of the "sin $1/x$ curves" is a monotone retract of M . Thus M does not satisfy (**).

5. Concluding Remarks

Consider the following weak version of condition (**).

(***) Each subcontinuum of M which is irreducible between a fixed point p and some other point is a monotone retract of M .

If M is a dendroid, then (***) is equivalent to (**) by [12], Theorem 3.

Question. Are conditions (**) and (***) equivalent for an arbitrary continuum M ?

We remark that it is possible to modify the proof of Theorem 1 to obtain an affirmative answer to this question in the special case when M is hereditarily unicoherent at the point p . Thus it suffices to determine whether a continuum M satisfying (***) must be hereditarily unicoherent at p .

References

1. J. J. Charatonik, *On irreducible smooth continua*, Proc. International Symposium in Topology, Budva, Yugoslavia 1972, 45-50.
2. G. R. Gordh, Jr., *Doctoral Dissertation*, University of California at Riverside, 1971.
3. _____, *Monotone decomposition of irreducible Hausdorff continua*, Pacific J. Math. 36 (1971), 647-658.
4. _____, *On decompositions of smooth continua*, Fund. Math. 75 (1972), 51-60.
5. _____, *Concerning closed quasi-orders on hereditarily unicoherent continua*, Fund. Math. 78 (1973), 61-73.
6. _____ and L. Lum, *Monotone retracts and some characterizations of dendrites*, Proc. Amer. Math. Soc. 59 (1976), 156-158.
7. J. G. Hocking and G. S. Young, *Topology*, Addison-Wesley, Reading, Mass., 1961.
8. F. B. Jones, *Aposyndesis revisited*, Proceedings of the University of Oklahoma Topology Conference, 1972, 64-78.
9. K. Kuratowski, *Topology II*, PWN--Academic Press, Warsaw--New York, 1968.
10. L. Lum, *A characterization of local connectivity in dendroids*, Studies in Topology, Academic Press, New York, 1975, 331-338.
11. _____, *Weakly smooth continua*, Trans. Amer. Math. Soc., 214 (1975), 153-167.
12. _____, *Order preserving and monotone retracts of a dendroid*, Proc. of the Auburn Topology Conference 1976.
13. T. Maćkowiak, *Some characterizations of smooth continua*, Fund. Math. 79 (1973), 173-186.
14. _____, *On smooth continua*, Fund. Math. 85 (1974), 79-95.

Salem College

Winston-Salem, North Carolina 27108