

TOPOLOGY PROCEEDINGS



Volume 5, 1980

Pages 209–213

<http://topology.auburn.edu/tp/>

Research Announcement:

U-EMBEDDED SUBSETS OF THE PLANE

by

RONNIE LEVY AND M. D. RICE

Topology Proceedings

Web: <http://topology.auburn.edu/tp/>

Mail: Topology Proceedings
Department of Mathematics & Statistics
Auburn University, Alabama 36849, USA

E-mail: topolog@auburn.edu

ISSN: 0146-4124

COPYRIGHT © by Topology Proceedings. All rights reserved.

U-EMBEDDED SUBSETS OF THE PLANE

Ronnie Levy and M. D. Rice

For a metric space M , let $U(M)$ denote the set of all real-valued uniformly continuous functions on M and let $U^*(M)$ denote the set of bounded members of $U(M)$. If X is a subset of M , then X is U -embedded [respectively U^* -embedded] in M if every element of $U(X)$ [respectively $U^*(X)$] is the restriction to X of an element of $U(M)$. The following theorem of Katětov answers all questions about U^* -embedding.

Theorem. (Katětov) If M is a metric space, every subset of M is U^ -embedded in M .*

The question of which subsets of a metric space are U -embedded is much more difficult. Not every subset of an arbitrary metric space is U -embedded--the set N of natural numbers is not U -embedded in the space \mathbf{R} of real numbers because the uniformly continuous function $f: N \rightarrow \mathbf{R}$ given by $f(n) = n^2$ does not extend to a uniformly continuous function on \mathbf{R} . This example provides the key to characterizing the U -embedded subsets of \mathbf{R} .

Theorem 1. X is U -embedded in \mathbf{R} if and only if X is not the union of an infinite uniformly discrete family of non-empty subsets.

Unlike the case of \mathbf{R} , the U -embedded subsets of \mathbf{R}^2 have not been characterized. The next theorem gives a

class of subsets which are not U-embedded in \mathbf{R}^2 , as well as a class of subsets which are U-embedded in \mathbf{R}^2 .

Theorem 2. Suppose M is a normed linear space and $X \subseteq M$.

a) If X is the union of an infinite uniformly discrete family of subsets, then X is not U-embedded in M .

b) If X is convex, then X is U-embedded in M .

Since the convex subsets of \mathbf{R}^2 are U-embedded, it is natural to ask which starlike regions of \mathbf{R}^2 are U-embedded. The answer to even this question is not known. The following examples indicate the types of problems which can arise. Let $(a_n)_{n=1}^{\infty}$ be an unbounded increasing sequence of numbers and let $X = \bigcup_{k=0}^{\infty} L_k$, where L_0 is the non-negative x-axis and for $k = 1, 2, \dots$, L_k is the segment joining the origin to $(a_k, 1)$. The sequences $a_n = n$ and $a_n = 2^n$ define the spaces X_1 and X_2 in Figure 1. Then X_1 is U-embedded in \mathbf{R}^2 , but X_2 is not U-embedded in \mathbf{R}^2 . The proof that X_1 is U-embedded combines the following facts: (i) L_0 is U-embedded in \mathbf{R}^2 . (Theorem 2b.) (ii) X_1 is U^* -embedded in \mathbf{R}^2 (Katětov's theorem). (iii) If $f \in U(X_1)$ and the restriction of f to L_0 is bounded, then $f \in U^*(X_1)$. The fact that X_2 is not U-embedded is established by showing that the function $f: X_2 \rightarrow \mathbf{R}$ defined by

$$f(x,y) = \begin{cases} 0 & \text{if } y \leq 1/2 \\ a_n & \text{if } (x,y) = (a_n, 1) \\ \text{linear between } (a_n/2, 1/2) \text{ and } (a_n, 1) & \end{cases}$$

is uniformly continuous. (See Figure 2.) By condition

(iii) above, f cannot be extended to an element of $U(X_1)$, so f certainly cannot be extended to an element of $U(\mathbb{R}^2)$. By generalizing these arguments, one can establish the following.

Theorem 3. If X is the star-like region defined above by the sequence (a_n) , then X is U -embedded in \mathbb{R}^2 if and only if $\lim_{n \rightarrow \infty} a_n/a_{n+1}$ (exists and) = 1.

References

- M. Katětov, *On real-valued functions in topological spaces*,
Fund. Math. 38 (1951), 85-91.
- R. Levy and M. D. Rice, *The approximation of uniformly
continuous mappings by Lipschitz and Hölder mappings*
(submitted manuscript).
- _____, *The extension of uniformly continuous mappings
defined on subsets of metric spaces* (submitted manu-
script).

George Mason University

Fairfax, Virginia 22030






FIGURE 1



Graph of $f: X_2 \rightarrow \mathbb{R}$

FIGURE 2