

---

# TOPOLOGY PROCEEDINGS



Volume 6, 1981

Pages 317–328

---

<http://topology.auburn.edu/tp/>

## EMBEDDING PIECEWISE LINEAR $\mathbf{R}^\infty$ -MANIFOLDS INTO $\mathbf{R}^\infty$

by

RICHARD E. HEISEY

---

### Topology Proceedings

**Web:** <http://topology.auburn.edu/tp/>

**Mail:** Topology Proceedings  
Department of Mathematics & Statistics  
Auburn University, Alabama 36849, USA

**E-mail:** [topolog@auburn.edu](mailto:topolog@auburn.edu)

**ISSN:** 0146-4124

COPYRIGHT © by Topology Proceedings. All rights reserved.

**EMBEDDING PIECEWISE LINEAR  
 $R^\infty$ -MANIFOLDS INTO  $R^\infty$**

**Richard E. Heisey**

It is well known that a compact piecewise linear manifold of dimension  $n$ ,  $n \geq 2$ , can be piecewise linearly embedded into  $R^{2n}$ . Here we establish an infinite-dimensional analogue. Let  $R^\infty = \varinjlim R^n$ , the countable direct limit of lines. We show that any separable, paracompact piecewise linear  $R^\infty$ -manifold can be piecewise linearly embedded onto a closed piecewise linear submanifold of  $R^\infty$ . As a consequence piecewise linear  $R^\infty$ -manifolds may be regarded as "polyhedra" in  $R^\infty$ .

**I. Definitions and Statement of the Main Theorem**

Let  $R^\infty = \varinjlim R^n$ , the countable direct limit of lines. We think of  $R^\infty$  as  $\{(x_i) : \text{all but finitely many } x_i \text{ are } 0\}$  and identify  $R^n$  with  $R^n \times \{0\} \times \{0\} \times \{0\} \times \dots \subset R^\infty$ . A straightforward observation, e.g. see Lemma III-6 of [1], shows that any compact subset of  $R^\infty$  is contained in some  $R^n$ . Let  $U$  and  $V$  be open subsets of  $R^\infty$ . A map  $f: U \rightarrow V$  is  $R^\infty$ -piecewise linear, hereafter  $R^\infty$ -p.l., if for every compact polyhedron  $C \subset U$  and for every choice of  $n$  such that  $f(C) \subset V \cap R^n$ , the restriction  $f|_C: C \rightarrow V \cap R^n$  is piecewise linear in the usual sense. (By *polyhedron* we mean a subset  $P \subset R^n$  such that every point  $x \in P$  has a cone neighborhood  $xL$ , where  $L$  is compact. For this and other basic definitions and results from piecewise linear topology see [3].)

A *piecewise linear  $R^\infty$ -atlas* for a space  $M$  is a collection of pairs  $\{(U_\alpha, \phi_\alpha)\}$  where  $\{U_\alpha\}$  is an open cover of  $M$  by nonempty sets,  $\phi_\alpha: U_\alpha \rightarrow \phi_\alpha(U_\alpha)$  is a homeomorphism onto an open subset of  $R^\infty$ , and where, if  $U_\alpha \cap U_\beta \neq \emptyset$ ,  $\phi_\beta \phi_\alpha^{-1}: \phi_\alpha(U_\alpha \cap U_\beta) \rightarrow \phi_\beta(U_\alpha \cap U_\beta)$  is  $R^\infty$ -p.l. A *piecewise linear  $R^\infty$ -structure* for  $M$  is a maximal p.l.  $R^\infty$ -atlas for  $M$ . Since any p.l.  $R^\infty$ -atlas for the space  $M$  is contained in a unique maximal p.l.  $R^\infty$ -atlas, a p.l.  $R^\infty$ -atlas for  $M$  determines a p.l.  $R^\infty$ -structure for  $M$ . A *piecewise linear  $R^\infty$ -manifold* is a paracompact space  $M$  together with a p.l.  $R^\infty$ -structure. A *piecewise linear  $R^\infty$ -atlas for the p.l.  $R^\infty$ -manifold  $M$*  is any p.l.  $R^\infty$ -atlas for the space  $M$  which is contained in the p.l.  $R^\infty$ -structure for  $M$ . An element  $(U, \phi)$  of some p.l.  $R^\infty$ -atlas for the p.l.  $R^\infty$ -manifold  $M$  is a *piecewise linear  $R^\infty$ -chart* for  $M$ . If  $(U, \phi)$  is such a chart and if  $\phi': U' \rightarrow \phi'(U')$  is the restriction of  $\phi$  to a nonempty open subset of  $U$  then, clearly,  $(U', \phi')$  is such a chart.

A map  $f: M \rightarrow N$  between two p.l.  $R^\infty$ -manifolds is  *$R^\infty$ -piecewise linear* if for each  $x \in M$  there is a p.l.  $R^\infty$ -chart  $(U, \phi)$  for  $M$  and a p.l.  $R^\infty$ -chart  $(V, \psi)$  for  $N$  such that  $x \in U$ ,  $f(x) \in V$  and  $\psi f \phi^{-1}: \phi(U \cap f^{-1}(V)) \rightarrow \psi(V)$  is  $R^\infty$ -p.l. It follows then that if  $f: M \rightarrow N$  is  $R^\infty$ -p.l. and  $(U, \phi)$  and  $(V, \psi)$  are any given  $R^\infty$ -p.l. charts with  $x \in U$  and  $f(x) \in V$  then  $\psi f \phi^{-1}: \phi(U \cap f^{-1}(V)) \rightarrow \psi(V)$  is  $R^\infty$ -p.l. An  $R^\infty$ -p.l. map  $f: M \rightarrow N$  is an  $R^\infty$ -p.l. *isomorphism* if  $f$  is a homeomorphism and  $f^{-1}: N \rightarrow M$  is  $R^\infty$ -p.l.

Let  $\tau: R^\infty \times R^\infty \rightarrow R^\infty$  be the natural linear homeomorphism  $\tau((x_i), (y_i)) = (x_1, y_1, x_2, y_2, x_3, y_3, \dots)$ . (That  $\tau$  is a

homeomorphism follows since  $R$  is locally compact [1, Corollary III-1].) We identify  $R^\infty \times R^\infty$  with  $R^\infty$  as p.l.  $R^\infty$ -manifolds via  $\tau$ . Thus, for a p.l.  $R^\infty$ -manifold  $M$  we may identify any given p.l.  $R^\infty$ -chart with image in  $R^\infty$  with one whose image is in  $R^\infty \times R^\infty$ . Let  $N$  be a subset of the p.l.  $R^\infty$ -manifold  $M$  such that for each  $x \in N$  there is a p.l.  $R^\infty$ -chart  $(U, \phi)$  for  $M$  with  $x \in U$  such that  $\phi(U) = U_1 \times U_2$ ,  $U_1$  open in  $R^\infty$ , and such that  $\phi(U \cap N) = U_1 \times \{0\}$ . (Here  $0 = (0, 0, 0, \dots)$ .) If we identify  $R^\infty \times \{0\}$  with  $R^\infty$ , then, for such a chart  $(U, \phi)$ ,  $\phi|_{(U \cap N)}: U \cap N \rightarrow U_1$  is a homeomorphism. Thus, charts of the form  $(U', \phi') = (U \cap N, \phi|_{(U \cap N)})$  form a p.l.  $R^\infty$ -atlas for  $N$  inducing a p.l.  $R^\infty$ -structure for  $N$ . With this p.l.  $R^\infty$ -structure we call  $N$  a p.l.  $R^\infty$ -submanifold of  $M$  (of infinite codimension).

We may now state our main theorem.

*Theorem.* If  $M$  is a separable, paracompact p.l.  $R^\infty$ -manifold then there is an  $R^\infty$ -p.l. isomorphism  $f: M \rightarrow N$ ,  $N$  a closed p.l.  $R^\infty$ -submanifold of  $R^\infty$ .

The proof of this theorem is given in §III.

There is a natural definition for  $R^\infty$ -polyhedra and p.l. maps between them.

*Definition.* A subset  $X$  of  $R^\infty$  is an  $R^\infty$ -polyhedron if for each compact polyhedron  $C$  in  $R^\infty$ ,  $C \cap X$  is a polyhedron. A map  $f: X \rightarrow Y$  between two  $R^\infty$ -polyhedra is  $R^\infty$ -piecewise linear if for each compact polyhedron  $C \subset X$  and any choice of  $n$  such that  $f(C) \subset Y \cap R^n$ ,  $f|_C: C \rightarrow Y \cap R^n$  is p.l.

We conclude our paper by showing, in §IV, that any p.l.  $\mathbb{R}^\infty$ -submanifold of  $\mathbb{R}^\infty$  is an  $\mathbb{R}^\infty$ -polyhedron and that for maps between two such submanifolds the two definitions of  $\mathbb{R}^\infty$ -piecewise linear agree. Thus, the study of p.l.  $\mathbb{R}^\infty$ -manifolds and  $\mathbb{R}^\infty$ -p.l. maps between them is a special case of the study of  $\mathbb{R}^\infty$ -polyhedra and  $\mathbb{R}^\infty$ -p.l. maps between them.

## II. Preliminary Results

Lemma 1 below is the crucial auxiliary result we will need. In addition we establish some helpful elementary results about  $\mathbb{R}^\infty$ -p.l. maps. First, a useful definition.

*Definition.* If  $U$  is an open subset of  $\mathbb{R}^\infty$  and  $P$  is a finite-dimensional polyhedron we say a map  $f: U \rightarrow P$  is *piecewise linear* (p.l.) if for every compact polyhedron  $C \subset U$ ,  $f|_C: C \rightarrow P$  is p.l. If  $M$  is a p.l.  $\mathbb{R}^\infty$ -manifold we say that  $f: M \rightarrow P$  is p.l. if  $f\phi^{-1}: \phi(U) \rightarrow P$  is p.l. for each p.l.  $\mathbb{R}^\infty$ -chart  $(U, \phi)$  for  $M$ .

*Lemma 1.* Let  $U$  be an open subset of  $\mathbb{R}^\infty$ . Let  $A \subset W \subset U$  where  $A$  is closed in  $U$  and  $W$  is open in  $U$ . Then there is a p.l. map  $\lambda: U \rightarrow I$  such that  $\lambda|_A = 0$  and  $\lambda|_{(U-W)} = 1$ .

*Proof.* The proof proceeds in the spirit of the proof of Proposition IV.2 in [2]. Let  $c = \{c_i\} = (c_1, c_2, \dots, c_{n_0}, 0, 0, \dots) \in \mathbb{R}^\infty$ , and let  $V = [(c_1 - \epsilon_1, c_1 + \epsilon_1) \times (c_2 - \epsilon_2, c_2 + \epsilon_2) \times \dots \times (c_{n_0} - \epsilon_{n_0}, c_{n_0} + \epsilon_{n_0}) \times (-\epsilon_{n_0+1}, \epsilon_{n_0+1}) \times \dots] \cap \mathbb{R}^\infty$ , where  $\epsilon_i > 0$ . Define  $V(2) = [(c_1 - 2\epsilon_1, c_1 + 2\epsilon_1) \times (c_2 - 2\epsilon_2, c_2 + 2\epsilon_2) \times \dots] \cap \mathbb{R}^\infty$ . Let  $\alpha_i: \mathbb{R} \rightarrow I$  be a p.l. map such

that  $\alpha_i | [c_i - \epsilon_i, c_i + \epsilon_i] = 1$  and  $\alpha_i | [R \setminus (c_i - 2\epsilon_i, c_i + 2\epsilon_i)] = 0$ . Define  $\psi_i: R^\infty \rightarrow I$  by  $\psi_i((x_1, x_2, \dots)) = \alpha_i(x_i)$  and then  $\psi(x) = \min\{\psi_1(x) | i = 1, 2, 3, \dots\}$ . If  $x \in R^n$ ,  $n \geq n_0$ , then  $\psi_i(x) = 1$ ,  $i > n$ , and  $\psi(x) = \min\{\psi_1(x), \dots, \psi_n(x)\}$ . Thus,  $\psi$  is continuous,  $\psi|_{R^n}$  is p.l.,  $n \geq 1$ ,  $\psi|_V = 1$ , and  $\psi|_{[R \setminus V(2)]} = 0$ . Note that sets of the form of  $V$  form a basis for  $R^\infty$  [1, Proposition II.1(a)].

Let  $A$ ,  $W$ , and  $U$  be as in Lemma 1. By elementary reasoning  $U = \varinjlim C_n$  where  $C_n \subset R^n$  is a compact polyhedron and  $C_n \subset \text{Int}_{R^{n+1}} C_{n+1}$ . Let  $A_k = A \cap C_k$ . Choose finitely many basic open sets of the type in the preceding paragraph,  $V_{1,1}, \dots, V_{1,k_1}$ , covering compact  $A_1$  and such that  $V_{1,i}(2) \subset W$ ,  $i = 1, 2, \dots, k_1$ . For each  $x \in A_2 \setminus C_1$ , choose a basic open set  $V_{2,x}$  such that  $x \in V_{2,x} \subset V_{2,x}(2) \subset W \setminus C_1$ . Then  $V_{1,1}, \dots, V_{1,k_1}$  together with  $\{V_{2,x} : x \in A_2 \setminus C_1\}$  form an open cover of  $A_2$ , so we may select a finite subcover  $V_{1,1}, \dots, V_{1,k_1}, V_{2,1}, \dots, V_{2,k_2}$ . Continuing, we obtain a sequence  $V_{1,1}, \dots, V_{1,k_1}, V_{2,1}, \dots, V_{2,k_2}, V_{3,1}, \dots, V_{3,k_3}, \dots$  covering  $A$  such that  $V_{i,j}(2) \subset W \setminus C_{i-1}$ ,  $i > 1$ . By the work in the preceding paragraph, for each  $(i,j)$  there is a p.l. map  $\phi_{i,j}: U \rightarrow I$  such that  $\phi_{i,j}|_{V_{i,j}} = 0$  and  $\phi_{i,j}|_{[U \setminus V_{i,j}(2)]} = 1$ . Let  $\phi_i = \min\{\phi_{i,1}, \dots, \phi_{i,k_i}\}$  and  $\phi = \min\{\phi_i | i = 1, 2, \dots\}$ . Let  $x \in C_n$ . Then  $x \notin V_{k,j}(2)$ ,  $k > n$ , so that  $\phi_{k,j}(x) = 1$ ,  $k > n$ . Therefore,  $\phi|_{C_n} = \min\{\phi_1, \dots, \phi_n\}$ . Thus,  $\phi|_{C_n}$  is p.l.,  $n \geq 1$ , and it follows that  $\phi$  is p.l. Also,  $\phi|_A = 0$  and  $\phi|(U \setminus W) = 1$ .

Lemma 2. (a) The composition of  $R^\infty$ -p.l. maps is  $R^\infty$ -p.l. Also, if  $f: M \rightarrow N$  is  $R^\infty$ -p.l. and  $g: N \rightarrow P$ ,  $P$  a

finite-dimensional polyhedron, is p.l., then  $gf$  is p.l.

(b) A map  $f = (f_1, f_2, f_3, \dots): M \rightarrow \mathbb{R}^\infty$ ,  $M$  a p.l.  $\mathbb{R}^\infty$ -manifold, is  $\mathbb{R}^\infty$ -p.l. if and only if each  $f_i$  is p.l.

*Proof.* The proof of (a) is straightforward, and we omit it. For (b) regard a p.l. map (on a finite-dimensional polyhedron) as one that is locally conical [3, p. 5]. Given a p.l.  $\mathbb{R}^\infty$ -chart  $(U, \phi)$  for  $M$  and  $\bar{a}$  compact polyhedron  $C \subset \phi(U)$ ,  $f\phi^{-1}(c) \subset \mathbb{R}^n$  some  $n$ . Thus, it is clear that if each  $f_i\phi^{-1}$  is locally conical so is  $f\phi^{-1}|_C$ . Thus,  $f\phi^{-1}$  is  $\mathbb{R}^\infty$ -p.l. Conversely, if  $f$  is  $\mathbb{R}^\infty$ -p.l. then  $f_i = \pi_i f$  where  $\pi_i: \mathbb{R}^\infty \rightarrow \mathbb{R}$  is the projection onto the  $i$ th-coordinate. Since  $\pi_i$  is p.l.,  $f_i$  is p.l. by (a).

*Lemma 3.* If  $\{(U_\alpha, \phi_\alpha) \mid \alpha \in A\}$  is a p.l.  $\mathbb{R}^\infty$ -atlas for the p.l.  $\mathbb{R}^\infty$ -manifold  $M$ , then there is a p.l.  $\mathbb{R}^\infty$ -atlas  $\{(U_\alpha, \psi_\alpha) \mid \alpha \in A\}$  for  $M$  such that  $\psi_\alpha(x) \in (-1, 1)^\infty = \varinjlim (-1, 1)^n$ , all  $\alpha \in A$ ,  $x \in U_\alpha$ .

*Proof.* Let  $\beta: \mathbb{R} \rightarrow (-1, 1)$  be a p.l. homeomorphism taking 0 to 0. Then  $\beta': \mathbb{R}^\infty \rightarrow (-1, 1)^\infty$ , defined by  $\beta'(x_1, x_2, x_3, \dots) = (\beta(x_1), \beta(x_2), \beta(x_3), \dots)$ , is an  $\mathbb{R}^\infty$ -p.l. isomorphism. Letting  $\psi_\alpha = \beta'\phi_\alpha$  gives the desired atlas.

*Lemma 4.* Let  $(U, \phi)$  be a p.l.  $\mathbb{R}^\infty$ -chart for the p.l.  $\mathbb{R}^\infty$ -manifold  $M$  such that  $\phi(U) \subset (-1, 1)^\infty$ . Let  $A$  be a closed subset of  $U$ ,  $V$  an open subset of  $U$  such that  $A \subset V \subset \bar{V} \subset U$ ,  $\bar{V}$  the closure of  $V$  in  $M$ . Then there is a p.l. map  $\lambda: M \rightarrow I$  and an  $\mathbb{R}^\infty$ -p.l. map  $\psi: M \rightarrow (-1, 1)^\infty \subset \mathbb{R}^\infty$  such that  $\lambda|_A = 1$ ,  $\lambda|_{M \setminus V} = 0$ ,  $\psi|_{\lambda^{-1}(1)} = \phi|_{\lambda^{-1}(1)}$ , and  $\psi|_{(U \setminus V)} = \mathbf{0}$ .

*Proof.* By Lemma 1 there is a p.l. map  $\lambda': \phi(U) \rightarrow I$  such that  $\lambda'|_{\phi(A)} = 1$ ,  $\lambda'|_{\phi(U \setminus \bar{V})} = 0$ . Define p.l.  $\lambda: M \rightarrow I$

by  $\lambda|U = \lambda'\phi$  and  $\lambda|(M \setminus \bar{V}) = 0$ . Write  $\phi(x) = (\phi_1(x), \phi_2(x), \dots)$ .

Define  $\psi = (\psi_1, \psi_2, \dots)$  by

$$\psi_i(x) = \begin{cases} \max\{-\lambda(x), \min\{\lambda(x), \phi_i(x)\}\}, & x \in U \\ 0, & x \in M \setminus \bar{V}. \end{cases}$$

That  $\psi$  is  $\mathbb{R}^\infty$ -p.l. follows from Lemma 2(b). The other desired properties are clear.

Finally, we will use the following.

*Proposition 5.* Let  $M$  and  $N$  be p.l.  $\mathbb{R}^\infty$ -manifolds. Let  $f: M \rightarrow N$  be an  $\mathbb{R}^\infty$ -p.l. map which is also a homeomorphism. Then  $f$  is an  $\mathbb{R}^\infty$ -p.l. isomorphism. I.e.  $f^{-1}$  is also  $\mathbb{R}^\infty$ -p.l.

*Proof.* Let  $(U, \phi)$  be a p.l.  $\mathbb{R}^\infty$ -chart at  $x \in M$ ,  $(V, \psi)$  a p.l.  $\mathbb{R}^\infty$ -chart at  $f(x)$  in  $N$  such that  $f^{-1}(V) \subset U$ . It suffices to show that  $\phi f^{-1} \psi^{-1}: \psi(V) \rightarrow \phi(f^{-1}(V))$  is  $\mathbb{R}^\infty$ -p.l. Let  $C \subset \psi(V)$  be a compact polyhedron. Then  $\phi f^{-1} \psi^{-1}(C)$  is compact. Hence, we may choose  $n$  and then a compact polyhedron  $P$  such that  $\phi f^{-1} \psi^{-1}(C) \subset P \subset \phi(f^{-1}(V)) \cap \mathbb{R}^n$ . On  $P$ ,  $\psi f \phi^{-1}$  is a p.l. homeomorphism, so  $Q = \psi f \phi^{-1}(P)$  is a compact polyhedron [3, p. 13] and  $\phi f^{-1} \psi^{-1}|Q$  is p.l. [3, p. 6]. Since  $C$  is a subpolyhedron of  $Q$ ,  $\phi f^{-1} \psi^{-1}|C$  is also p.l., as required.

In relation to the above proposition we remark that if  $f: M \rightarrow N$  is an  $\mathbb{R}^\infty$ -p.l. map such that  $f(M) \subset Q$  where  $Q$  is a p.l.  $\mathbb{R}^\infty$ -submanifold of  $N$  then  $f: M \rightarrow Q$  is also (clearly)  $\mathbb{R}^\infty$ -p.l. Thus, if  $f$  is also a topological embedding onto  $Q$ , then  $f: M \rightarrow Q$  is a p.l.  $\mathbb{R}^\infty$ -isomorphism.

### III. Proof of the Theorem

Let  $M$  be as in the theorem. Let  $\rho: \mathbb{R}^\infty \rightarrow (\mathbb{R}^\infty)^\infty = \varinjlim (\mathbb{R}^\infty)^n$  be the map obtained by Cantor diagonalization.



That is,  $\rho((x_1, x_2, x_3, \dots)) = ((x_1, x_3, x_6, x_{10}, \dots), (x_2, x_5, x_9, \dots), (x_4, x_8, \dots), \dots)$ . Then  $\rho$  is a linear homeomorphism [1, Corollary III-3]. Identify  $\mathbb{R}^\infty$  with  $(\mathbb{R}^\infty)^\infty$  as p.l.  $\mathbb{R}^\infty$ -manifolds via  $\rho$ . It suffices, then, to show that there is an  $\mathbb{R}^\infty$ -p.l. isomorphism  $f: M \rightarrow (\mathbb{R}^\infty)^\infty$  onto a closed  $\mathbb{R}^\infty$ -submanifold of  $(\mathbb{R}^\infty)^\infty$ . Note that, by Lemma 2(b), with our identification a map  $f: M \rightarrow (\mathbb{R}^\infty)^\infty$  is  $\mathbb{R}^\infty$ -p.l. if and only if each of its projections to  $\mathbb{R}$  is p.l. and, hence, if and only if each of its projections to  $\mathbb{R}^\infty$  is  $\mathbb{R}^\infty$ -p.l.

Let  $m \in M$ . By Lemma 3 there is a p.l.  $\mathbb{R}^\infty$ -chart  $(U_m, \phi_m)$  with  $m \in U_m$  and  $\phi_m(U_m) \subset (-1, 1)^\infty$ . If we choose an open set  $G$  such that  $\phi(m) \in G \subset \bar{G} \subset \phi(U_m)$ ,  $\bar{G}$  the closure of  $G$  in  $\mathbb{R}^\infty$ , then for any  $U$  with  $\bar{U} \subset \phi^{-1}(G)$  we have  $\phi(\bar{U}) = \overline{\phi(U)}$ . Thus, there is a p.l.  $\mathbb{R}^\infty$ -atlas  $\{(U_\alpha, \phi_\alpha)\}$  for  $M$  such that each  $\phi_\alpha(U_\alpha) \subset (-1, 1)^\infty$  and each  $\phi_\alpha$  extends to a closed embedding  $\phi_\alpha: \bar{U}_\alpha \rightarrow \overline{\phi_\alpha(U_\alpha)}$  into  $\mathbb{R}^\infty$ . Since  $M$  is paracompact and Lindelöf we thus obtain a countable, locally finite, p.l.  $\mathbb{R}^\infty$ -atlas  $\{(U_i, \phi_i)\}$  for  $M$  such that, for each  $i$ ,  $\phi_i(U_i) \subset (-1, 1)^\infty$  and  $\phi_i$  extends to a closed embedding into  $\mathbb{R}^\infty$ .

Let  $\{W_i\}, \{V_i\}$  be precise open refinements of  $U_i$  such that

$$\phi \neq W_i \subset \bar{W}_i \subset V_i \subset \bar{V}_i \subset U_i$$

(the closures in  $M$ ). By Lemma 4 there is, for each  $i$ , a p.l. map  $\lambda_i: M \rightarrow I$  and an  $\mathbb{R}^\infty$ -p.l. map  $\psi_i: M \rightarrow \mathbb{R}^\infty$  such that  $\lambda_i|_{\bar{W}_i} = 1$ ,  $\lambda_i|(M - V_i) = 0$ ,  $\psi_i|_{\lambda_i^{-1}(1)} = \phi_i$  and  $\lambda_i|(M - V_i) = 0$ .

Choose a nonzero point  $e \in \mathbb{R}^\infty$ . Define  $f: M \rightarrow (\mathbb{R}^\infty)^\infty$  by  $f(m) = (\sum_{i=1}^\infty i \lambda_i(m) e, \psi_1(m), \lambda_1(m) e, \psi_2(m), \lambda_2(m) e, \psi_3(m), \lambda_3(m) e, \dots)$ .

We will show that  $f$  is the desired  $\mathbb{R}^\infty$ -p.l. isomorphism. The local finiteness of  $\{U_i\}$  guarantees that the sum is finite and that  $f$  is well defined, i.e. that  $f(m) \in (\mathbb{R}^\infty)^\infty$ . Let  $f_i$  be the projection of  $f$  onto the  $i$ -th copy of  $\mathbb{R}^\infty$ . Since each  $f_i$  is  $\mathbb{R}^\infty$ -p.l.,  $f$  is an  $\mathbb{R}^\infty$ -p.l. map. Let  $x$  and  $y$  be distinct elements of  $M$ . Choose  $i$  such that  $x \in W_i$ . If  $\lambda_i(x) \neq \lambda_i(y)$ , then clearly  $f(x) \neq f(y)$ . Otherwise  $\lambda_i(y) = \lambda_i(x) = 1$  which implies  $\psi_i(x) = \phi_i(x) \neq \phi_i(y) = \psi_i(y)$ . Thus,  $f$  is one-to-one.

To see that  $f$  is a closed topological embedding let  $f(m_\alpha) \rightarrow y = (y_1, y_2, \dots) \in (\mathbb{R}^\infty)^\infty$ ,  $\{m_\alpha | \alpha \in A\}$  a net in  $A$ , a closed subset of  $M$ . Then  $f_1(m_\alpha) \rightarrow y_1$  implies that for some  $n$  and some  $\beta \in A$ ,  $\sum_{i=1}^\infty i\lambda_i(m) \leq n$ ,  $\alpha > \beta$ . Since  $M \subset \cup_{i=1}^\infty \lambda_i^{-1}(1)$ , it follows that  $\{m_\alpha | \alpha > \beta\} \subset \lambda_1^{-1}(1) \cup \dots \cup \lambda_n^{-1}(1)$ . Thus, for some cofinal  $D \subset A$  and some  $k$ ,  $\{m_\alpha | \alpha \in D\} \subset \lambda_k^{-1}(1)$ . But on  $\lambda_k^{-1}(1)$ ,  $f_{2k} = \phi_k$  is a closed embedding into  $\mathbb{R}^\infty$ . Thus, since  $f_{2k}(m_\alpha) = \phi_k(m_\alpha) \rightarrow y_{2k}$  we have that  $y_{2k} = \phi_k(m)$  some  $m \in A$  and that  $\{m_\alpha | \alpha \in D\} \rightarrow m$ . Thus,  $\{f(m_\alpha) | \alpha \in D\} \rightarrow f(m)$  so that  $y = f(m) \in f(A)$ . We have shown that  $f(A)$  is closed, and it follows that  $f$  is a closed topological embedding.

Let  $N = f(M)$ . To see that  $N$  is a p.l.  $\mathbb{R}^\infty$ -submanifold of  $(\mathbb{R}^\infty)^\infty$  let  $m_0 \in M$ . Find  $j$  such that  $m_0 \in W_j$  and then a neighborhood  $O$  of  $m_0$  such that  $O \subset W_j$ . Then on  $O$ ,  $f_{2j} = \phi_j$ . Let  $Z = (\mathbb{R}^\infty \times \mathbb{R}^\infty \times \dots \times \mathbb{R}^\infty \times \phi_j(O) \times \mathbb{R}^\infty \times \mathbb{R}^\infty \times \dots) \cap (\mathbb{R}^\infty)^\infty$ , where  $\phi_j(O)$  occurs in the  $2j$  factor. Define  $\gamma: Z \rightarrow Z$  by  $\gamma((x_i)) = (y_i)$  where  $y_i = x_i - f_i \phi_j^{-1}(x_{2j})$ ,  $i \neq 2j$ , and  $y_{2j} = x_{2j}$ . Then  $\gamma$  is an  $\mathbb{R}^\infty$ -p.l. isomorphism, and  $\gamma(Z \cap f(M)) = \mathbf{0} \times \mathbf{0} \times \dots \times \mathbf{0} \times \phi_j(O) \times \mathbf{0} \times \mathbf{0} \dots$ . Define  $\delta: Z \rightarrow \phi_j(O) \times \mathbb{R}^\infty$  by  $\delta((x_i)) = (x_{2j}, \rho^{-1}(x_1, \dots, x_{2j-1}, x_{2j+1}, x_{2j+2}, \dots))$

where  $\rho: \mathbb{R}^\infty \rightarrow (\mathbb{R}^\infty)^\infty$  is the homeomorphism given at the beginning of this proof. Then  $\delta$  is a p.l.  $\mathbb{R}^\infty$ -isomorphism. Thus,  $(Z, \delta\gamma)$  is a p.l.  $\mathbb{R}^\infty$ -chart for  $(\mathbb{R}^\infty)^\infty$  with  $f(m_0) \in Z$ ,  $\gamma\delta(Z) = \phi_j(0) \times \mathbb{R}^\infty$ , and  $\delta\gamma(Z \cap f(M)) = \phi_j(0) \times \{0\}$ . Thus,  $N = f(M)$  is an  $\mathbb{R}^\infty$ -p.l. submanifold of  $(\mathbb{R}^\infty)^\infty$ .

We thus have an  $\mathbb{R}^\infty$ -p.l. map  $f: M \rightarrow (\mathbb{R}^\infty)^\infty$  which is a topological embedding onto a closed p.l.  $\mathbb{R}^\infty$ -submanifold of  $(\mathbb{R}^\infty)^\infty$ . While it is relatively easy to see directly that  $f^{-1}$  is  $\mathbb{R}^\infty$ -p.l., for this we refer, instead, to the remark following Proposition 5. Thus,  $f$  is the desired  $\mathbb{R}^\infty$ -p.l. isomorphism.

#### IV. $\mathbb{R}^\infty$ -Polyhedra

In this section we show that any p.l.  $\mathbb{R}^\infty$ -submanifold of  $\mathbb{R}^\infty$  is an  $\mathbb{R}^\infty$ -polyhedron (see definition in §I) and relate the two definitions of  $\mathbb{R}^\infty$ -p.l. maps on such spaces.

*Lemma 6.* Let  $M$  be a p.l.  $\mathbb{R}^\infty$ -manifold. Let  $C \subset M$  be compact. Then any p.l.  $\mathbb{R}^\infty$ -atlas for  $M$  contains finitely many p.l.  $\mathbb{R}^\infty$ -charts  $\{(U_i, \phi_i) \mid i = 1, \dots, n\}$  such that there are cubes  $D_1, \dots, D_n$  in  $\mathbb{R}^{k_1}, \dots, \mathbb{R}^{k_n}$ , respectively, with  $D_i \subset \phi_i(U_i)$  and  $C \subset \bigcup_{i=1}^n \phi_i^{-1}(\text{Int}_{\mathbb{R}^{k_i}}(D_i))$ .

*Proof.* Let  $c \in C$ . Let  $(U, \phi)$  be a p.l.  $\mathbb{R}^\infty$ -chart from the given atlas. Choose an open set  $V$  such that  $c \in V \subset \bar{V} \subset U$ . Then compact  $\phi(C \cap \bar{V}) \subset \mathbb{R}^n$  some  $n$ . Let  $y = (y_i) = \phi(c)$  and choose  $D' = (\prod_{i=1}^n [y_i - \epsilon_i, y_i + \epsilon_i]) \cap \mathbb{R}^\infty$  such that  $D' \subset \phi(U)$ . Then, if  $D = \prod_{i=1}^n [y_i - \epsilon_i, y_i + \epsilon_i]$ ,  $\phi^{-1}(\text{Int}_{\mathbb{R}^n} D) \supset \phi^{-1}((\prod_{i=1}^n (y_i - \epsilon_i, y_i + \epsilon_i)) \cap \phi(C \cap V))$ , a neighborhood of  $c$  in  $C \cap V$  and, hence, also in  $C$ . The lemma now follows from

the compactness of  $C$ .

*Proposition 7.* If  $M$  is a closed p.l.  $\mathbb{R}^\infty$ -submanifold of  $\mathbb{R}^\infty$ , then  $M$  is an  $\mathbb{R}^\infty$ -polyhedron.

*Proof.* Let  $C \subset \mathbb{R}^\infty$  be a compact polyhedron. From the definition of submanifold for each  $x \in M$  there is a p.l.  $\mathbb{R}^\infty$ -chart  $(U, \phi)$  with  $x \in U$  and such that  $\phi(U) = U_1 \times U_2$ ,  $\phi(U \cap M) = U_1 \times \{0\} \equiv U_1$ . The corresponding charts  $(U', \phi') = (U \cap M, \phi|_{(U \cap M)})$  form a p.l.  $\mathbb{R}^\infty$ -atlas for  $M$ . By the preceding lemma there are finitely many such charts  $\{(U'_i, \phi'_i) \mid i = 1, \dots, n\}$  and compact cubes  $D_i \subset \phi'_i(U'_i)$  such that  $C \cap M \subset \bigcup_{i=1}^n (\phi'_i)^{-1}(D_i)$ . Thus,  $C \cap M = C \cap \bigcup_{i=1}^n (\phi'_i)^{-1}(D_i)$ . Since finite unions and intersections of compact polyhedra are again polyhedra it suffices to show that each  $(\phi'_i)^{-1}(D_i)$  is a polyhedron in  $\mathbb{R}^\infty$ . But  $(\phi'_i)^{-1}(D_i) = \phi_i^{-1}(D_i \times 0)$ , a compact polyhedron since  $\phi_i^{-1}: \phi_i(U) \rightarrow U$  is an  $\mathbb{R}^\infty$ -p.l. isomorphism.

*Proposition 8.* If  $M$  and  $N$  are closed p.l.  $\mathbb{R}^\infty$ -submanifolds of  $\mathbb{R}^\infty$  then  $f: M \rightarrow N$  is  $\mathbb{R}^\infty$ -p.l. in the manifold sense if and only if  $f$  is  $\mathbb{R}^\infty$ -p.l. in the polyhedral sense.

*Proof.* First note that if  $(U, \phi)$  is any p.l.  $\mathbb{R}^\infty$ -chart in  $\mathbb{R}^\infty$ , and if  $Q \subset U$  is a compact polyhedron then  $\phi|_Q = (\text{id}_{\mathbb{R}^\infty})^{-1} \phi|_Q$  is p.l. and, hence,  $\phi(Q)$  is a compact polyhedron.

Now let  $f$  be  $\mathbb{R}^\infty$ -p.l. in the manifold sense. Let  $C \subset M$  be a compact polyhedron,  $x \in C$ . Let  $(U, \phi), (V, \psi)$  be charts for  $M$  and  $N$ , respectively, with  $x \in U, f(x) \in V$  and such that  $(U', \phi') = (U \cap M, \phi|_{(U \cap M)})$  and  $(V', \psi') = (V \cap N, \psi|_{(V \cap N)})$  are p.l.  $\mathbb{R}^\infty$ -charts for  $M$  and  $N$ . Choose a compact polyhedral neighborhood  $P$  of  $x$  in  $C \cap U' \cap f^{-1}(V')$ . Then

$\phi'(P) = \phi(P)$  is a compact polyhedron, and, since  $f$  is  $R^\infty$ -p.l. in manifold sense,  $\psi'f(\phi')^{-1}|_{\phi(P)}$  is p.l. Thus,  $f|_P = \psi^{-1}(\psi'f(\phi')^{-1})\phi|_P$  is p.l. Thus,  $f|_C$  is locally p.l. and, hence, p.l. as required.

Conversely, if  $f: M \rightarrow N$  is  $R^\infty$ -p.l. in the polyhedral sense and  $x \in M$ , let  $(U, \phi)$ ,  $(V, \psi)$ ,  $(U', \phi')$ ,  $(V', \psi')$  be as above. If  $P \subset \phi'(U' \cap f^{-1}(V'))$  is a compact polyhedron then  $(\phi')^{-1}(P)$  is a compact polyhedron in  $M$ . Thus  $f|_{(\phi')^{-1}(P)}$  is p.l. Hence  $\psi'f(\phi')^{-1}|_P$  is p.l. as required.

### References

1. R. E. Heisey, *Contracting spaces of maps on the countable direct limit of a space*, Trans. Amer. Math. Soc. 193 (1974), 389-411.
2. \_\_\_\_\_, *Partitions of unity and a closed embedding theorem for  $(C, b^*)$ -manifolds*, Trans. Amer. Math. Soc. 206 (1975), 281-294.
3. C. P. Rourke and B. J. Sanderson, *Introduction to piecewise-linear topology*, Ergebnisse Math. Grenzgebiete, Band 69, Springer-Verlag, New York, 1972. MR. #3236.

Vanderbilt University

Nashville, Tennessee 37235