
TOPOLOGY PROCEEDINGS



Volume 11, 1986

Pages 25–27

<http://topology.auburn.edu/tp/>

A LOCALLY COMPACT, HOMOGENEOUS METRIC SPACE WHICH IS NOT BIHOMOGENEOUS

by
H. COOK

Topology Proceedings

Web: <http://topology.auburn.edu/tp/>

Mail: Topology Proceedings
Department of Mathematics & Statistics
Auburn University, Alabama 36849, USA

E-mail: topolog@auburn.edu

ISSN: 0146-4124

COPYRIGHT © by Topology Proceedings. All rights reserved.

A LOCALLY COMPACT, HOMOGENEOUS METRIC SPACE WHICH IS NOT BIHOMOGENEOUS

H. Cook

Suppose S is a topological space. If, for each two points a and b of S , there is a homeomorphism of S onto S throwing a to b , then S is said to be homogeneous. If, for each two points a and b of S , there is a homeomorphism of S onto S throwing a to b and b to a , then S is said to be bihomogeneous. In [2], Kuratowski gave an example of a connected metric homogeneous space which is not bihomogeneous. His example is not locally compact (nor, indeed, is it topologically complete). Here, we give a locally compact example.

The Space X . Let P denote a pseudo-arc and $C(P)$ denote the space of all subcontinua of P . Let X denote the subspace of $C(P)$ comprising the nondegenerate proper subcontinua of P .

Theorem 0. *The space X is connected and locally compact.*

Proof. There is a continuous collection H of proper subcontinua of P filling up P , [1]. Now, H is a continuum in X and each point of X is connected to H by an arc, [1]. Thus, X is connected.

Since X is an open subset of the compact space $C(P)$, X is locally compact.

Theorem 1. The space X is homogeneous.

Proof. Lehner has shown [4] that, if a and b are two proper subcontinua of P , then there is a homeomorphism h of P onto P that takes a onto b . Then h^* , the homeomorphism of $C(P)$ onto $C(P)$ induced by h , throws X onto X and $h^*(a) = b$. Thus, X is homogeneous.

Theorem 2. The space X is not bihomogeneous.

Proof. Let a and b be two points of X such that b is a proper subcontinuum of a . Further, let b_1, b_2, b_3, \dots be a sequence of proper subcontinua of a (points of X) converging to b such that no two of them lie in the same composant of a and no one of them lies in the composant of a containing b . There is only one arc ab in X from a to b ; for each i , there is only one arc ab_i from a to b_i ; and if tb is an arc in X from a point $t \neq b$ of X to b such that $tb \cap ab = b$ then t is a proper subcontinuum of b (in P), [1].

Now, suppose that h is a homeomorphism from X onto X such that $h(a) = b$ and $h(b) = a$. For each i , let $t_i = h(b_i)$. Now, h throws the arc ab onto itself, and, for each i , throws the arc ab_i onto an arc $t_i b$ such that $t_i b \cap ab = b$. Then, for each i , t_i is a proper subcontinuum (in P) of b . But the sequence t_1, t_2, t_3, \dots converges to $h(b) = a$. But no sequence of proper subcontinua of b converges to a .

Thus, X is not bihomogeneous.

Notes. For a general exposition of the pseudo-arc, see [3]. For a thorough discussion of hyperspaces such as $C(P)$, see [5].

Kuratowski's example, mentioned in the opening paragraph, is a 1-dimensional subspace of the plane. The space X is a 2-dimensional subspace of Euclidean 3-space, [6]. K. Kuperberg tells me that she has an example (to appear) of a 3-dimensional compact metric continuum which is homogeneous but not bihomogeneous.

References

- [1] J. L. Kelley, *Hyperspaces of a continuum*, Trans. Amer. Math. Soc. 52 (1942), 22-36.
- [2] K. Kuratowski, *Un probleme sur les ensembles homogenes*, Fund. Math. 3 (1922), 14-19.
- [3] _____, *Topology*, Vol. II, Academic Press, New York and London, 1968.
- [4] G. R. Lehner, *Extending homeomorphisms on the pseudo-arc*, Trans. Amer. Math. Soc. 98 (1961), 369-394.
- [5] S. B. Nadler, Jr., *Hyperspaces of sets*, Marcel Dekker, Inc., New York, 1978.
- [6] W. R. R. Transue, *On the hyperspace of continua of the pseudo-arc*, Proc. Amer. Math. Soc. 18 (1967), 1074-1075.

University of Houston

Houston, Texas 77004