

TOPOLOGY PROCEEDINGS



Volume 12, 1987

Pages 111–116

<http://topology.auburn.edu/tp/>

ON THE SUBGROUPS OF THE FUNDAMENTAL GROUP AND THE REPRESENTATIONS

by

JINGYAL PAK

Topology Proceedings

Web: <http://topology.auburn.edu/tp/>

Mail: Topology Proceedings
Department of Mathematics & Statistics
Auburn University, Alabama 36849, USA

E-mail: topolog@auburn.edu

ISSN: 0146-4124

COPYRIGHT © by Topology Proceedings. All rights reserved.

ON THE SUBGROUPS OF THE FUNDAMENTAL GROUP AND THE REPRESENTATIONS

Jingyal Pak

1. Introduction

Let X be a compact, connected ANR. Let $E(X)$ and $H(X)$ be the space of self-homotopy equivalences and the group of homeomorphisms of X , respectively. Let $\gamma: E(X) \rightarrow X$ and $\gamma': H(X) \rightarrow X$ be the evaluation maps at $x \in X$. Then γ and γ' induce $\gamma_{\#}: \pi_1(E(X), id) \rightarrow \pi_1(X, x)$ and $\gamma'_{\#}: \pi_1(H(X), id) \rightarrow \pi_1(X, x)$ such that if $i: H(X) \rightarrow E(X)$ denotes the inclusion map, then we have the following commutative diagram

$$\begin{array}{ccc} \pi_1(H(X), id) & \xrightarrow{i_{\#}} & \pi_1(E(X), id) \\ \gamma'_{\#} \searrow & & \swarrow \gamma_{\#} \\ & \pi_1(X, x) & \end{array}$$

McCarty [6] has shown that $\gamma'_{\#}(\pi_1(H(X), id))$ lies in the center $Z(\pi_1(X, x))$ and each element $\alpha \in \gamma'_{\#}(\pi_1(H(X), id))$ acts trivially on $\pi_k(X, x)$ for all $k \geq 1$ if X is an admissible space, i.e., X is at least locally compact, locally connected and Hausdorff. Therefore, the natural question is whether $\alpha \in \gamma'_{\#}(\pi_1(H(X), id))$ if any $\alpha \in \pi_1(X, x)$ acts trivially on $\pi_k(X, x)$, for all $k \geq 1$. This question arose while I was studying the Nielsen fixed point theorems, which heavily depend on the structure of the fundamental group [1].

All those elements $\alpha \in \pi_1(X, x)$ which act trivially on $\pi_k(X, x)$ are called k -simple elements of $\pi_1(X, x)$ [4]. Then our question can be rephrased as: if $\alpha \in \pi_1(X, x)$ k -simple for all $k \geq 1$, then is $\alpha \in \gamma'_{\#}(\pi_1(H(X), id))$?

Let $P(X, x) = \{\alpha \in \pi_1(X, x) \mid \alpha \text{ is } k\text{-simple for all } k \geq 1\}$. Then $P(X, x) \subset Z(\pi_1(X, x))$, the center of $\pi_1(X, x)$. In 1965, Gottlieb [3] showed that if X is aspherical, then $\gamma_{\#}(\pi_1(E(X), id)) \simeq P(X, x) \simeq Z(\pi_1(X, x))$. Now our question can be rephrased as follows: if X is aspherical then under what conditions $\gamma'_{\#}(\pi_1(H(X), id)) \simeq P(X, x) \simeq Z(\pi_1(X, x))$, that is, whether $\gamma'_{\#}$ hits the center.

Let $E(X, x)$ and $H(X, x)$ be the based self-homotopy equivalences and the based homeomorphisms at $x \in X$. For any homeomorphism $g \in H(X, x)$, we have an induced automorphism $g_{\#}: \pi_k(X, x) \rightarrow \pi_k(X, x)$. If g is based homotopic to g' then the induced automorphisms agree, i.e., $g_{\#} = g'_{\#}$, and yield a representation

$$\psi: \pi_0(H(X, x), id) \rightarrow \text{Aut}(\pi_k(X, x)).$$

We answer the above question in the following form. Let $X = M$ be a closed, connected aspherical manifold. Then $\gamma'_{\#}$ hits the center if and only if the representation $\psi: \pi_0(H(M, x), id) \rightarrow \text{Aut}(\pi_1(M, x))$ is faithful. This implies that if $\pi_1(M, x)$ is centerless then the representation ψ is faithful. At the end we will give some examples satisfying our hypothesis.

Finally I would like to thank Gottlieb for his comments made on the original version of this paper.

2. On the Homeomorphism Groups and Representations

Let $X = M$ be a closed, connected aspherical manifold. A manifold M is called aspherical if its universal covering space $\tilde{M}$ is contractible, i.e., M is a $K(\pi, 1)$ -space. As before, let $E(M)$ and $H(M)$ be the self-homotopy equivalences

and the based homeomorphisms at $x \in M$ respectively. The evaluation maps $\gamma: E(M) \rightarrow M, \gamma': H(M) \rightarrow M$ defined by $\gamma(h) = h(x)$ and $\gamma'(g) = g(x)$ at $x \in M$ are fiberings [2], [6], and we have the following fiber-homotopy commutative diagram:

$$\begin{array}{ccccccc} \cdots \rightarrow \pi_1(H(M), id) & \xrightarrow{\gamma'_\#} & \pi_1(M, x) & \xrightarrow{d'_\#} & \pi_0(H(M, x), id) & \rightarrow & \pi_0(H(M), id) \rightarrow 0 \\ & \downarrow & \downarrow & & \downarrow & & \downarrow \\ \cdots \rightarrow \pi_1(E(M), id) & \xrightarrow{\gamma_\#} & \pi_1(M, x) & \xrightarrow{d_\#} & \pi_0(E(M, x), id) & \rightarrow & \pi_0(E(M), id) \rightarrow 0 \end{array}$$

Lemma 1. Let M be a closed, connected aspherical manifold. If the induced homomorphism $0i_\#: \pi_0(H(M, x), id) \rightarrow \pi_0(E(M, x), id)$ is a monomorphism then $1i_\#: \pi_1(H(M), id) \rightarrow \pi_1(E(M), id)$ is an epimorphism.

Proof. From the Gottlieb theorem [3], we know $\pi_1(H(M, x), id) = 0$ and $\pi_1(E(M, x), id) = 0$, and we have the following commutative diagram:

$$\begin{array}{ccccccc} 0 \rightarrow \pi_1(H(M), id) & \rightarrow & \pi_1(M, x) & \rightarrow & \pi_0(H(M, x), id) & \rightarrow & \pi_1(H(M), id) \rightarrow 0 \\ & 1i_\# \downarrow & & \downarrow \parallel & 0i_\# \downarrow & & \downarrow \\ 0 \rightarrow \pi_1(E(M), id) & \rightarrow & \pi_1(M, x) & \rightarrow & \pi_0(E(M, x), id) & \rightarrow & \pi_0(E(M), id) \rightarrow 0 \end{array}$$

We can see that $1i_\#$ is a monomorphism, since it factors through

$$\begin{array}{ccc} 0 \rightarrow \pi_1(H(M), id) & \xrightarrow{r'_\#} & \pi_1(M, x) \\ & 1i_\# \downarrow & \uparrow r_\# \\ & \pi_1(E(M), id) & \end{array}$$

Now by diagram chasing, i.e., by the Weak Four Lemma [7], we know that $1i_\#$ onto if $0i_\#$ is a monomorphism.

Corollary 2. $\gamma'_\#(\pi_1(H(M), id)) \simeq Z(\pi_1(M, x))$ if $0i_\#$ is a monomorphism.

Proof. Since $\gamma_{\#}(E(M), id) \simeq Z(\pi_1(M, x))$ [3] and $1i_{\#}$ is onto from the lemma 1, we have the result.

Lemma 3. Let M be a closed, connected aspherical manifold such that $\pi_0(H(M), id) \rightarrow \pi_0(E(M), id)$ is a monomorphism. If $1i_{\#}$ is an epimorphism then $0i_{\#}: \pi_0(H(M, x), id) \rightarrow \pi_0(E(M, x), id)$ is a monomorphism.

Proof. This lemma again follows from diagram chasing. This time we apply The Five Lemma [7]. Note that $0i_{\#}$ is an epimorphism if $\pi_0(H(M), id) \rightarrow \pi_0(E(M), id)$ is onto.

Corollary 4. With the hypothesis of lemma 3, we have a representation $\psi: \pi_0(H(M, x), id) \rightarrow \text{Aut}(\pi_1(M, x))$, which is faithful.

Proof. Let $\psi = 0i_{\#} \phi$, where $\phi: \pi_0(E(M, x), id) \rightarrow \text{Aut}(\pi_1(M, x))$ is an isomorphic representation [2]. Since $0i_{\#}$ becomes monic, the result follows.

Combining these two lemmas, we have

Theorem 5. Let M be a closed, connected aspherical manifold such that $\pi_0(H(M), id) \simeq \pi_0(E(M), id)$. Then $1i_{\#}$ is an isomorphism if and only if $0i_{\#}$ is an isomorphism i.e., there is an isomorphic representation $\psi: \pi_0(H(M, x), id) \rightarrow \text{Aut}(\pi_1(M, x))$.

Corollary 6. Let M be a closed, connected aspherical manifold. If $\pi_1(M, x)$ has no non-trivial center, then there is a faithful representation. $\psi: \pi_0(H(M, x), id) \rightarrow \text{Aut}(\pi_1(M, x))$.

Remark. Let M be an arbitrary manifold. If $[\alpha] \in \pi_1(M, x)$ then we can lift the loop α to α^* in $H(M)$ such that α^* is a path from the identity homeomorphism to $g \in H(M, x)$. McCarty [6] has shown that the induced automorphism $g_{\#}: \pi_k(M, x) \rightarrow \pi_k(M, x)$ is the same as the standard action of $[\alpha]$ on higher homotopy groups for all k [4]. Thus if $[\alpha] \in P(M, x)$, then $g_{\#}$ becomes the identity automorphism for all $k \geq 1$. If this implies that g is isotopic to the identity homeomorphism relative to x on M then g belongs to the identity path component in $\pi_0(H(M, x))$. Let β be a path from g to identity map in $H(M, x)$. Then $\beta \circ \alpha^*$ is a loop in $H(X)$ such that $\gamma'(\beta \circ \alpha^*) = \alpha$. This implies $\gamma'(\pi_1(H(M), id) = P(M, x)$. We ask the following question. If $g_{\#}: \pi_k(M, x) \rightarrow \pi_k(M, x)$ is the identity automorphism for all $k \geq 1$, what conditions are necessary to ensure that g belongs to the path component of the identity homeomorphism in $H(M, x)$.

Example 1. Let M be a closed, connected 3-manifold which is irreducible and sufficiently large [8]. These manifolds are aspherical and Waldhausen has shown $\pi_0(E(M)) \cong \pi_0(H(M))$. On the other hand Laudenbach [5], pushing further Waldhausen's result, has shown $\pi_1(E(M), id) \cong \pi_1(H(M), id)$ if in addition M is P^2 -irreducible. Thus these manifolds satisfy our hypothesis, and there is an isomorphic representation

$$\psi: \pi_0(H(M, x), id) \rightarrow \text{Aut } \pi_1(M, x).$$

Example 2. For the higher dimensional examples, we show "The model aspherical manifolds" from [2]. Let (W, N) be a properly discontinuous action of a discrete group N on a contractible manifold W so that W/N is compact. Then for each torsion free extension $1 \rightarrow Z^k \rightarrow \pi \rightarrow N \rightarrow 1$ the space $M = (T^k \times W)/N$ is an aspherical manifold and the map $M \rightarrow W/N$ is a Seifert fibering. These manifolds M satisfy our hypothesis; $\pi_0(E(M)) \simeq \pi_0(H(M))$ and $\pi_1(E(M), id) \simeq \pi_1(H(M), id)$. Thus we have an isomorphic representation $\psi: \pi_0(H(M, x), id) \rightarrow \text{Aut}(\pi_1(M, x))$.

References

1. R. F. Brown, *The Lefschetz fixed point theorem*, Scott Foresman and Co., Glenview, Ill., 1971.
2. P. E. Conner and F. Raymond, *Deforming homotopy equivalences to homeomorphisms in aspherical manifolds*, Bulletin of A.M.S. 83 (1977), 36-85.
3. D. H. Gottlieb, *A certain subgroup of the fundamental group*, Amer. J. Math. 87 (1965), 840-856.
4. S. T. Hu, *Homotopy theory*, Academic Press, New York, 1959.
5. F. Laudenbach, *Topologie de la dimension trois: homotopie et isotopie*, Astérisque, Soc. Math. de Franc 12 (1974).
6. G. S. McCarty, Jr., *Homeotopy groups*, Trans. of A.M.S. 106 (1963), 293-304.
7. S. MacLane, *Homology*, Academic Press, New York, 1963.
8. F. Waldhausen, *On irreducible 3-manifolds which are sufficiently large*, Ann. of Math. 18 (1968), 56-88.

Wayne State University
 Detroit Michigan 48202