


<http://topology.auburn.edu/tp/>

TOPOLOGY PROCEEDINGS



Volume 53, 2019

Pages 201–207

<http://topology.nipissingu.ca/tp/>

COARSE DIRECT PRODUCTS AND PROPERTY C

by

G. BELL AND A. LAWSON

Electronically published on January 18, 2019

Topology Proceedings

Web: <http://topology.auburn.edu/tp/>

Mail: Topology Proceedings
Department of Mathematics & Statistics
Auburn University, Alabama 36849, USA

E-mail: topolog@auburn.edu

ISSN: (Online) 2331-1290, (Print) 0146-4124

COPYRIGHT © by Topology Proceedings. All rights reserved.



COARSE DIRECT PRODUCTS AND PROPERTY C

G. BELL AND A. LAWSON

ABSTRACT. We show that coarse property C is preserved by finite coarse direct products. We also show that the coarse analog of Dydak's countable asymptotic dimension is equivalent to the coarse version of straight finite decomposition complexity and is therefore preserved by direct products.

1. INTRODUCTION AND PRELIMINARIES

The coarse category was described by Roe [9] as a generalization of the large-scale approach to discrete groups begun by Gromov [7]. Coarse spaces are sets that are equipped with a so-called coarse structure that provides a measure of proximity without referring to a metric. Coarse structures can be derived from metric structures [9, 11], topological structures [9], or group structures [8]. Coarse versions of asymptotic dimension [6, 9] as well as property C and finite decomposition complexity [1] have been established and studied [12].

The primary goal of this short note is to show that coarse property C is stable with respect to finite coarse direct products (defined below); this was shown in the metric case recently [2, 3]. We also show that the coarse analog of Dydak's countable asymptotic dimension [4] coincides with the coarse version of straight finite decomposition complexity (sFCDC); as a result, this notion is also stable with respect to coarse direct products.

A coarse structure can be defined on any set X . Take the multiplication (referred to here as composition) and inverse operations from the pair groupoid structure on the product $X \times X$. A collection \mathcal{E} of subsets of $X \times X$ is called a **coarse structure** on X if it contains the diagonal and is closed under subsets, finite unions, inverses, and compositions [9].

2010 *Mathematics Subject Classification.* Primary 54F45; Secondary 54E15.

Key words and phrases. Coarse geometry, property C.

©2019 Topology Proceedings.

Elements of \mathcal{E} will be called **entourages** and we call the pair (X, \mathcal{E}) a **coarse space**.

Suppose (X_1, \mathcal{E}_1) and (X_2, \mathcal{E}_2) are coarse spaces. Let p_i denote the projection map $p_i : X_1 \times X_2 \rightarrow X_i$ for $i \in \{1, 2\}$. We define the **(direct) product coarse structure** [6] on the product $X_1 \times X_2$ by

$$\mathcal{E}_1 * \mathcal{E}_2 = \{E \subseteq (X_1 \times X_2)^2 : (p_i \times p_i)(E) \in \mathcal{E}_i \text{ for each } i \in \{1, 2\}\}.$$

If X and Y are metric spaces in the bounded coarse structure [9, Example 2.5], the coarse space obtained from taking the bounded coarse structure on $X \times Y$ is the same as the coarse product of the spaces X and Y equipped with bounded coarse structures; by contrast, the C_0 -coarse structure (defined by Wright [11]) is not necessarily preserved in this way. For example, the C_0 -coarse structure on the metric product $\mathbb{N} \times [0, 1]$ is easily seen to be a proper subset of the coarse product of the C_0 -coarse structures on the individual spaces. Therefore, a coarse structure on the product of two spaces does not necessarily correspond to the coarse product of those same types of coarse structure on the factors.

To complete this section, we recall the definitions of coarse property **C** and sFCDC [1].

Definition 1.1. [1] A coarse space (X, \mathcal{E}) has **coarse property C** if and only if for any sequence $E_1 \subseteq E_2 \subseteq \dots$ of entourages there is a finite sequence of families $\mathcal{U}_1, \mathcal{U}_2, \dots, \mathcal{U}_n$ of subsets of X satisfying

- (1) $\bigcup_{i=1}^n \mathcal{U}_i$ is a cover of X ;
- (2) each \mathcal{U}_i is **E_i -disjoint**; i.e., for any pair of distinct elements U, V of \mathcal{U}_i we have $U \times V \cap E_i = \emptyset$; and
- (3) each \mathcal{U}_i is **uniformly bounded**; i.e., $\bigcup_{U \in \mathcal{U}_i} U \times U \in \mathcal{E}$.

Let Y be a subset of a coarse space (X, \mathcal{E}) and let \mathcal{U} be a family of subsets of X . Let n be a positive integer and let $E \in \mathcal{E}$ be an entourage. We say that Y admits an (E, n) -**decomposition** over \mathcal{U} if Y can be expressed as a union of n sets Y^1, Y^2, \dots, Y^n in such a way that each Y^i can be expressed as an E -disjoint union of sets from \mathcal{U} . Here, by an **E -disjoint union of sets from \mathcal{U}** , we mean that each $Y^i = \sqcup_j Y_j^i$, where $Y_j^i \times Y_{j'}^i \cap E = \emptyset$ if $j \neq j'$ and $Y_j^i \in \mathcal{U}$ for all j .

Definition 1.2. ([1]) The coarse space (X, \mathcal{E}) is said to have **straight finite coarse decomposition complexity** (sFCDC) if for any infinite sequence of entourages L_1, L_2, \dots there is a finite sequence of families $\mathcal{V}_1, \dots, \mathcal{V}_n$ of subsets of X so that

- (1) $\mathcal{V}_1 = \{X\}$;
- (2) for every $i \geq 1$, each $U \in \mathcal{V}_i$ admits an $(L_i, 2)$ -decomposition over \mathcal{V}_{i+1} ; and
- (3) \mathcal{V}_n is uniformly bounded.

K_{10}	K_{14}	K_{19}	K_{25}	K_{32}	K_{40}	$K_{1,4}$	$K_{2,4}$	$K_{3,4}$	$K_{4,4}$	$K_{5,4}$	$K_{6,4}$
K_6	K_9	K_{13}	K_{18}	K_{24}	K_{31}	$K_{1,3}$	$K_{2,3}$	$K_{3,3}$	$K_{4,3}$	$K_{5,3}$	$K_{6,3}$
K_3	K_5	K_8	K_{12}	K_{17}	K_{23}	$K_{1,2}$	$K_{2,2}$	$K_{3,2}$	$K_{4,2}$	$K_{5,2}$	$K_{6,2}$
K_1	K_2	K_4	K_7	K_{11}	K_{16}	$K_{1,1}$	$K_{2,1}$	$K_{3,1}$	$K_{4,1}$	$K_{5,1}$	$K_{6,1}$

FIGURE 1. We give a possible rearrangement of the sequence K_1, K_2, \dots into a two-dimensional array $K_{i,j}$.

It is known that sFCDC is preserved by coarse direct products [1, Theorem 4.17].

2. RESULTS

The proof that asymptotic property C is preserved by direct products of metric spaces [2, 3] is based on the technique used to prove the corresponding theorem for topological property C when one of the two factors is compact [10]. The proof of the following theorem is similar in spirit to the earlier works, but the absence of a metric means that more care and bookkeeping is required. In particular the single distance parameter must be replaced by two sequences of entourages in the factors.

Theorem 2.1. *Let (X, \mathcal{E}) and (Y, \mathcal{F}) be coarse spaces with coarse property C. Then $(X \times Y, \mathcal{E} * \mathcal{F})$ has coarse property C.*

Proof. Let $E_1 \subseteq E_2 \subseteq \dots$ be a sequence of entourages in $\mathcal{E} * \mathcal{F}$. For each i , put $K_i = (p_1 \times p_1)(E_i)$ and $L_i = (p_2 \times p_2)(E_i)$. Then, by the definition of $\mathcal{E} * \mathcal{F}$, each $K_i \in \mathcal{E}$ and $L_i \in \mathcal{F}$. Observe that since $E_i \subseteq E_{i+1}$, we have $K_i \subseteq K_{i+1}$ and $L_i \subseteq L_{i+1}$. Arrange the indices $1, 2, 3, \dots$ into a two-dimensional array with the property that the indices are increasing from left to right along rows and from bottom to top along columns. In Figure 1 we give one example of such an arrangement, which was first used in the metric proof [2].

For each i , we apply the coarse property C definition to the column $K_{i,1}, K_{i,2}, \dots$ to find an n_i and a cover $\mathcal{U}_{i,1}, \mathcal{U}_{i,2}, \dots, \mathcal{U}_{i,n_i}$ of X by uniformly bounded families of subsets of X with the property that each $\mathcal{U}_{i,j}$ is $K_{i,j}$ -disjoint. Then, consider the sequence $L_{1,n_1}, L_{2,n_2}, \dots$. We may assume that the sequence is increasing by replacing L_{i,n_i} by an entourage that occurs higher in the i -th column, if necessary.

Using this sequence, and the fact that Y has coarse property C, we find a cover of Y by families $\mathcal{V}_1, \mathcal{V}_2, \dots, \mathcal{V}_m$ of subsets of Y that are uniformly bounded with the property that \mathcal{V}_i is L_{i,n_i} -disjoint.

Using the same rearrangement of indices as above, we construct the doubly indexed collection $\{E_{i,j}\}$ of entourages from the given sequence $E_1 \subseteq E_2 \subseteq \dots$. We claim that the family $\mathcal{W}_{i,j} = \{U \times V : U \in \mathcal{U}_{i,j}, V \in \mathcal{V}_i\}$ covers $X \times Y$, consists of uniformly bounded sets, and has the property that $\mathcal{W}_{i,j}$ is $E_{i,j}$ -disjoint. To finish the proof, we simply need to unravel the re-indexing to arrive at the original sequence, which may now include some empty families.

First we check that the collection $\mathcal{W}_{i,j}$ covers $X \times Y$. To this end, let $(x, y) \in X \times Y$ be given. Since the families $\mathcal{V}_1, \mathcal{V}_2, \dots, \mathcal{V}_m$ cover Y , there is a subset V in one such family (say) \mathcal{V}_{i_0} such that $y \in V$ and $V \in \mathcal{V}_{i_0}$. Now, the families $\mathcal{U}_{i_0,1}, \mathcal{U}_{i_0,2}, \dots, \mathcal{U}_{i_0,n_{i_0}}$ cover X , so there is some index (say) j_0 so that the family \mathcal{U}_{i_0,j_0} contains a subset U of X with $x \in U$.

Fix a pair (i, j) and consider the family $\mathcal{W}_{i,j}$. To show that $\mathcal{W}_{i,j}$ is uniformly bounded, we must show that

$$\bigcup_{W \in \mathcal{W}_{i,j}} W \times W$$

is an entourage in the product coarse structure $\mathcal{E} * \mathcal{F}$. By the definition, we need only show the projection of this union to each factor is an entourage in that factor. Since each $W \in \mathcal{W}_{i,j}$ can be expressed as a product $W = U \times V$ with $U \in \mathcal{U}_{i,j}$ and $V \in \mathcal{V}_i$, we observe that,

$$\begin{aligned} (p_1 \times p_1) \left[\bigcup_{W \in \mathcal{W}_{i,j}} W \times W \right] &= (p_1 \times p_1) \left[\bigcup_{W=U \times V} ((U \times V) \times (U \times V)) \right] \\ &= \bigcup_{W=U \times V} [(p_1 \times p_1)((U \times V) \times (U \times V))] \\ &= \bigcup_{W=U \times V} [U \times U] \in \mathcal{E}. \end{aligned}$$

The conclusion for the projection to the second factor is similar.

Finally, we check that $\mathcal{W}_{i,j}$ is $E_{i,j}$ -disjoint. To this end, take distinct $U_1 \times V_1$ and $U_2 \times V_2$ in $\mathcal{W}_{i,j}$. Assume that there were some $(a, c, b, d) \in E_{i,j} \cap ((U_1 \times V_1) \times (U_2 \times V_2))$. Then, in particular, $a \in U_1$, $b \in U_2$, $c \in V_1$ and $d \in V_2$. Thus, $(a, b) \in (p_1 \times p_1)(E_{i,j}) = K_{i,j}$ and $(c, d) \in (p_2 \times p_2)(E_{i,j}) = L_{i,j}$. Since $U_1 \times V_1 \neq U_2 \times V_2$, we either have $U_1 \neq U_2$ or $V_1 \neq V_2$. In the first case, the $K_{i,j}$ -disjointness of $\mathcal{U}_{i,j}$ does not allow (a, b) to be in $K_{i,j}$. In the second case, the fact that \mathcal{V}_i is L_{i,n_i} -disjoint and the fact that $L_{i,j} \subset L_{i,n_i}$ for all $j \leq n_i$ means that (c, d) cannot be in $L_{i,j}$. Thus, there can be no such point (a, c, b, d) . We conclude that the intersection is empty, which is what we needed to show. \square




FIGURE 2. We find covers $\{\mathcal{U}_{i,j}\}_{j=1}^{n_i}$ of X for each column $K_{i,1}, K_{i,2}, \dots$, and then construct a cover $\{\mathcal{V}_i\}_{i=1}^m$ of Y corresponding to L_{i,n_i}

Dydak defined countable asymptotic dimension for metric spaces [4] and it was shown to be equivalent to straight finite decomposition complexity by Dydak and Virk [5]. We show that the analogous result also holds for coarse spaces. As before, some care is needed to work with entourages in the absence of a metric.

Proposition 2.2. *Let (X, \mathcal{E}) be a coarse space. The following are equivalent:*

- (1) *there is a sequence (n_i) of integers such that for every sequence of entourages K_i there is a finite sequence of families $\mathcal{V}_1, \dots, \mathcal{V}_r$ of subsets of X such that $\mathcal{V}_1 = \{X\}$, every $V \in \mathcal{V}_i$ admits an (K_i, n_i) -decomposition over \mathcal{V}_{i+1} and such that \mathcal{V}_r is uniformly bounded.*
- (2) *for every sequence L_i of entourages there is a finite sequence of families $\mathcal{U}_1, \dots, \mathcal{U}_s$ of subsets of X such that $\mathcal{U}_1 = \{X\}$, every $U \in \mathcal{U}_i$ admits a $(L_i, 2)$ -decomposition over \mathcal{U}_{i+1} and such that \mathcal{U}_s is uniformly bounded.*

Proof. Clearly (2) implies (1).

To see the other implication, let (n_i) be the sequence of positive integers satisfying (1) for X . Let L_1, L_2, \dots be a sequence of entourages. By taking unions we may assume $L_i \subseteq L_{i+1}$. Put $K_1 = L_{n_1}$, $K_2 = L_{n_1+n_2}$, and in general, put $K_j = L_{n_1+\dots+n_j}$. Apply (1) with the sequence (K_i) to obtain $\mathcal{V}_1, \mathcal{V}_2, \dots$ such that $\mathcal{V}_1 = \{X\}$ and such that \mathcal{V}_i admits a (K_i, n_i) -decomposition over \mathcal{V}_{i+1} .

We will define a sequence \mathcal{U}_i of families of subsets of X with the property that $\mathcal{U}_1 = \{X\}$ and \mathcal{U}_i admits an $(L_i, 2)$ -decomposition over \mathcal{U}_{i+1} . To begin, we observe that we can write $X = X^1 \cup X^2 \cup \dots \cup X^{n_1}$, with each $X^i = \bigsqcup_{K_1\text{-disj}} X_j^i$, and each $X_j^i \in \mathcal{V}_2$. Therefore, we take $\mathcal{U}_2 = \{X_1^1, X_2^1, X_3^1, \dots\} \cup \{X^2 \cup X^3 \cup \dots \cup X^{n_1}\}$. Then, it is clear that X can be $(K_1, 2)$ -decomposed over \mathcal{U}_2 and since $L_1 \subseteq L_{n_1} = K_1$, we see that there is an $(L_1, 2)$ -decomposition of any set in \mathcal{U}_1 over the family \mathcal{U}_2 . For \mathcal{U}_3 , we take $\{X_1^1, X_2^1, \dots\} \cup \{X_1^2, X_2^2, \dots\} \cup \{X^3 \cup X^4 \cup \dots \cup X^{n_1}\}$; we also observe that any set in \mathcal{U}_2 admits an $(L_2, 2)$ -decomposition over \mathcal{U}_3 since $L_2 \subseteq L_{n_1} = K_1$. Continue to define families this way to obtain $\mathcal{U}_1, \mathcal{U}_2, \dots, \mathcal{U}_{n_1}$ with an $(L_i, 2)$ -decomposition of each \mathcal{U}_i over \mathcal{U}_{i+1} for $1 \leq i < n_1$. We observe that in this way, $\mathcal{U}_{n_1} = \mathcal{V}_2$.

To define \mathcal{U}_j for $j \geq n_1 + 1$, we observe first that we may assume each family \mathcal{V}_i is a partition of X itself, (cf. [5, Corollary 8.3]). We then repeat the above procedure to arrive at $\mathcal{U}_{n_1+n_2} = \mathcal{V}_3$. We repeat this entire process until we arrive at $\mathcal{U}_{n_1+\dots+n_{r-1}} = \mathcal{V}_r$, which is uniformly bounded. \square

A coarse space has sFCDC precisely when it satisfies condition (2) of Proposition 2.2. Condition (1) of Proposition 2.2 is the coarse analog of countable asymptotic dimension. Since sFCDC is preserved by coarse direct products [1, Theorem 4.17], we obtain the following.

Corollary 2.3. *The coarse version of Dydak's countable asymptotic dimension is preserved by coarse direct products.* \square

The authors wish to express their sincere appreciation to the referee for several helpful suggestions.

REFERENCES

- [1] G. Bell, D. Moran, and A. Nagórko. *Coarse property C and decomposition complexity*, Topology Appl. **227** (2017), 30–50.
- [2] G. C. Bell and A. Nagórko. *On the stability of asymptotic property C for products and some group extensions*, Algebr. Geom. Topol. **18**(1) (2018), 221–245.
- [3] T. Davila. *On asymptotic property C*, ArXiv e-prints, Nov. 2016.
- [4] J. Dydak. *Coarse amenability and discreteness*, J. Aust. Math. Soc. **100**(1) (2016), 65–77.
- [5] J. Dydak and Ž. Virk. *Preserving coarse properties*, Rev. Mat. Complut. **29**(1) (2016), 191–206.
- [6] B. Grave. *Coarse geometry and asymptotic dimension*. PhD thesis, Georg-August-Universität Göttingen, 2005.
- [7] M. Gromov. *Asymptotic invariants of infinite groups*. In *Geometric group theory, Vol. 2 (Sussex, 1991)*, volume 182 of *London Math. Soc. Lecture Note Ser.*, pages 1–295. Cambridge Univ. Press, Cambridge, 1993.

- [8] A. Nicas and D. Rosenthal. *Coarse structures on groups*, Topology Appl. **159**(14) (2012), 3215–3228.
- [9] J. Roe. *Lectures on coarse geometry*, volume 31 of *University Lecture Series*. American Mathematical Society, Providence, RI, 2003.
- [10] D. M. Rohm. *Products of infinite-dimensional spaces*, Proc. Amer. Math. Soc. **108**(4) (1990), 1019–1023.
- [11] N. J. Wright. *$C(0)$ coarse geometry*. ProQuest LLC, Ann Arbor, MI, 2002. Thesis (Ph.D.)—The Pennsylvania State University.
- [12] T. Yamauchi. *Straight finite decomposition complexity implies property A for coarse spaces*, Topology Appl. **231** (2017), 329–336.

DEPARTMENT OF MATHEMATICS AND STATISTICS, THE UNIVERSITY OF NORTH
CAROLINA AT GREENSBORO, GREENSBORO, NC 27402, USA

Email address, Bell: gcbell@uncg.edu

Email address, Lawson: azlawson@uncg.edu