


<http://topology.auburn.edu/tp/>

TOPOLOGY PROCEEDINGS



Volume 55, 2020

Pages 123–126

<http://topology.nipissingu.ca/tp/>

A FINITELY GENERATED GROUP THAT DOES NOT SATISFY THE GENERALIZED BURGHELEA CONJECTURE

by

A. DRANISHNIKOV AND M. HULL

Electronically published on May 27, 2019

Topology Proceedings

Web: <http://topology.auburn.edu/tp/>

Mail: Topology Proceedings
Department of Mathematics & Statistics
Auburn University, Alabama 36849, USA

E-mail: topolog@auburn.edu

ISSN: (Online) 2331-1290, (Print) 0146-4124

COPYRIGHT © by Topology Proceedings. All rights reserved.

A FINITELY GENERATED GROUP THAT DOES NOT SATISFY THE GENERALIZED BURGHELEA CONJECTURE

A. DRANISHNIKOV AND M. HULL

ABSTRACT. We construct a finitely generated group that does not satisfy the generalized Burghlelea conjecture.

1. GENERALIZED BURGHELEA CONJECTURE

In [2], Burghlelea gave an explicit formula for the periodic cyclic homology of groups algebras with rational coefficients (and more generally with coefficients in fields of characteristic zero):

$$PHC_*(\mathbb{Q}G) = \bigoplus_{[x] \in \langle G \rangle^{fin}, n \geq 0} H_{2n+*}(N_x, \mathbb{Q}) \oplus \bigoplus_{[x] \in \langle G \rangle^\infty} T_*(x; \mathbb{Q})$$

where the group $T_*(x; \mathbb{Q}) = \lim_{\leftarrow} \{H_{*+2n}(N_x, \mathbb{Q})\}$.

Here G_x denotes the centralizer of x in G , $N_x = G_x / \langle x \rangle$ is the reduced centralizer, $\langle G \rangle^{fin}$ is the set of conjugacy classes of elements of finite order, and $\langle G \rangle^\infty$ the set of conjugacy classes of elements of infinite order. The bonding maps in the inverse sequences are the Gysin homomorphisms $S : H_{m+2}(N_x, \mathbb{Q}) \rightarrow H_m(N_x, \mathbb{Q})$ corresponding to the fibration $B\langle x \rangle \simeq S^1 \rightarrow BG_x \rightarrow BN_x$.

Conjecture 1.1 (Generalized Burghlelea Conjecture). *Let G be a discrete group, then $T_*(x; \mathbb{Q}) = 0$ for all $x \in \langle G \rangle^\infty$.*

2010 *Mathematics Subject Classification.* 20J05, 20F65.

Key words and phrases. Burghlelea conjecture.

©2019 Topology Proceedings.

Burghelea stated the above conjecture for groups G which admit a finite $K(G, 1)$ [2]. In the same paper Burghelea constructed a countable group that does not satisfy the Generalized Burghelea Conjecture. There are still no known counterexamples to the original version of Burgelea's conjecture, and it is known to hold for many classes of groups [3, 4].

The following is our main result.

Theorem 1.2. *There is a finitely generated group G that does not satisfy the Generalized Burghelea Conjecture.*

Our strategy is to show that the countable group constructed by Burghelea can be embedded in a finitely generated group in a way that preserves centralizers. This embedding is based on the theory of small cancellation over relatively hyperbolic groups developed by Osin [7].

Remark 1.3. Shortly after this paper was written the authors became aware that Engel and Marcinkowski also had constructions of finitely generated counterexamples to the generalized Burghelea Conjecture, including a finitely presented counterexample and a counterexample of type F_∞ , using completely different methods. These constructions have since been added to [4].

2. MALNORMAL EMBEDDINGS

For a torsion-free group G hyperbolic relative to a subgroup H , a subgroup S of G is called *suitable* if S contains two infinite order elements f and g which are not conjugate to any elements of H and such that no non-trivial power of f is conjugate to a non-trivial power of g . This is equivalent to [7, Definition 2.2] since G is torsion-free. Indeed, the maximal virtually cyclic subgroups containing f and g respectively are both cyclic, and since no power of f is equal to a power of g these cyclic subgroups must intersect trivially.

The following is a special case of [7, Theorem 2.4].

Theorem 2.1. *Let G be a torsion-free group hyperbolic relative to a subgroup H , let $t \in G$, and let S be a suitable subgroup of G . Then there exists a group $\overline{G}$ and an epimorphism $\gamma: G \rightarrow \overline{G}$ such that:*

- (1) $\gamma|_H$ is injective (so we identify H with its image in $\overline{G}$).
- (2) $\overline{G}$ is hyperbolic relative to H .
- (3) $\gamma(t) \in \gamma(S)$.
- (4) $\gamma(S)$ is a suitable subgroup of $\overline{G}$.
- (5) $\overline{G}$ is torsion-free.

We inductively apply the previous theorem to construct the desired embedding. This can be extracted from the proof of [7, Theorem 2.6], but since it is not explicitly stated there we include the proof below.

Recall that a subgroup H of a group G is called *malnormal* if for all $x \in G \setminus H$, $x^{-1}Hx \cap H = \{1\}$.

Theorem 2.2. *Let H be a torsion-free countable group. Then there exists a finitely generated group Γ which contains H as a malnormal subgroup.*

Proof. Let $H = \{1 = h_0, h_1, h_2, \dots\}$. We inductively define a sequence of quotients as follows: Let $G_0 = H * F$, where $F = F(x, y)$ is the free group on $\{x, y\}$. Then G_0 is torsion-free, hyperbolic relative to H , and F is a suitable subgroup of G_0 . Let $\alpha_0: G_0 \rightarrow G_0$ be the identity map. Suppose now we have constructed a torsion-free group G_i together with an epimorphism $\alpha_i: G_0 \rightarrow G_i$ such that:

- (1) $\alpha_i|_H$ is injective (so we identify H with its image in G_i).
- (2) G_i is hyperbolic relative to H .
- (3) $\alpha_i(h_j) \in \alpha_i(F)$ for all $0 \leq j \leq i$.
- (4) $\alpha_i(F)$ is a suitable subgroup of G_i .
- (5) G_i is torsion-free.

Given such G_i , we can apply Theorem 2.1 to G_i , H , $t = h_{i+1}$, and $S = \alpha_i(F)$. Let $G_{i+1} = \overline{G_i}$ be the quotient provided by Theorem 2.1. Define $\alpha_{i+1} = \gamma \circ \alpha_i$, where γ is the epimorphism given by Theorem 2.1. Then Theorem 2.1 implies that $\alpha_{i+1}: G_0 \rightarrow G_{i+1}$ satisfies conditions (1)–(5).

Let Γ be the direct limit of the sequence $G_0 \rightarrow G_1 \rightarrow G_2 \dots$, that is $\Gamma = G_0 / \bigcup \ker(\alpha_i)$. Let $\beta: G_0 \rightarrow \Gamma$ be the natural quotient map. Note that $\beta|_F$ is surjective by construction. Indeed, G_0 is generated by $H \cup \{x, y\}$ and for each $h_i \in H$, $\alpha_i(h_i) \in \alpha_i(F)$, hence $\beta(h_i) \in \beta(F)$. Thus Γ is generated by $\{\beta(x), \beta(y)\}$.

Now $\beta|_H$ is injective, so H embeds in Γ ; we identify H with its image in Γ . Suppose $x \in \Gamma$ such that $x^{-1}Hx \cap H \neq \{1\}$. Then there exist $g, h \in H \setminus \{1\}$ such that $x^{-1}gxh^{-1} = 1$. Let $\tilde{x} \in G_0$ such that $\beta(\tilde{x}) = x$. Then for some $i \geq 1$, $\tilde{x}^{-1}g\tilde{x}h^{-1} \in \ker \alpha_i$. This means that $\alpha_i(\tilde{x})^{-1}H\alpha_i(\tilde{x}) \cap H \neq \{1\}$. Since G_i is hyperbolic relative to H , H is malnormal in G_i by [7, Lemma 8.3b]. Hence $\alpha_i(\tilde{x}) \in H$, which means that $x = \beta(\tilde{x}) \in H$. Therefore H is malnormal in Γ . $\square$

3. PROOF OF THEOREM 1.2

Proof of Theorem 1.2. We start by reviewing the counterexample constructed by Burghelea. By the Kan-Thurston theorem [1, 5], there exists a group G and a map $t: K(G, 1) \rightarrow \mathbb{C}P^\infty$ which induces an isomorphism on homology and cohomology. Burghelea observes that the group G can be chosen to be torsion-free. The idea behind this observation is that since $\mathbb{C}P^\infty = \bigcup \mathbb{C}P^n$, $K(G, 1)$ can be constructed inductively as a union

of the form $K(G, 1) = \bigcup K(G_i, 1)$, where $K(G_i, 1) = t^{-1}(\mathbb{C}P^n)$ and each $K(G_i, 1)$ is a finite CW-complex (see, for example, the proof in [6] of the Kan-Thurston theorem). Since $G = \bigcup G_i$ and each G_i is torsion-free, G is also torsion-free.

Note that $H_{2n}(\mathbb{C}P^\infty; \mathbb{Q}) \cong \mathbb{Q}$ and the Gysin homomorphism $S: H_{2n+2}(\mathbb{C}P^\infty; \mathbb{Q}) \rightarrow H_{2n}(\mathbb{C}P^\infty; \mathbb{Q})$ for the canonical S^1 -bundle $S^\infty \rightarrow \mathbb{C}P^\infty$ is an isomorphism, hence $\lim_{\leftarrow} \{H_{2n}(\mathbb{C}P^\infty; \mathbb{Q}), S\} \cong \mathbb{Q}$.

Let

$$1 \rightarrow \mathbb{Z} = \langle x \rangle \rightarrow H \rightarrow G \rightarrow 1$$

be the central extension that corresponds to a generator $a \in H^*(G) = \mathbb{Z}[a]$, $\deg(a) = 2$. Note that $H \cong \pi_1(Y)$, where Y is the pull-back of the bundle $S^\infty \rightarrow \mathbb{C}P^\infty$ along t . Hence $N_x = H/\langle x \rangle \cong G$, and $T_0(x; \mathbb{Q}) \cong \lim_{\leftarrow} \{H_{2n}(G, \mathbb{Q})\} \cong \lim_{\leftarrow} \{H_{2n}(\mathbb{C}P^\infty; \mathbb{Q}), S\} \cong \mathbb{Q}$. H is the group constructed by Burghlelea.

We now apply Theorem 2.2 to obtain a malnormal embedding $H \rightarrow \Gamma$ into a finitely generated group Γ , and we identify H with its image in Γ . Since H is malnormal, no elements of $\Gamma \setminus H$ will centralize x . Hence $\Gamma_x = H$ and $N_x \cong G$. Then as before, we get that $T_0(x; \mathbb{Q}) \cong \mathbb{Q} \neq 0$ in the group Γ . $\square$

REFERENCES

- [1] G. Baumslag, E. Dyer and A. Heller, *The topology of discrete groups*, J. Pure Appl. Algebra **16** (1980), 1–47.
- [2] D. Burghlelea, *The cyclic homology of the group rings*, Comment. Math. Helvetici **60** (1985), 354–365.
- [3] A. Dranishnikov, *On Burghlelea Conjecture*, arXiv:1612.08700.
- [4] A. Engel, M. Marcinkowski, *Burghlelea conjecture and asymptotic dimension of groups*, Journal of Topology and Analysis, <https://doi.org/10.1142/S1793525319500559>.
- [5] D. M. Kan and W. P. Thurston, *Every connected space has the homology of a $K(\pi, 1)$* , Topology **15** (1976), 253–258.
- [6] C.R.F. Maunder, *A Short Proof of a Theorem of Kan and Thurston*, Bull. London Math. Soc. (1981) **13** (4), 325–327.
- [7] D. Osin, *Small cancellations over relatively hyperbolic groups and embedding theorems*, Ann. of Math. **172** (2010), no. 1, 1–39.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF FLORIDA, GAINESVILLE, FL 32611, U.S.A.

Email address, Dranishnikov: dranish@ufl.edu

Email address, Hull: mbhull@ufl.edu